

2013

Moments of Matrix Variate Skew Elliptically Contoured Distributions

Shimin Zheng

East Tennessee State University; Nanjing Audit University, zhengs@etsu.edu

Jeff Knisley

East Tennessee State University, knisleyj@etsu.edu

Chunming Zhang

University of Wisconsin

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>



Part of the [Statistics and Probability Commons](#)

Citation Information

Zheng, Shimin; Knisley, Jeff; and Zhang, Chunming. 2013. Moments of Matrix Variate Skew Elliptically Contoured Distributions. *Advances and Applications in Statistics*. Vol.36(1). 13-27. <http://www.pphmj.com/abstract/7903.htm> ISSN: 0972-3617

This Article is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

Moments of Matrix Variate Skew Elliptically Contoured Distributions

Copyright Statement

© Pushpa Publishing House. This document was published with permission by Pushpa Publishing House. It was originally published in *Advances and Applications in Statistics*.



MOMENTS OF MATRIX VARIATE SKEW ELLIPTICALLY CONTOURED DISTRIBUTIONS

Shimin Zheng^{1,2}, Jeff Knisley³ and Chunming Zhang⁴

¹Institute for Quantitative Biology

Department of Biostatistics and Epidemiology

East Tennessee State University

Box 70259, Johnson City, TN 37614, U. S. A.

²Department of Finance

Nanjing Audit University

Nanjing, P. R. China

³Institute for Quantitative Biology

Department of Mathematics and Statistics

East Tennessee State University

Box 70663, Johnson City, TN 37614, U. S. A.

⁴Department of Statistics

University of Wisconsin

1300 University Avenue

Madison, WI 53706, U. S. A.

Abstract

Matrix variate skew elliptically contoured distributions generalize several classes of important distributions. This paper defines and

Received: May 31, 2013; Accepted: June 24, 2013

2010 Mathematics Subject Classification: Primary 62H10; Secondary 62H05.

Keywords and phrases: matrix variate, skew elliptically contoured distribution, skew Pearson type VII distribution, skew normal distribution, stochastic representation, moment.

explores matrix variate skew elliptically contoured distributions. In particular, we discuss the first two moments of the matrix variate skew elliptically contoured distributions.

1. Introduction

The moments of matrix variate distributions are necessary both in theory and in applications, but often, their derivation is not straightforward and can constitute a work in and of itself. For example, Wen and Zhu derived stochastic representations and the first two moments, among other properties, of the multivariate skew Pearson type VII distribution and the skew t -distribution [1, 2]. In addition, Chen and Gupta extended several important properties of the multivariate skew normal distributions to the matrix variate case [3]. Also, Gupta et al. derived the first two moments of the multivariate skew normal distributions [4, 5, 6]. Furthermore, Akdemir and Gupta discussed more general matrix variate skew normal distributions, including moments [7].

In a previous work [8], we obtained the stochastic representations of the matrix variate skew elliptically contoured distributions (MSE). This paper extends that work by deriving the first two moments of the matrix variate skew elliptically contoured distributions (MSE).

In Section 2, we recall several definitions pertinent to the paper as well as the results from [8] for stochastic representations of matrix variate skew elliptically contoured distributions (MSE). In Section 3, we obtain moments of matrix variate skew elliptically contoured distributions.

2. Notation and Definitions

Matrix variate distributions have been studied by Gupta and Nagar, among others [9]. Below we include the results from [8] necessary for the derivation of moments. We refer the reader to [9] and [8] for additional definitions and theorems.

Let X be a $p \times m$ random matrix, which has a matrix variate normal distribution, i.e., $X \sim N_{p,m}(M, \Sigma \otimes \Psi)$, where $M \in \mathbb{R}^{p \times m}$ is a $p \times m$ mean matrix, Σ is a $p \times p$ positive definite matrix and Ψ is an $m \times m$ positive definite matrix. We use bold face upper and lower case letters to denote vectors, uppercase letters without bold face for matrices, and lowercase letters without bold face for elements of a vector or a matrix.

We list below the theorems and corollaries from [8] which are used in the following section.

Theorem 1 (Theorem 2 in [8]). *Assume*

$$V = \begin{pmatrix} \mathbf{V}_0 \\ V_1 \end{pmatrix} \sim E_{p+1,m}(0, \Sigma^* \otimes \Psi, \psi),$$

$$\Sigma^* = \begin{pmatrix} 1 & 0 \\ 0 & \Sigma \end{pmatrix}.$$

Σ is $p \times p$, Ψ is $m \times m$, \mathbf{V}_0 is $1 \times m$, V_1 is $p \times m$.

For $\delta_j \in (-1, 1)$, let

$$\left\{ \begin{array}{l} \mathbf{X}_0 = |\mathbf{V}_0|, \mathbf{X}_j = \delta_j |\mathbf{V}_0| + (1 - \delta_j^2)^{1/2} \mathbf{V}_j, j = 1, 2, \dots, p, \\ |\mathbf{V}_0| = (|v_{01}|, |v_{02}|, \dots, |v_{0m}|), \\ Z = (\mathbf{X}'_0, X'), X = (\mathbf{X}'_1, \dots, \mathbf{X}'_p)', \\ V^* = \begin{pmatrix} |\mathbf{V}_0| \\ V_1 \end{pmatrix}, \\ p_c = \int_{\mathbf{X}_0 < 0} |\Psi|^{-1/2} h(\text{tr} \mathbf{X}_0 \Psi^{-1} \mathbf{X}'_0) d\mathbf{X}_0. \end{array} \right.$$

Then the random matrix X has a matrix variate skew elliptically contoured distribution. Specifically,

$$X \sim MSE_{p,m}(\Sigma_\delta^{-1} \delta, \Sigma_\delta, \Psi, \Omega_{11.2} \Psi, \psi),$$

where

$$\delta = (\delta_1, \delta_2, \dots, \delta_p)', \Sigma_\delta = \delta\delta' + \Delta\Sigma\Delta, \Delta = \text{diag}\left((1 - \delta_1^2)^{\frac{1}{2}}, \dots, (1 - \delta_p^2)^{\frac{1}{2}}\right),$$

and $\Omega_{11.2} = 1 - \delta'\Sigma_\delta^{-1}\delta$.

In the following corollaries, we use the notation and definitions found in Theorem 1.

Corollary 1. Assume $V = \begin{pmatrix} \mathbf{V}_0 \\ V_1 \end{pmatrix} \sim N_{p+1,m}(0, \Sigma^* \otimes \Psi)$. Then the random matrix X has a matrix variate skew normal distribution. The pdf of the random matrix X can be written as

$$\begin{aligned} f(X) &= \left[\int_{\mathbf{X}_0 < 0} \det(\Psi)^{-\frac{1}{2}} (2\pi)^{-\frac{1}{2}m} \exp\left\{-\frac{1}{2} \mathbf{X}_0 \Psi^{-1} \mathbf{X}_0'\right\} d\mathbf{X}_0 \right]^{-1} \\ &\quad \times \det(\delta\delta' + \Delta\Sigma\Delta)^{-\frac{1}{2}m} \det(\Psi)^{-\frac{1}{2}p} (2\pi)^{-\frac{1}{2}pm} \exp\left\{-\frac{1}{2} \text{tr}(X' \Sigma_\delta^{-1} X \Psi^{-1})\right\} \\ &\quad \times \int_{\mathbf{X}_0 < 0} \det(\Omega_{11.2})^{-\frac{1}{2}m} \det(\Psi)^{-\frac{1}{2}} (2\pi)^{-\frac{1}{2}m} \\ &\quad \times \exp\left\{-\frac{1}{2} \text{tr}((\mathbf{X}_0 - \delta'\Sigma_\delta^{-1} X')' \Omega_{11.2}^{-1} (\mathbf{X}_0 - \delta'\Sigma_\delta^{-1} X) \Psi^{-1})\right\} d\mathbf{X}_0 \\ &= \left[\int_{\mathbf{X}_0 < 0} \phi_m(X_0, 0, \Psi) d\mathbf{X}_0 \right]^{-1} \phi_{p,m}(X, 0, \Sigma_\delta \otimes \Psi) \\ &\quad \times \Phi_m(X' \Sigma_\delta^{-1} \delta, 0, \Omega_{11.2} \Psi) \end{aligned}$$

which is denoted by $MSN_{p,m}(\Sigma_\delta^{-1}\delta, \Sigma_\delta, \Psi, \Omega_{11.2}\Psi)$.

Corollary 2. Assume $\begin{pmatrix} \mathbf{V}_0 \\ V_1 \end{pmatrix} \sim PVII_{p+1,m}(0, \Sigma^*, \Psi, q, r)$. Then the random matrix X has a matrix variate skew Pearson type VII distribution. The pdf of the random matrix X can be written as

$$\begin{aligned}
 f(X) &= \left[\int_{\mathbf{X}_0 < 0} \frac{\Gamma(q)}{(\pi\lambda)^{m/2} \Gamma(q - m/2) |\Psi|^{1/2}} \left(1 + \frac{\mathbf{X}_0 \Psi^{-1} \mathbf{X}'_0}{\lambda} \right)^{-q} d\mathbf{X}_0 \right]^{-1} \\
 &\quad \times \frac{\Gamma(q)}{(\pi\lambda)^{pm/2} \Gamma(q - pm/2) |\Sigma_\delta|^{m/2} |\Psi|^{p/2}} \left(1 + \frac{\text{tr}(X' \Sigma_\delta^{-1} X \Psi^{-1})}{\lambda} \right)^{-q} \\
 &\quad \times \int_{\mathbf{X}_0 < 0} \frac{\Gamma(q)}{(\pi\lambda)^{m/2} \Gamma(q - m/2) |\Omega_{11.2} \Psi|^{1/2}} \\
 &\quad \times \left\{ 1 + \frac{[\mathbf{X}_0 - \delta'(\Sigma_\delta)^{-1} X](\Omega_{11.2} \Psi)^{-1} [\mathbf{X}_0 - \delta'(\Sigma_\delta)^{-1} X]'}{\lambda} \right\}^{-q} d\mathbf{X}_0.
 \end{aligned}$$

This matrix variate skew Pearson type VII distribution is denoted by

$$MSPVII_{p,m}(\Sigma_\delta^{-1} \delta, \Sigma_\delta, \Psi, \Omega_{11.2} \Psi, q, \lambda).$$

In the next theorem, we use the same notation and definitions as in Theorem 1.

Theorem 2 (Theorem 3 in [8]). *Assume*

$$V = \begin{pmatrix} \mathbf{V}_0 \\ \mathbf{V}_1 \end{pmatrix} \sim E_{p+1,m}(0, \Sigma^* \otimes \Psi, \psi).$$

Also assume that $\Sigma = AA'$, and $\Psi = BB'$. For $\delta_j \in (-1, 1)$, let

$$\begin{pmatrix} \mathbf{X}_1 \\ \vdots \\ \mathbf{X}_p \end{pmatrix} = \begin{pmatrix} \delta_1 \\ \vdots \\ \delta_p \end{pmatrix} |\mathbf{V}_0| + \Delta \begin{pmatrix} \mathbf{V}_1 \\ \vdots \\ \mathbf{V}_p \end{pmatrix}, \text{ or } X = \delta |\mathbf{V}_0| + \Delta \mathbf{V}_1.$$

Then the random matrix X has the following stochastic representation:

$$X = R(\delta |\mathbf{U}_1| + \Delta \mathbf{A} \mathbf{U}_2) \mathbf{B}', \quad (1)$$

where R is a nonnegative random variable, \mathbf{U}_1 is $1 \times m$ and uniformly distributed on sphere S_m , \mathbf{U}_2 is $p \times m$ and $\text{vec}(\mathbf{U}_2')$ is uniformly distributed

on sphere S_{pm} . In addition, R , \mathbf{U}_1 , and U_2 are independent. Furthermore, $\psi(x) = \int_0^\infty \Omega_{pm}(r^2 x) dF(r)$, $x \geq 0$, where $\Omega_{pm}(\mathbf{t}'\mathbf{t})$, $\mathbf{t} \in \mathbb{R}^{pm}$ denotes the characteristic function of $\text{vec}(U_2')$, and $F(r)$ denotes the distribution function of r .

The next corollary follows immediately. Moreover, its statement uses the same notation as in Theorem 2 and also assumes that $\Sigma = AA'$, $\Psi = BB'$.

Corollary 3. Assume $V = \begin{pmatrix} \mathbf{V}_0 \\ \mathbf{V}_1 \end{pmatrix} \sim N_{p+1,m}(0, \Sigma^* \otimes \Psi)$. For $\delta_j \in (-1, 1)$,

let

$$\begin{pmatrix} \mathbf{Z}_1 \\ \vdots \\ \mathbf{Z}_p \end{pmatrix} = \begin{pmatrix} \delta_1 \\ \vdots \\ \delta_p \end{pmatrix} | \mathbf{V}_0 | + \Delta \begin{pmatrix} \mathbf{V}_1 \\ \vdots \\ \mathbf{V}_p \end{pmatrix}, \text{ or } Z = \delta | \mathbf{V}_0 | + \Delta \mathbf{V}_1.$$

Then the random matrix Z is distributed as matrix variate skew normal and has the following stochastic representation:

$$Z = R_0(\delta | \mathbf{U}_1 | + \Delta \mathbf{A}U_2)B', \quad (2)$$

where \mathbf{U}_1 , U_2 , A , B are the same as in Theorem 2.

Based on the definitions and the stochastic representation obtained in [8], the moments of matrix variate skew elliptically contoured distribution can be derived, as we now show.

3. Moments of Matrix Variate Skew Elliptically Contoured Distribution

The moments of matrix variate skew elliptical distributions can be found using the moments of the matrix variate skew normal distribution since the latter is easy to find. Suppose that the random matrix X is distributed as a general matrix variate skew elliptically contoured distribution (MCD) such as in Theorem 2 and also suppose that the random matrix Z is distributed as a matrix variate skew normal as in Corollary 3. Then the moments of the skew

MCD X are associated with the moments of skew normal Z . Specifically, we have the following relationship between them:

$$\mathbb{E}\left(\prod_{i=1}^p \prod_{j=1}^m X_{ij}^{s_{ij}}\right) = \frac{\mathbb{E}\left(R^{\sum_{i=1}^p \sum_{j=1}^m s_{ij}}\right)}{\mathbb{E}\left(R_0^{\sum_{i=1}^p \sum_{j=1}^m s_{ij}}\right)} \mathbb{E}\left(\prod_{i=1}^p \prod_{j=1}^m Z_{ij}^{s_{ij}}\right), \quad (3)$$

where X_{ij} is the (i, j) element of random matrix X , Z_{ij} is the (i, j) coefficient of the random matrix Z , and R_0^2 is distributed as χ^2 with degrees of freedom pm . Equation (3) leads to the following results:

$$\mathbb{E}(X) = \frac{\mathbb{E}(R)}{\mathbb{E}(R_0)} \times \mathbb{E}(Z), \quad \mathbb{E}(XX') = \frac{\mathbb{E}(R^2)}{\mathbb{E}(R_0^2)} \times \mathbb{E}(ZZ'), \quad (4)$$

$$\mathbb{E}(\text{vec}(X') \otimes \text{vec}(X')) = \frac{\mathbb{E}(R^2)}{\mathbb{E}(R_0^2)} \times \mathbb{E}(\text{vec}(Z') \otimes \text{vec}(Z')). \quad (5)$$

It also leads to our next theorem.

Theorem 3. *Suppose that the random matrix X is distributed as a matrix variate skew Pearson type VII defined in Corollary 2, i.e., $X \sim MSPVII_{p,m}(\Sigma_\delta^{-1}\delta, \delta\delta' + \Delta\Sigma\Delta, \Psi, \Omega_{11,2}\Psi, q, \lambda)$. Suppose also that the random matrix Z is distributed as a matrix variate skew normal defined in Corollary 1, i.e., $Z \sim MSN_{p,m}(\Sigma_\delta^{-1}\delta, \Sigma_\delta, \Psi, \Omega_{11,2}\Psi)$. Then we have*

$$\mathbb{E}(X) = \frac{\Gamma\left(q - \frac{(p+1)m+1}{2}\right)}{\Gamma\left(q - \frac{(p+1)m}{2}\right)} \sqrt{\frac{\lambda}{2}} \times \mathbb{E}(Z), \quad (6)$$

$$\mathbb{E}(XX') = \frac{\lambda}{2q - (p+1)m - 2} \times \mathbb{E}(ZZ'), \quad (7)$$

$$\mathbb{E}[\text{vec}(X') \otimes \text{vec}(X')] = \frac{\lambda}{2q - (p+1)m - 2} \times \mathbb{E}[\text{vec}(Z') \otimes \text{vec}(Z')]. \quad (8)$$

Proof. Based on (2.5.17) on p. 59 of [10], the pdf of the random variable R can be found using the function h defined in (6) of [8]:

$$\begin{aligned}
g(r) &= \frac{2\pi^{\frac{1}{2}(p+1)m}}{\Gamma\left(\frac{1}{2}(p+1)m\right)} r^{(p+1)m-1} h(r^2) \\
&= \frac{2\pi^{\frac{1}{2}(p+1)m}}{\Gamma\left(\frac{1}{2}(p+1)m\right)} r^{(p+1)m-1} \frac{\Gamma(q)}{(\pi\lambda)^{\frac{(p+1)m}{2}} \Gamma\left(q - \frac{(p+1)m}{2}\right)} \left(1 + \frac{r^2}{\lambda}\right)^{-q} \\
&= \frac{2}{\Gamma\left(\frac{1}{2}(p+1)m\right)} \frac{\Gamma(q) r^{(p+1)m-1}}{\lambda^{\frac{(p+1)m}{2}} \Gamma\left(q - \frac{(p+1)m}{2}\right)} \left(1 + \frac{r^2}{\lambda}\right)^{-q}.
\end{aligned}$$

Thus,

$$\begin{aligned}
\mathbb{E}(R) &= \int_0^{+\infty} r g(r) dr \\
&= \frac{2\Gamma(q)}{\lambda^{\frac{(p+1)m}{2}} \Gamma\left(\frac{(p+1)m}{2}\right) \Gamma\left(q - \frac{(p+1)m}{2}\right)} \int_0^{+\infty} r^{(p+1)m} \left(1 + \frac{r^2}{\lambda}\right)^{-q} dr \\
&= \frac{\Gamma\left(q - \frac{(p+1)m+1}{2}\right) \Gamma\left(\frac{(p+1)m+1}{2}\right)}{\Gamma\left(q - \frac{(p+1)m}{2}\right) \Gamma\left(\frac{(p+1)m}{2}\right)} \lambda^{\frac{1}{2}}.
\end{aligned}$$

Next, we find the expectation of R_0 . By (2.5.17) on p. 59 of [10], we have

$$g(r_0) = \frac{2\pi^{\frac{1}{2}(p+1)m}}{\Gamma\left(\frac{1}{2}(p+1)m\right)} r_0^{(p+1)m-1} h(r_0^2)$$

$$= \frac{2\pi^{\frac{1}{2}(p+1)m}}{\Gamma\left(\frac{1}{2}(p+1)m\right)} r_0^{(p+1)m-1} \left(\frac{1}{(2\pi)^{\frac{(p+1)m}{2}}} \right) e^{-\frac{1}{2}r_0^2}.$$

Let $s = r_0^2$. Then

$$g(s) = \left(\Gamma\left(\frac{m}{2}\right) 2^{\frac{1}{2}(p+1)m} \right)^{-1} s^{\frac{(p+1)m}{2}-1} e^{-\frac{1}{2}s}.$$

Therefore, S or $R_0^2 \sim \chi^2((p+1)m)$ and

$$\mathbb{E}(R_0) = \mathbb{E}(\sqrt{S})$$

$$= \int_0^{+\infty} \left(\Gamma\left(\frac{1}{2}(p+1)m\right) 2^{\frac{1}{2}(p+1)m} \right)^{-1} s^{\frac{(p+1)m}{2}-1+\frac{1}{2}} e^{-\frac{1}{2}s} ds$$

$$= \frac{\Gamma\left(\frac{(p+1)m+1}{2}\right)}{\Gamma\left(\frac{(p+1)m}{2}\right)} \sqrt{2},$$

$$\frac{\mathbb{E}(R)}{\mathbb{E}(R_0)} = \frac{\Gamma\left(q - \frac{(p+1)m+1}{2}\right)}{\Gamma\left(q - \frac{(p+1)m}{2}\right)} \sqrt{\frac{\lambda}{2}}. \quad (9)$$

On the other hand,

$$\mathbb{E}(R^2) = \int_0^{+\infty} r^2 g(r) = \frac{(p+1)m}{2q - (p+1)m - 2} \lambda, \quad \text{and} \quad \mathbb{E}(R_0^2) = (p+1)m,$$

$$\frac{\mathbb{E}(R^2)}{\mathbb{E}(R_0^2)} = \frac{\lambda}{2q - (p+1)m - 2}. \quad (10)$$

Hence, equations (6), (7) and (8) are proved using (9) and (10). \square

In order to use Theorem 3 to find the first two moments of any matrix variate skew elliptically contoured distribution, we have to first find the first two moments of matrix variate skew normal distributions. Gupta et al. derived the first two moments of multivariate skew normal distributions [4, 5, 6]. To generalize these results to matrix variate skew normal distribution, we utilize moment generating functions. In particular, Arellano-Valle and Azzalini derived the moment generating function (mgf) of a multivariate unified skew normal distribution [11]. Harrar and Gupta presented the mgf for general matrix variate skew normal distributions [12]. The following lemma is from Harrar and Gupta [12].

Lemma 1. *Assume $X \sim MSN_{p,m}(\mathbf{b}, \Sigma, \Psi, \Omega)$. Then the moment generating function of X can be written as*

$$M_X(T) = c \times \text{etr} \left\{ \frac{1}{2} \Sigma T \Psi T' \right\} \Phi_m(\Psi T' \Sigma \mathbf{b}, \Omega + \mathbf{b}' \Sigma \mathbf{b} \Psi), \quad (11)$$

where $c = [\Phi_m(0; \Omega + \mathbf{b}' \Sigma \mathbf{b} \Psi)]^{-1}$.

In order to simplify notation in the sequel, we let $\Xi = \Omega + \mathbf{b}' \Sigma \mathbf{b} \Psi$ and $\boldsymbol{\theta} = \Psi T' \Sigma \mathbf{b}$. Note that $\boldsymbol{\theta} = \text{vec}(\Psi T' \Sigma \mathbf{b}) = [(\mathbf{b}' \Sigma) \otimes \Psi] \times \text{vec}(T')$. Furthermore, we let $\mathbf{t} = \text{vec}(T')$, $D = (\mathbf{b}' \Sigma) \otimes \Psi$.

Theorem 4. *Suppose $X \sim MSN_{p,m}(\mathbf{b}, \Sigma, \Psi, \Omega)$. Then the expectation of X can be written as*

$$\mathbb{E}(X) = [\Phi_m(0; \Xi)]^{-1} \times \sum_{i=1}^m \sum_{j=1}^p G_{ij}^{mp}(0; D, \Xi) \times H_{ij}, \quad (12)$$

where the matrix H_{ij} ($m \times p$) has unit element at the (i, j) th place and zero elsewhere and that

$$\begin{aligned} G_{ij}^{mp}(0; D, \Xi) &= \frac{\partial}{\partial t_{ij}} \Phi_m(D\mathbf{t}; \Xi) \Big|_{T=0} \\ &= \frac{1}{\sqrt{2\pi} |\Xi|^{1/2}} \sum_{k=1}^m D_{k, m(i-1)+j} (\Xi_{(k)}^{-1})^{-1} |^{1/2} \Phi_{m-1}[0; (\Xi_{(k)}^{-1})^{-1}], \end{aligned}$$

where $D_{k,m(i-1)+j}$ is the element $(k, m(i-1)+j)$ of matrix D , $\Xi_{(k)}^{-1}$ is the matrix constructed by eliminating the k th row and the k th column of Ξ^{-1} , with the convention that $\Xi_{(k)}^{-1} = (\Xi^{-1})_{(k)}$.

Proof. Since $\frac{\partial \text{tr}(\Sigma T \Psi T')}{\partial T} = 2\Sigma T \Psi$, the derivative with respect to T of the moment generating function (11) can be written in the form

$$\begin{aligned} \frac{\partial M_X(T)}{\partial T} &= c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} (\Sigma T \Psi) \times \Phi_m(\boldsymbol{\theta}, \Xi) \\ &\quad + c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \times \frac{\partial}{\partial T} \Phi_m(\boldsymbol{\theta}, \Xi). \end{aligned} \quad (13)$$

After a lengthy calculation (the reader is referred to similar derivations in [4, 5, 6] for details), we have

$$\begin{aligned} \mathbb{E}(X) &= \left. \frac{\partial M_X(T)}{\partial T} \right|_{T=0} \\ &= [\Phi_m(0; \Xi)]^{-1} \times \left. \frac{\partial}{\partial T} \Phi_m(\boldsymbol{\theta}, \Xi) \right|_{T=0}, \end{aligned}$$

where

$$\left. \frac{\partial}{\partial T} \Phi_m(\boldsymbol{\theta}, \Xi) \right|_{T=0} = \left. \frac{\partial}{\partial T} \Phi_m(D\mathbf{t}; \Xi) \right|_{T=0}$$

and

$$\left. \frac{\partial}{\partial T} \Phi_m(D\mathbf{t}; \Xi) \right|_{T=0} = \sum_{i=1}^m \sum_{j=1}^p G_{ij}^{mp}(0; D, \Xi) H_{ij}.$$

Therefore, the theorem is proved. \square

Finally, we have the following theorem, whose proof – like that of Theorem 4 – requires a long elementary calculation of a second derivative which is similar in details to derivations in [4, 5, 6].

Theorem 5. Suppose $X \sim MSN_{p,m}(\mathbf{b}, \Sigma, \Psi, \Omega)$. Then the expectation of $\text{vec}(X') \otimes \text{vec}(X')'$ can be written as

$$\begin{aligned} & \mathbb{E}(\text{vec}(X') \otimes \text{vec}(X')') \\ &= \Sigma \otimes \Psi + [\Phi_m(0; \Xi)]^{-1} \times \sum_{i'=1}^{pm} \sum_{j'=1}^{pm} G_{i'j'}^{mp}(0; D, \Xi) \times H_{i'j'}, \end{aligned}$$

where the matrix $H_{i'j'}(pm \times pm)$ has unit element at the (i', j') th place and zero elsewhere,

$$i' = m(i-1) + j, \quad j' = m(a-1) + b; \quad i, a = 1, \dots, m; \quad j, b = 1, \dots, p;$$

and

$$\begin{aligned} & G_{i'j'}^{pm}(0, D, \Xi) \\ &= \frac{\partial}{\partial t_{ij} \partial t_{ab}} \Phi_m(D\mathbf{t}, \Xi) \Big|_{\mathbf{t}=0} \\ &= \frac{1}{2\pi |\Xi|^{1/2}} \sum_{k=1}^m \sum_{l \neq k}^m D_{l, m(i-1)+j} D_{k, m(a-1)+b} |\Xi_{(k,l)}^{-1}| \Phi_{m-2}(0; [\Xi_{(k,l)}^{-1}]^{-1}) \\ & \quad + \sum_{k=1}^m \sum_{\substack{i=1 \\ i \neq k}}^m \int_{-\infty}^0 \cdots \int_{-\infty}^0 \cdots \int_{-\infty}^0 D_{l, m(i-1)+j} D_{k, m(a-1)+b} \Xi^{ki} x_i \\ & \quad \times \phi_{m-1}[x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_m; (\Xi_{(k)}^{-1})^{-1}] dx_1 \cdots dx_{k-1} dx_{k+1} \cdots dx_m, \end{aligned}$$

where Ξ^{kk} is the element (k, k) of Ξ^{-1} and $\Xi_{(i,j)}^{-1}$ is the matrix constructed by eliminating the i th and j th rows and the i th and j th columns of Ξ^{-1} , with $\Xi_{(i,j)}^{-1} = (\Xi^{-1})_{(i,j)}$.

Proof. Based on (11), we have

$$\begin{aligned} \frac{\partial M_X(T)}{\partial \text{vec}(T')} &= c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \frac{\partial \text{tr}\left(\frac{1}{2}\Sigma T \Psi T'\right)}{\partial \text{vec}(T')} \Phi_m(\boldsymbol{\theta}, \Xi) \\ &= c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \frac{\partial}{\partial \text{vec}(T')} \Phi_m(\boldsymbol{\theta}, \Xi). \end{aligned}$$

Thus,

$$\begin{aligned} &\frac{\partial M_X(T)}{\partial \text{vec}(T') \partial \text{vec}(T')} \\ &= c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \frac{\partial \text{tr}\left(\frac{1}{2}\Sigma T \Psi T'\right)}{\partial \text{vec}(T')} \otimes \frac{\partial \text{tr}\left(\frac{1}{2}\Sigma T \Psi T'\right)}{\partial \text{vec}(T')} \Phi_m(\boldsymbol{\theta}, \Xi) \\ &\quad + c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \frac{\partial \text{tr}\left(\frac{1}{2}\Sigma T \Psi T'\right)}{\partial \text{vec}(T') \partial \text{vec}(T')} \Phi_m(\boldsymbol{\theta}, \Xi) \\ &\quad + c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \frac{\partial \Phi_m(\boldsymbol{\theta}, \Xi)}{\partial \text{vec}(T')} \otimes \frac{\partial \text{tr}\left(\frac{1}{2}\Sigma T \Psi T'\right)}{\partial \text{vec}(T')} \\ &\quad + c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \frac{\partial^2 \Phi_m(\boldsymbol{\theta}, \Xi)}{\partial \text{vec}(T') \partial \text{vec}(T')} \\ &= c \times \text{etr}\left\{\frac{1}{2}\Sigma T \Psi T'\right\} \left\{ \text{vec}(\Psi T' \Sigma) \otimes (\text{vec}(\Psi T' \Sigma))' \Phi_m(\boldsymbol{\theta}, \Xi) \right. \\ &\quad + (\Sigma \otimes \Psi) \Phi_m(\boldsymbol{\theta}, \Xi) + \frac{\partial \Phi_m(\boldsymbol{\theta}, \Xi)}{\partial \text{vec}(T')} \otimes (\text{vec}(\Psi T' \Sigma))' \\ &\quad \left. + \frac{\partial^2 \Phi_m(\boldsymbol{\theta}, \Xi)}{\partial \text{vec}(T') \partial \text{vec}(T')} \right\}. \end{aligned}$$

Hence,

$$\begin{aligned} \mathbb{E}[\text{vec}(X') \otimes \text{vec}(X')'] &= \left. \frac{\partial M_X(T)}{\partial \text{vec}(T') \partial \text{vec}(T')'} \right|_{T=0} \\ &= \Sigma \otimes \Psi + c \times \left. \frac{\partial^2 \Phi_m(\boldsymbol{\theta}, \Xi)}{\partial \text{vec}(T') \partial \text{vec}(T')'} \right|_{T=0}. \quad \square \end{aligned}$$

4. Concluding Remarks

The set of skew elliptically contoured distributions contains the elliptically contoured distributions. In [8], we defined and explored the matrix variate skew elliptically contoured distribution, obtaining in the process two stochastic representations. In this work, we have extended that effort by obtaining the first two moments of a subfamily of this large family of distributions.

In particular, the moment results obtained in this paper are useful in both theory and application. Future directions include expanding upon the utility and importance of these results.

Acknowledgments

The authors graciously acknowledge the financial and collaborative support of the Institute for Quantitative Biology at East Tennessee State University and Zhang's research is supported by the US NSF grant DMS-1106586 and Wisconsin Alumni Research Foundation. The authors would also like to thank the referees for many useful suggestions and comments which improved the presentation in this paper.

References

- [1] Y. J. Wen and D. Y. Zhu, New multivariate skew t distributions generated from skew Pearson vii distributions, *Chinese Journal of Probability and Statistics* 25(5) (2009), 485-498.

- [2] Y. J. Wen and D. Y. Zhu, Properties of new multivariate skew t distributions generated from skew Pearson VII distributions, *Chinese Journal of Probability and Statistics* 26(26) (2010), 367-383.
- [3] J. T. Chen and A. K. Gupta, Matrix variate skew normal distributions, *Statistics* 39(3) (2005), 247-253.
- [4] G. González-Farías, J. A. Domínguez-Molina and A. K. Gupta, A general multivariate skew normal distribution, Technical Report 01-09, Department of Mathematics and Statistics, Bowling Green State University, 2001.
- [5] A. K. Gupta, G. González-Farías and J. A. Domínguez-Molina, A multivariate skew normal distribution, Technical Report No. 01-10, Department of Mathematics and Statistics, Bowling Green State University, 2001.
- [6] A. K. Gupta, G. González-Farías and J. A. Domínguez-Molina, A multivariate skew normal distribution, *Journal of Multivariate Analysis* 89 (2004), 181-190.
- [7] D. Akdemir and A. K. Gupta, A matrix variate skew distribution, *European Journal of Pure and Applied Mathematics* 3(10) (2010), 128-140.
- [8] S. Zheng, C. Zhang and J. Knisley, Stochastic representations of the matrix variate skew elliptically contoured distributions, *Advances and Applications in Statistics* 33(2) (2013), 83-98.
- [9] A. K. Gupta and D. K. Nagar, *Matrix Variate Distributions*, Chapman & Hall/CRC, Boca Raton, London, New York, Washington, D.C., 2000.
- [10] K. T. Fang and Y. T. Zhang, *Generalized Multivariate Analysis*, Springer-Verlag, Berlin Heidelberg, New York, London, Paris, Tokyo, Hong Kong, 1990.
- [11] R. B. Arellano-Valle and A. Azzalini, On the unification of families of skew-normal distributions, *Scand. J. Statist.* 33 (2006), 561-574.
- [12] S. Harrar and A. K. Gupta, On matrix variate skew normal distribution, *Statistics* 42 (2008), 179-194.