

Apr 12th, 11:00 AM - 11:15 AM

Association of Age, Gender and Race in Chronic Kidney Disease Patients with and without Dialysis

Busayo Onatolu

Shimin Zheng
East Tennessee State University

Hemang Panchal
East Tennessee State University

Edward Leinaar
East Tennessee State University

Follow this and additional works at: <https://dc.etsu.edu/asrf>

Onatolu, Busayo; Zheng, Shimin; Panchal, Hemang; and Leinaar, Edward, "Association of Age, Gender and Race in Chronic Kidney Disease Patients with and without Dialysis" (2019). *Appalachian Student Research Forum*. 23.
<https://dc.etsu.edu/asrf/2019/schedule/23>

This Oral presentation is brought to you for free and open access by the Events at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in Appalachian Student Research Forum by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

ASSOCIATION OF AGE, GENDER AND RACE IN CHRONIC KIDNEY DISEASE PATIENTS WITH AND WITHOUT DIALYSIS

Busayo Onatolu, Hemang Panchal, Edward Francis Leinaar, *Shimin Zheng, Timir K. Paul

East Tennessee State University

*Sponsoring faculty

Background

- Studies have shown that chronic kidney disease (CKD) is common among adults in the United States.
- The Center for Disease Control and Prevention (CDC) states that 30 million people or 15% of US adults are estimated to have CKD.
- 48% of those with severely reduced kidney function not aware of having CKD and therefore do not receive hemodialysis.

Purpose

- The purpose of this study was to investigate the association of age, gender and race in chronic kidney diseased patients with and without dialysis.

Methods

- A nationwide inpatient sample database from 2012-2014 was used to identify all patients admitted to the hospital using International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes (n= 534,845).
- Patients with dialysis dependent CKD (n=8,100) and CKD without dialysis (n=51,285) were compared to non-CKD patients (n=475,460).
- Hierarchical logistic regression was performed and $p < 0.05$ was considered as the level of significance.

Variables

- Explanatory Variables
 - Age
 - Gender
 - Race

Results

- The sample included 534,845 patients admitted to the hospital.
- 11.1% of the sample had CKD
 - 9.59% had CKD without dialysis and 1.51% had CKD with dialysis
 - Among patients with CKD, 13.64% were on HD
 - 86.34% with CKD were non HD patients
- Age
 - Mean age for total population; male - 61.36, female - 68.41
 - Mean age for patients with CKD,
 - Non-HD - males - 71.12, females - 76.96
 - With HD - males - 66.21, females - 67.62
 - Higher proportion of patients with CKD without HD in the ≥ 80 years age group ($\geq 80 = 37.84\%$, $65-79 = 36.94\%$, $50-64 = 20.80\%$, $35-49 = 4.12\%$ and $18-34 = 0.30\%$).
 - Among patients with CKD patients with HD, 65-79 years of age were the highest in the age group. ($\geq 80 = 16.30\%$, $65-79 = 41.79\%$, $50-64 = 33.09\%$, $35-64 = 8.09\%$ and $18-34 = 1.29\%$).
 - 9.59% had CKD without dialysis and 1.51% had CKD with dialysis.

Results(contd.)

- Gender
 - Males had higher rates of CKD with and without HD
 - 54.51% of males had CKD with non-HD, 8.03% were on HD
 - 31.85% of females had CKD with non-HD, 5.62% were on HD
- Race
 - 65.09% whites with mean age of 74.57 had CKD without HD, 7.17% with mean age of 69.20 had CKD with HD
 - 9.53% Blacks with mean age of 67.38 had CKD without HD, 3.24% with mean age of 63.62 had CKD with HD
 - 2.52% Asians with mean age of 73.26 had CKD without HD, 0.72% with mean age of 67.20 had CKD with HD
 - 0.40% Native Americans/ AK natives with mean age 70.47 had CKD without HD, 0.13% with age of 65.14 had CKD with HD
 - 8.65% from other races had CKD without HD, 2.56% had CKD with HD

Logistic regression of variables

Variable	Odds ratio	95% confidence interval	P-value
Age			
18 - 34 vs ≥ 80	4.023	2.941 - 5.504	0.0010
35 - 49 ≥ 80	3.802	3.404 - 4.246	<.0001
50 - 64 vs ≥ 80	3.366	3.123 - 3.628	<.0001
65 - 79 vs ≥ 80	2.546	2.375 - 2.730	0.2733
Gender			
Female vs male	1.448	1.375 - 1.525	<.0001
Race			
Asian vs white	2.486	2.210 - 2.797	<.0001
Black vs white	2.479	2.324 - 2.644	<.0001
Hispanic vs white	2.727	2.531 - 2.938	<.0001
Native American vs white	2.288	1.740 - 3.008	0.1234
Other vs white	1.265	1.090 1.469	<.0001

Conclusion

- From this study, males had higher rates of CKD with and without HD than females
- The age group ≥ 80 years had higher proportion of CKD without HD and those between 65-79 years had higher number of CKD with HD.
- Among the race, whites had higher rates of CKD with and without HD than other races.