



SCHOOL of
GRADUATE STUDIES
EAST TENNESSEE STATE UNIVERSITY

East Tennessee State University
Digital Commons @ East
Tennessee State University

Electronic Theses and Dissertations

Student Works

5-2005

Graph Theory for the Secondary School Classroom.

Dayna Brown Smithers
East Tennessee State University

Follow this and additional works at: <https://dc.etsu.edu/etd>



Part of the [Algebra Commons](#)

Recommended Citation

Smithers, Dayna Brown, "Graph Theory for the Secondary School Classroom." (2005). *Electronic Theses and Dissertations*. Paper 1015.
<https://dc.etsu.edu/etd/1015>

This Thesis - Open Access is brought to you for free and open access by the Student Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

Graph Theory for the Secondary School Classroom

A thesis
presented to
the faculty of the Department of Mathematics
East Tennessee State University

In partial fulfillment
of the requirements for the degree
Master of Science in Mathematical Sciences

by
Dayna Brown Smithers
May 2005

Teresa Haynes, Ph.D., Chair
Debra Knisley, Ph.D.
Michel Helfgott, Ed.D
Janice Huang, Ph.D.

Keywords: vertex coloring, spanning tree, domination,
hamiltonian, high school

ABSTRACT

Graph Theory for the Secondary School Classroom

by

Dayna Brown Smithers

After recognizing the beauty and the utility of Graph Theory in solving a variety of problems, the author concluded that it would be a good idea to make the subject available for students earlier in their educational experience. In this thesis, the author developed four units in Graph Theory, namely Vertex Coloring, Minimum Spanning Tree, Domination, and Hamiltonian Paths and Cycles, which are appropriate for high school level.

Copyright by Dayna Brown Smithers 2005

DEDICATION

This thesis is dedicated to the many people who have supported me throughout my educational journey. To my parents, David and Yoshie Brown, your constant love, support, and prayers are what have gotten me through this journey. To my husband, Hillard Smithers, I could not have finished without your unconditional love and never ending encouragement. You have brought so much joy and laughter into my life. I love you with all of my heart! To Dr. Overtoun Jenda for starting me on my mathematics journey and for always believing in me. To the many friends that I have made along the way I would like to thank you for making this journey unforgettable. Most importantly I would like to thank my Heavenly Father. “The Lord is my strength and my shield; my heart trusts in him, and I am helped.” Psalm 28:7

ACKNOWLEDGEMENTS

I would like to truly thank Dr. Teresa Haynes and Dr. Janice Haung. This thesis would not have happened without your encouragement and support. I would also like to thank Travis Coake for his LATEX skills and the three pictures that he created for me.

CONTENTS

ABSTRACT	2
DEDICATION	4
ACKNOWLEDGEMENTS	5
LIST OF FIGURES	9
1 INTRODUCTION	10
1.1 The Idea	10
1.2 Available Options	10
1.3 Selecting the Units	11
2 MATHEMATICAL MODELING	14
3 GRAPH THEORY INTRODUCTION UNIT	19
3.1 Introduction	19
3.2 In Class Examples	20
3.3 Homework Exercises	24
3.4 Answer Key	25
4 VERTEX COLORING UNIT	27
4.1 Introduction	27
4.2 In Class Exercises	29
4.3 Problem Solving	30
4.4 Applications	32
4.5 Addendum	39
4.6 Homework Exercises	42
4.7 Answer Key	48

5	MINIMUM SPANNING TREE UNIT	58
5.1	Introduction	58
5.2	In Class Exercises	60
5.3	Problem Solving	62
5.4	Applications	68
5.5	Homework Exercises	75
5.6	Answer Key	80
6	DOMINATION UNIT	89
6.1	Introduction	89
6.2	In Class Exercises	91
6.3	Problem Solving	92
6.4	Applications	94
6.5	Homework Exercises	101
6.6	Answer Key	107
7	HAMILTONIAN PATHS AND CYCLES UNIT	116
7.1	Introduction	116
7.2	In Class Exercises	119
7.3	Problem Solving	120
7.4	Application	123
7.5	Homework Exercises	129
7.6	Answer Key	134
8	SUMMARY	145
	BIBLIOGRAPHY	148

VITA	150
------------	-----

LIST OF FIGURES

1	Fence Problem	15
2	Process of Mathematical Modeling	18
3	A graph G	20
4	A graph H ; H is a subgraph of G	21
5	Complete Graphs	22
6	Paths	22
7	Cycles	23
8	Vertex Coloring Example	27
9	Chromatic Number Example	30
10	Trees	58
11	Spanning Trees	60
12	Minimum Spanning Trees	64
13	Domination Example	89
14	The Five Queens Graph	92
15	Hamiltonian Paths and Cycles	117
16	Hamilton's Icosian Game	122

1 INTRODUCTION

1.1 The Idea

“Among the themes that have been central to mathematics education during the last 30 years are relations between mathematics and the real world (or better, according to Pollak, 1979, the “rest of the world”)” [11]. It is with this theme in mind that we shall approach the teaching of Graph Theory to high school students.

During the author’s Fall 2003 semester at East Tennessee State University [ETSU], the author enrolled in MATH 5340, Graph Theory and its Applications, under Dr. Teresa Haynes. Graph Theory is a relatively new area in mathematics that is only touched upon in some discrete mathematics classes in high schools and in the applied mathematics track in many universities. Due to the author’s undergraduate degree in pure mathematics, she had never taken a course in Graph Theory. During the course, the author discovered how important mathematical modeling is in our society and how Graph Theory is such a useful tool in this area. The author began to think how she would have loved to have been introduced to this area of math long before her graduate studies. This led to the idea of creating Graph Theory units for high school students, so that they can have an earlier start to discovering this wonderful area of mathematics.

1.2 Available Options

The author first researched to see if a high school mathematics class that taught Graph Theory already existed. She discovered that, according to the Mathematics

Curriculum Standards, the class Discrete Mathematics with Statistics and Probability, Course Number 3135, does include a small amount of Graph Theory. The class is structured around three subjects, Discrete Mathematics, Probability, and Statistics. In the Discrete Mathematics section, the students “use vertex-edge graphs to solve network problems such as finding circuits, critical paths, minimum spanning trees, and adjacency matrices” [6].

Upon learning this information, the author wanted to see if the high schools in her surrounding area had this class in their school curriculum. She visited the web sites of local high schools and discovered that this course was not in the local high schools’ curricula. The author followed-up by contacting some local high school teachers to see if there would be an interest either in a new course in Graph Theory or incorporating a section of Graph Theory into an existing course.

The consensus of the teachers was that their schools are so over populated at this current time that they do not have the manpower to add a course, or are having trouble filling the required courses, much less find room for an elective math course. However, some teachers did explain that they sometimes have extra time at the end of the year where they can teach supplementary material and they would like to teach a section of Graph Theory to their classes.

1.3 Selecting the Units

The teachers’ response led the author to create independent units of Graph Theory that can be used in a high school classroom when extra time permits. The units are designed for a teacher to be able to cover a selected topic in Graph Theory in one week.

The Introduction Unit is designed to be taught before any of the other units. It gives basic definitions and terminology in Graph Theory, necessary background to the other units. The other four units, Vertex Coloring, Minimum Spanning Tree, Domination, and Hamiltonian Paths and Cycles are independent of each other, thus allowing the mathematics teachers to teach any unit they feel fits in their lesson plan. Ideally, if a teacher has two weeks available at the end of the semester, the Introduction Unit and another unit could be covered, hence giving the students exposure to a different type of mathematics.

The author and her chair decided that Vertex Coloring would interest high school students because of the physical coloring of a graph to reach a solution to a problem. The Minimum Spanning Tree Unit has many algorithms throughout the unit for the students to use to guide them to a solution. Domination is a personal favorite of the chair and the author, thus making it an automatic choice to include. Lastly, the Hamiltonian Unit presents a very popular unsolvable problem to students, but shows them methods to try and solve the problem as best they can.

In all four of the units, an Introduction Section defines new terminology for the particular unit along with examples. An In Class Section follows, which gives the teachers examples for the students to try on their own or together in groups. The Problem Solving Section contains different relevant material for each unit. The next section, the Application Section, shows how each area of Graph Theory can be applied to the real world. Many times students question the importance of mathematics and if they will ever use the information. The author felt that it was important to include this section to show high school students how useful this type of mathematics can

be. They are shown how to solve a number of problems that they as students could possibly face now or as an adult in the real world. Lastly, Homework Exercises with an Answer Key are provided for the teacher.

2 MATHEMATICAL MODELING

The first question that arises from the given theme is “What exactly is mathematical modeling?” According to Niss, mathematical modeling can be defined as “certain objects, relations between them, and structures belonging to the area under consideration are selected and translated into mathematical objects, relations and structures, which are then said to represent the original ones” [8]. In other words, a mathematical model involves a relation between some kind of mathematical object and a situation of a non-mathematical nature. So when a non-mathematical situation presents itself and mathematics is applied, some kind of a mathematical model has to be involved explicitly or implicitly.

To better understand mathematical modeling, let us look at a very simple example. Suppose that a farmer wants to build a rectangular fence with 300 meters of wood, so that the area it encloses is a maximum. The fence only needs to have three sides because the fourth side lies along a river. An approach to solving this question uses mathematical modeling. The students solving this problem should draw a representation of the fence with the two opposite sides of the fence having a length of x and the other side with a length of y because we don't know how small or large to cut the sides of the fence, so we just give the sides arbitrary numbers.

Next, the students should be able to come up with the equation $2x + y = 300$ from the given information. Which results in $y = 300 - 2x$. The students should next remember that the area of a rectangle is $A = xy$. Solving for y we get $A(x) = x(300 - 2x)$. Thus, by using completion of squares, the remaining part of the solution goes as follows:




Figure 1: Fence Problem

$$\begin{aligned}
 A(x) &= -2x^2 + 300x \\
 &= -2(x^2 - 150x) \\
 &= -2(x^2 - 150x + 75^2 - 75^2) \\
 &= -2[(x - 75)^2 - 75^2] \\
 &= -2(x - 75)^2 + 2 * 75^2
 \end{aligned}$$

Therefore, the maximum is adopted at $x = 75$, because we do not want to subtract anything positive. The corresponding value of y is 150.

Understanding exactly what a mathematical model is is just the first step in discovering relations between mathematics and the real world. Ideally, behind every mathematical model there is a modeling process involved. The modeling process can consist of the following six sub-processes [7].

- a.** Formulation of a task (more or less explicit) that guides you to identify the characteristics of the perceived reality that is to be modeled.
- b.** Selection of the relevant objects, relations etc. from the resulting domain of in-

quiry, and idealization of these in order to make possible a mathematical representation.

- c. Translation of these objects and relations from their initial mode of appearance to mathematics.
- d. Use of mathematical methods to achieve mathematical results and conclusions.
- e. Interpretation of these as results and conclusion regarding the initiating domain of inquiry.
- f. Evaluation of the validity of the model by comparison with data (observed or predicted) and/or with knowledge (theoretically based or shared/personal experience based).

Thus, going through the entire process described above is an explanation of mathematical modeling. Please note that the mathematical modeling process does not have to strictly start with step (a) and chronologically move to step (f). Sometimes it will make more sense to go backwards and repeat some of the processes, or to go through all of them several times. In general, mathematical modeling means working with all aspects from the six sub-processes one way or the other.

However, the majority of mathematical modeling done at the middle school or high school level starts out giving the problem already pre-structured. The classical word problem in the secondary school system “is nothing more than a “dressing up” of a purely mathematical problem in the words of a segment of the real world” [11]. In this case the students are just “undressing” the problem and the mathematical modeling process simply consists of this undressing, the use of mathematics and an

easy interpretation. In terms of the steps given in the mathematical process, the students' would only use the sub-processes (c) and (d). The previous fence example is an example of a mathematical modeling problem that only uses the steps (c) and (d). This problem starts with step (c) because the problem is given in a mathematical nature and the students just need to create a mathematical picture to help visualize how to solve the problem. Coming up with the equations, using the completing the square step, and finishing the math of the problem, all involve step (d). The interpretation is not a true interpretation that would happen in step (e) because the values of x and y were the answers to the problem. The students did not have to take those values and interpret them some other way.

Graph Theory is a valuable mathematical modeling tool. The units designed in this thesis incorporate all the steps in the mathematical modeling process in the Application examples. The following model of the Process of Mathematical Modeling is used with the high school students to make it a little easier for them to follow the steps. The students are encouraged to use this process in every application problem to help them solve the problems.

Process of Mathematical Modeling




Figure 2: Process of Mathematical Modeling

Note the following references were used in the development of the following Graph Theory Units for the High School classroom: [10, 5, 4, 3, 9, 1, 2]. Next, we present the actual units designed for use in the high school classroom. Then, we will conclude the thesis with a summary.

3 GRAPH THEORY INTRODUCTION UNIT

3.1 Introduction

A **graph** G is a finite nonempty set of objects called **vertices** together with a set of unordered pairs of distinct vertices called **edges**. We denote the vertex set of a graph G by $V(G)$ and the edge set by $E(G)$. The number of elements in the vertex set of a graph G is called the **order** of G , denoted n , and the number of elements in the edge set of a graph G is called the **size** of G , denoted m . A pair of vertices v_i and v_j in V are **adjacent** if they are connected by an edge; otherwise, v_i and v_j are **nonadjacent**. The **degree of v** , denoted $\deg(v)$, is the number of vertices adjacent to v . Note that a vertex of degree zero is called an **isolated vertex**. The **minimum degree** of G , denoted $\delta(G)$, is the minimum degree among the vertices of G and the **maximum degree** of G , denoted $\Delta(G)$, is the maximum degree among the vertices of G . A vertex u is said to be **connected** to a vertex v in a graph G if there exists a sequence of edges from u to v in G . A graph G is **connected** if every two of its vertices are connected.

3.2 In Class Examples




Figure 3: A graph G

1. List the vertex set and edge set of G .
2. List the degrees of the vertices of G .
3. What is the minimum degree of G ?
4. What is the maximum degree of G ?
5. Is the graph G connected?


Solution

1. $V(G) = \{a, b, c, d, e, f, g\}$ and $E(G) = \{ab, ac, af, bf, be, bc, cd, df\}$
2. The degrees are as follows: $\deg(a) = 3$, $\deg(b) = 4$, $\deg(c) = 3$, $\deg(d) = 2$, $\deg(e) = 1$, $\deg(f) = 3$, and $\deg(g) = 0$.
3. The minimum degree of G is 0.

4. The maximum degree of G is 4.
5. No, the graph G is not connected. The vertex g is not connected to any vertex in the graph.

For a graph G with vertex set $V(G)$ and edge set $E(G)$, we call a graph H a **subgraph** of G if the vertex set $V(H)$ and edge set $E(H)$ are subsets of $V(G)$ and $E(G)$ where each edge $e = uv$ is in $E(H)$, and both u and v are in $V(H)$. We can obtain H from G by deleting edges and/or vertices from G . Note that when you remove each vertex v , all edges incident with v must also be removed.

Example



A graph G




Figure 4: A graph H ; H is a subgraph of G .

A graph is called a **complete** graph if every two of its vertices are adjacent, denoted K_n .

Examples




Figure 5: Complete Graphs

A **path** is graph in which all but two vertices have degree 2, and the other two vertices have degree 1, denoted P_n .

Examples




Figure 6: Paths

A **cycle** is a graph in which every vertex has degree 2, denoted C_n . Note that cycles must have at least three vertices.

Examples





Figure 7: Cycles

3.3 Homework Exercises

Use the following graph to answer the questions.




1. List the vertex set and edge set of G .
2. List the degrees of the vertices of G .
3. What is the minimum degree of G ?
4. What is the maximum degree of G ?
5. Is the graph connected?
6. Give an example of a subgraph of G consisting of four vertices.


Draw the following graphs.

7. P_8
8. K_6
9. C_7

3.4 Answer Key




1. $V(G) = \{a, b, c, d, e, f, g, h, i\}$ and
 $E(G) = \{ab, ai, bc, bh, ch, cd, de, dg, dh, eg, ef, gf, hi\}$
2. The degrees are as follows: $\deg(a) = 2$, $\deg(b) = 3$, $\deg(c) = 3$, $\deg(d) = 4$,
 $\deg(e) = 3$, $\deg(f) = 2$, $\deg(g) = 3$, $\deg(h) = 4$, and $\deg(i) = 2$.
3. The minimum degree of G is 2.
4. The maximum degree of G is 4.
5. Yes, the graph G is connected.
6. A possible subgraph of G . (Note: there can be other correct answers.)




7.



8.



9.



4 VERTEX COLORING UNIT

4.1 Introduction

In this unit we will examine a concept in Graph Theory called **vertex coloring**. This concept can be very useful in real life applications, such as how to manage conflicts of interest. For example, we will later see how graph coloring techniques can be applied to assigning frequencies to radio stations, scheduling club meetings, and coloring the countries of a map. By a **coloring** of a graph G , we mean the assignment of colors (numbers) to the vertices of G , one color to each vertex, so that adjacent vertices are assigned different colors. A k -**coloring** of G is a coloring of G using k colors. For example, Figure 8 shows a 5-coloring of the graph G_1 , as well as a 4-coloring of the graph G_2 .




Figure 8: Vertex Coloring Example





Notice that a graph G with n vertices can always have an n -coloring. But what if

we wanted to determine the smallest number of colors needed for a graph? In other words, we want to find a smallest number k , where $k \leq n$, for which a k -coloring of G exists. Looking back at Figure 8, we can find a 4-coloring, 3-coloring, and a 2-coloring for the graph G_1 . However, there is no k -coloring where $k < 4$ for the graph of G_2 since $G_2 = K_4$ and all edges are present between all vertices.

This brings us to the question of finding the **chromatic number** of a graph G . The chromatic number of a graph G is the minimum value k for which a k -coloring of G exists. The chromatic number of G is denoted $\chi(G)$. Thus, if we look once more at the graphs in Figure 8, we can see that $\chi(G_1) = 2$ and $\chi(G_2) = 4$. Note that we have only given the chromatic numbers of these two graphs, we have not proved that we know the chromatic numbers.

4.2 In Class Exercises


Find the chromatic number of the graphs below.



4.3 Problem Solving


Let us explore further how to show that we have determined the chromatic number, in other words show that we cannot find a smaller number of colors. For example find the chromatic number of H .

H :



(a)

H :



(b)

Figure 9: Chromatic Number Example

We claim that the chromatic number is at most 4 because we can find a 4-coloring. Suppose the vertices of H are labeled as in Figure 9(b). For example, v_1 and v_4 can be colored B-blue, v_2 colored R-red, v_3 colored G-green, and v_5 colored Y-yellow. Thus, we know that the chromatic number of H is at most 4, $\chi(H) \leq 4$. Now, if we can show that there is no 3-coloring of H , then we can without a doubt say that the chromatic number of H is 4. Notice that the vertices v_1 , v_2 , and v_3 form a triangle, therefore three colors are required to color these vertices. Say that we assign color B to v_1 , color R to v_2 , and color G to v_3 . Since v_4 is adjacent to both v_2 and v_3 , we

cannot assign R or G to v_4 . However, v_4 can be colored B. We can now see that v_5 is adjacent to a vertex colored R, a vertex colored B, and a vertex colored G. Thus, we need to introduce a new color to color v_5 , so there exists no 3-coloring of H . Hence, we have shown that $\chi(H) = 4$.

4.4 Applications

Now that we have an understanding of the coloring concept, how can we use vertex coloring in day-to-day situations?

Potential Problems

1. A club scheduling conflict where some students are members of more than one club.
2. A radio station conflict where frequencies would interfere with each other if the stations were too close.
3. A map conflict where two countries need to be painted differently if they share a border.


Let us do an example of Problem 1. Suppose that you were given the responsibility to schedule the meeting times of all the clubs in your High School. The first problem is that some students belong to more than one club, so not all of the clubs can meet on the same day. Secondly, the school does not want to be open every day for after-school clubs. Thus, you need to schedule as few days of the week for the clubs as possible. Below is the list of clubs and club members who belong to more than one club.

Clubs and Members


Clubs	Students in Multi-Clubs
Math Club	Dayna, Dale, Kristy
Debate Club	Kristy, Dayna, Travis
Science Club	Dale
Computer Club	Kristy, Rachel, Travis
Art Club	Rachel, Dale
Spanish Club	Dale

The question is: What is the minimum number of days needed so that no two clubs sharing a member meet on the same day?

Remembering our process of mathematical modeling, we can see that we are given a real-world situation, so now we need to represent our problem with a graph model. Let us have the vertices of our graph represent the different clubs, where two vertices are adjacent if the clubs they represent share a member. For example, the Debate Club would be adjacent to the Math Club because Dayna and Kristy are in both clubs. Here is a mathematical model where MC - Math Club, DC - Debate Club, SC - Science Club, CC - Computer Club, AC - Art Club, and SpC - Spanish Club:



Now we want to use as few colors as possible to color the graph, that is we want to assign a color to each vertex so that any two adjacent vertices have different colors. Let us have the colors represent the days of the week, so Day 1 is color 1, Day 2 is color 2 and so on.



We can see that four colors are needed to color the graph. So, let us interpret our solution in terms of our real-world problem. We can see that four days are needed in order for every club to meet once a week. Note that because of Dale's involvement in so many clubs, four days is the absolute fewest. Here is a chart to see the schedule more clearly.

Day 1	Day 2	Day 3	Day 4
Math Club	Debate Club Spanish Club	Science Club Computer Club	Art Club

Thus, the minimum number of colors used to color the graph gives the minimum

number of days needed to resolve all time conflicts.


Now let us do an example of Problem 2. The Federal Communications Commission (FCC) makes sure that the broadcast from one radio station does not interfere with the broadcast from any other radio station. Assigning an appropriate frequency to each station does this. The FCC requires that stations within transmitting range of each other must use different frequencies. Suppose that the FCC approves a new law where stations within 500 miles of each other must be assigned different frequencies. The locations of the seven stations are given in the grid below with the distances between the stations in miles.

	A	B	C	D	E	F	G
A	-	450	550	700	600	850	900
B	450	-	500	300	250	600	750
C	550	500	-	100	530	800	900
D	700	300	100	-	470	650	700
E	600	250	530	470	-	350	490
F	850	600	800	650	350	-	530
G	900	750	900	700	490	530	-

The FCC wants you to assign a frequency to each station so that no two stations interfere with each other. The FCC also wants you to assign the fewest possible number of frequencies.


Through our process of mathematical modeling, we begin with a real-world situation. Now we need to translate the above grid into a graph model. Since the stations within 500 miles of each other must be assigned different frequencies, we only need to concern ourselves with the stations that are 500 miles or less apart. Suppose the

vertices represent the stations and two vertices are adjacent whenever the stations they represent are 500 miles or less apart.



Now we want to use as few colors as possible to color the graph, that is, we want to assign a color to each vertex so that any two adjacent vertices have different colors.

Let us have the colors represent the different radio frequencies.




We can see that three colors are needed to color the graph. So, let us interpret our solution in terms of our real-world problem. We can see that three radio frequencies are needed so that stations within 500 miles of each other get different frequencies.

Let us conclude this section by doing an example of Problem 3. You may have noticed in your geography or social studies class that maps are always colored so that neighboring countries do not have the same color. This is done so that the countries are easily distinguished and do not blend into each other. Suppose that a mapmaker gives you a map of South America.



In this given real-world situation we can represent our problem with a graph model.

Let each country be a vertex where vertices are adjacent if they share a border.



We can see that four colors are needed to color this map.

4.5 Addendum


Note that as the number of vertices in a graph increases, it will become more difficult to label the vertices. Thus, you might think that a coloring algorithm would be the answer. However, there is no known efficient algorithm for coloring any graph with the fewest number of colors possible. You might try one of the following approaches:

- Color the vertex of highest degree first and keep coloring, trying to color as many vertices with a given color as possible. Make sure that adjacent vertices have different colors.
- Use the Welsh and Powell algorithm.

Welsh and Powell Algorithm


1. Begin by making a list of all the vertices starting with those of highest degree and ending with those of lowest degree.
2. Color the highest uncolored vertex on your list with an unused color.
3. Go down the list coloring as many uncolored vertices with the current color as you can.
4. If all the vertices are now colored, you are finished. If not, go back to Step 2.

However, the Welsh and Powell algorithm does not always give the fewest number of colors. For example look at the graph below. We will first use the Welsh and Powell algorithm, and then color the graph on our own.




Using the Welsh and Powell algorithm three colors will be used.

- Step 1: List the vertices according to degree: A, B, C, D, E, F, G, H and I .
- Step 2: A is at the top of the list. Color A green.
- Step 3: Going down the list, we color B green.
- Step 4: Choose a new color, red, for the highest uncolored vertex, C .
- Step 5: Going down the list, we color E, G, H , and I red.
- Step 6: Choose a new color, blue, for the highest uncolored vertex, D .
- Step 7: Going down the list, we color F blue.



However, we can color the graph with two colors. Let A, D, F , and I be red and C, E, G, H , and B , be blue.







In fact, finding such a method for using an algorithm to find the chromatic number is a famous unsolved problem! [3]

4.6 Homework Exercises

Mathematical Computations


1. Find the chromatic number for each of the graphs below.



2. (a) Draw a connected graph that has five vertices and a chromatic number of four.

(b) Draw a connected graph that has five vertices and a chromatic number of two.

3. Color the following a map using only three colors.



Application Computations

Note: In the following problems, remember to use the Process of Mathematical Modeling when possible!

1. Below is a list of chemicals together with a list of other chemicals with which each cannot be stored.

Chemicals	Cannot Be Stored With
1	2,5,7
2	1,3,5,4
3	2,4,6
4	2,3,7
5	1,2,6,7
6	5,3
7	1,4,5

How many different storage facilities are necessary in order to keep all seven chemicals?

2. A local zoo wants to take visitors on animal feeding tours. They offer the following tours:

Tour 1: Visit lions, elephants, and giraffes.

Tour 2: Visit monkeys, hippos, and flamingos.

Tour 3: Visit elephants, flamingos, and bears.

Tour 4: Visit hippos, reptiles, and bears.

Tour 5: Visit kangaroos, monkeys, and reptiles.

The animals should not be fed more than once a day. Also, there is only room for one tour group at a time at any one site. Can these tours be scheduled using only Monday, Wednesday, and Friday?

3. Draw graphs to represent the maps below. Color the graphs and find the minimum number of colors needed to color each map.



(a)




(b)

Problem Solving Computations


1. Prove that $\chi(G_1) = 2$ for the graph G_1 of Figure 6.
2. Prove that $\chi(G_2) = 4$ for the graph G_2 of Figure 6.

4.7 Answer Key


In Class Exercises (see page 29)



The chromatic number is 2. We know that we need at most two colors because of the given coloring, and because we can not color the graph with one color because there is an edge (K_2).




The chromatic number is 3. We know that we need at most three colors to color the graph of the given coloring, and because of the odd cycle C_5 subgraph.



(c)

The chromatic number is 4. We know that we need at most four colors to color the graph because of the given coloring, and because the graph contains a K_4 subgraph.




(d)

The chromatic number is 3. We know that we need at most three colors to color the graph because of the given coloring, and because there is a K_3 subgraph.


Homework Exercises

Mathematical Computations (see page 42)


1.




The chromatic number is 2.



The chromatic number is 3.




The chromatic number is 3.




The chromatic number is 3.


2. (a) A possible connected graph with 5 vertices and chromatic number 4.




(b) A possible connected graph with 5 vertices and chromatic number 2.



3. A possible coloring of the map using only 3 colors.




1. Let us represent our problem with a graph model. Let us have the vertices represent our chemicals where two vertices are connected if the chemicals cannot be stored together. The colors can represent the different storage facilities.




We can see that three colors are needed to color the graph. Now let us interpret our solution in terms of our real-world problem. Three different storage facilities are needed in order to keep all seven chemicals. One storage facility can hold chemicals 1 and 4, another can hold chemicals 2, 6, and 7, and the last facility can hold chemicals 3 and 5.

2. Let us represent our problem with a graph model. Let us have the vertices represent the different tours where two vertices are connected if the tours visit a common animal. The colors can represent the days of the week.




We can see that three colors are needed to color the graph. Now let us interpret our solution in terms of our real-world problem. Three days of the week are needed in order to schedule the different tour groups. A possible schedule could consist of Monday's visitors going on Tour 1 and Tour 4, Wednesday's visitors going on Tour 3 and Tour 5, and Friday's visitors going on Tour 2.

3. (a) Let us represent our problem with a graph model. Let us have the vertices represent the states where two vertices are connected if they share a border.




We can see that four colors are needed to color this map.

- (b) Let us represent our problem with a graph model. Let us have the vertices represent the states where two vertices are connected if they share a border.




Problem Solving Computations (see page 47)

1.



We claim that the chromatic number is at most 2 because we can find a 2-coloring. Let v_1 be B-blue and $v_2, v_3, v_4, v_5,$ and v_6 be R-red. Hence, we know that $\chi(G) \leq 2$. Since v_1 is adjacent to every vertex in G a second color would need to be introduced to color any of the vertices left in G . Thus, there exists no 1-coloring of G . So, $\chi(G) = 2$.

2.



We claim that the chromatic number of G is at most 4 because we can find a 4-coloring. Let v_1 be B-blue, v_2 be R-red, v_3 be G-green, and v_4 be Y-yellow. Hence we know that $\chi(G) \leq 4$. Since the vertices v_1, v_2 and v_3 form a triangle, three colors are required to color these vertices. Say that we assign color B to v_1 , color R to v_2 , and color G to v_3 . However, v_4 is adjacent to a vertex colored B, a vertex colored R, and a vertex colored G. Thus, we would need to introduce a new color to color v_4 . Therefore, there exists no 3-coloring of G . So, $\chi(G_2) = 4$.

5 MINIMUM SPANNING TREE UNIT

5.1 Introduction

In this unit we will examine a concept in Graph Theory called **minimum spanning trees**. This concept can be very useful in learning how to find the best network. We will later see how minimum spanning trees can be applied to optimizing a computer network, optimizing a road network, and optimizing cost. But first we need to define what a tree is and define its properties. The definition of a **walk** is the course taken from one vertex to another vertex along edges of a graph. A **path** is a walk in which no vertex nor edge is repeated. Hence, in a **cycle**, or a closed path, no edges may be repeated and only the beginning and ending vertices may be the same. A **tree** is a connected graph with no cycles. For example, which graphs in Figure 10 are trees and why?




Figure 10: Trees

We can see that Figure 10a is a tree because it is a connected graph with no cycles. Figure 10b is not a tree because it has a cycle, and Figure 10c is not a tree because it is not connected.

This now brings us to define a spanning tree. A **spanning tree** of a connected graph G is a tree that is a subgraph of G and contains every vertex of G .

5.2 In Class Exercises

How many different spanning trees can you find for the graph below?






Figure 11: Spanning Trees

To get you started, we will walk through the construction of two spanning trees. For our first spanning tree, let us begin at vertex b . Do not forget that we want to connect all the vertices in the graph such that we have no cycles, i.e., using the smallest number of edges possible. For this spanning tree, we will connect b to a and then a to e , creating a ba edge and an ae edge. Now we have another choice: we can either take the ed edge, or the ef edge. Let us take the ef edge. In order to include the vertices g and h , both the fg edge and the fh edge must be included. Thus, the only vertices that are not in our spanning tree thus far are d and c . We can connect d by choosing the db edge or the de edge. Let us take the de edge. To connect c we can choose the cb edge. Let us take the cd edge; hence we have constructed our first spanning tree with 7 edges. Note that this spanning tree is not unique. Our choices of edges determined it.



Now let us construct another spanning tree. This time we will begin with vertex c . Observe that we can either take the cb edge or the cd edge. For this spanning tree, let us take the cd edge. We are now left with two choices for d , either the de edge or the db edge, let us take the db edge. Next we will take the ba edge followed by the ae edge. Notice that we took the ba edge because c is already in our spanning tree, so the only vertex adjacent to b in our original graph is a , thus resulting in the ba edge choice. Now, the remaining vertices that need to be connected to our spanning tree are f, g , and h . Thus, we can add the ef edge, fg edge, and the fh edge and our second spanning tree with 7 edges is constructed.



See if you can find the remaining 9 spanning trees for the graph in Figure 11!

5.3 Problem Solving

Spanning Trees


From the previous example, we can see that it is not always easy to find a spanning tree, especially for a large graph. Thankfully, we have a couple of algorithms to help us find a spanning tree for a graph if one exists. The first method is called the **breadth-first search algorithm**.

Breadth-First Algorithm for Finding Spanning Trees

1. Pick a starting vertex, S , and label it with a 0.
2. Find all vertices that are adjacent to S and label them with a 1.
3. For each vertex labeled with a 1, find an edge that connects it with the vertex labeled 0. Darken those edges.
4. Look for unlabeled vertices adjacent to those with the label 1 and label them 2. For each vertex labeled 2, find an edge that connects it with a vertex labeled 1. Darken that edge. If more than one edge exists, choose one arbitrarily.
5. Continue this process until there are no more unlabeled vertices adjacent to labeled ones. If not all of the vertices of the graph are labeled, then a spanning tree for the graph does not exist. If all vertices are labeled, the vertices and darkened edges are a spanning tree of the graph.

Example using the Breadth-First Algorithm

Let us return to Figure 11 and apply the Breadth-First Algorithm. Under Step 1, we can pick vertex a to be our starting vertex, label it 0. For Step 2, we can see that the only vertices adjacent to a are e and b , so we can label them with a 1. Step 3 tells us to darken the ae edge and the ab edge. Through Step 4 we will label d , c , and f with a 2 because they are adjacent to vertices labeled as a 1, e and b . So we can darken the ef edge and the bc edge. Now we have a choice, d is adjacent to e and b , which are both labeled with a 1, so we can either darken the ed edge or the bd edge. Let us choose the ed edge. Under Step 5, label g and h with a 3 and darken the fg edge and the fh edge, thus constructing a spanning tree. Note, this spanning tree should appear as one of your 11 spanning trees from the in-class example. If you apply the same steps with a different vertex, or choose a different edge when you have a choice, you will construct a different spanning tree.



Minimum Spanning Tree

In applications involving spanning trees, there are sometimes numbers called **weights**, associated with each edge of a graph. Thus, a **minimum spanning tree** is a spanning tree for the graph for which the total of the weights in the tree is minimum. Note that a graph can have more than one minimum spanning tree, but all of the minimum spanning trees must have the same minimum weight sum. Let us now see how we can construct a minimum spanning tree from a graph.

We will take the same graph in Figure 11 and add weights to the edges.




Figure 12: Minimum Spanning Trees

One algorithm for finding a minimum spanning tree for a graph is known as **Prim's Algorithm**.


Prim's Minimum Spanning Tree Algorithm

1. Find the edge with the smallest weights in the graph. Darken it and circle its two vertices. Ties are broken arbitrarily.
2. Find the edge with the smallest weight from the remaining undarkened edges having one circled vertex and one uncircled vertex. Darken this edge and circle its uncircled vertex.
3. Repeat Step 2 until all vertices are circles.

Example using Prim's Algorithm

Let us look at Figure 12 and apply Prim's Algorithm. Under Step 1, we can see that there are two edges that have the smallest weight; the ef edge and the ab edge both have a weight of 1. We will decide to use the ef edge, hence darken the ef edge and circle f and e . Under Step 2, we can see that the ae edge with weight 2, $ae - 2$, is the smallest weight among the remaining undarkened edges have one circled vertex and one uncircled vertex. Therefore, we can darken the ae edge and circle a . Step 3 tells us to repeat Step 2, so next we can darken the $ab - 1$ edge and circle b . Now darken the $bc - 2$ edge and circle c . Notice that we are now given a choice, the $fg - 3$ edge, $fh - 3$ edge, and the $cd - 3$ edge are all possible choices with the same weight. It does not matter which order you decide to add them to our construction of a minimum spanning tree. Let us darken the cd edge and circle d , then the fg edge and circle g , followed by the fh edge and circle h . All of the vertices have been circled and are connected, thus we have a minimum spanning tree with a minimal weight of $3 + 3 +$

$$1 + 2 + 1 + 2 + 3 = 15.$$




Another algorithm for finding a minimum spanning tree for a graph is known as **Kruskal's Algorithm**.

Kruskal's Minimum Spanning Tree Algorithm

1. Examine the graph. If it is not connected, there will be no minimum spanning tree.
2. List the edges in order from the smallest weight to the largest weight. Ties are broken arbitrarily.
3. Darken the first edge on the list.
4. Select the next edge on the list. If it does not form a cycle with the darkened edges, darken it.
5. For a graph with n vertices, continue Step 4 until $n - 1$ edges of the graph have been darkened. The vertices and the darkened edges are a minimum spanning tree for the graph.

Example using Kruskal's Algorithm

Let us return to Figure 12 and apply Kruskal's Algorithm. Step 1 tells us to make sure that our graph is connected, which we can easily see. Under Step 2 we can create a list: $ef - 1, ab - 1, ae - 2, bc - 2, cd - 3, fg - 3, fh - 3, ed - 4$, and $bd - 5$. Note, for the edges that have the same weight, it does not matter which edge comes first in your listing. Through Step 3 and 4 we can darken ef, ab, ae, bc, cd, fg , and fh . We do not want to darken the ed edge or the bd edge because we will create a cycle. From Step 5, we can see that we have constructed our minimum spanning tree with a minimal weight of 15 because Figure 12 has 8 vertices, and we have selected 7 edges to be in our minimum spanning tree.



5.4 Applications

Now that we have an understanding of the minimum spanning tree concept, how can we use minimum spanning trees in day-to-day situations?

Potential Problems

1. Optimizing a computer network using the least amount of wire.
2. Optimizing a road network for the least amount of mileage.
3. Optimizing a cell phone network for the least total cost.


Let us do an example of Problem 1. Suppose that at your high school, six computers in six different offices need to be networked. Your school wants the “best” possible network, i.e. use the least amount of wire to link all the computers. Note that the connection between two computers can either be linked directly or indirectly through another computer. The grid below shows which computers can be linked directly as well as how much wire in meters is needed, where the computers are letters and the distances are in meters.

	A	B	C	D	E	F
A	-	9	-	-	-	3
B	9	-	8	-	8	11
C	-	8	-	3	5	-
D	-	-	3	-	6	11
E	-	8	5	6	-	9
F	3	11	-	11	9	-

The question is: What is the minimum amount of wire needed to connect all six


computers so that every computer is linked directly or indirectly to every other computer?

Remembering our process of mathematical modeling, we begin with a given real-world situation, so now we need to represent our problem with a graph model. Let us have the vertices of our graph represent the computers, where two vertices are connected by an edge if the computers have a direct connection. The weights attached to each edge are the distance between the two points on the grid.




Now we can solve our mathematical problem by using Prim's Algorithm or Kruskal's Algorithm to find the minimum spanning tree. Let us decide on Prim's algorithm. Using this algorithm, a possible spanning tree is dc, ce, cb, ba , and the af edge with a minimum weight of 28. Another consists of dc, ce, eb, ef , and the fa edge with a minimum weight of 28. Now let us interpret our solution in terms of our real world problem. We have found two minimum spanning trees, which mean that we have two computer networks that use the minimum amount of wire, 28 meters, to connect all

six computers.




Now we will do an example of Problem 2. Suppose that, in making plans for winter storms, your local county government needs a design for repairing the county roads in case of an emergency. You are given a map of the towns in your county and the existing major roads between them.



Map of your county with mileage

You are then asked to devise a plan that repairs the least number of miles of road but keeps a route open between each pair of towns.

We begin with our real-world situation. For our graph model, we can simply use the given map. Let us solve our mathematical problem this time by using Kruskal's Algorithm to find a minimum spanning tree. This algorithm creates the list: $AI - 5$, $DE - 5$, $AB - 6$, $DC - 7$, $AD - 8$, $BC - 9$, $AE - 9$, $GH - 10$, $GF - 12$, $HI - 13$, $GI - 15$, $GA - 15$, $FE - 15$, $AF - 16$, $GE - 20$. Using this algorithm, a possible minimum spanning tree consists of AI , DE , AB , DC , AD , GH , GF , and the HI edge with a minimum weight of 66. Now let us interpret our solution in terms of our real world problem. Through our minimum spanning tree we have discovered a plan that connects the towns with the minimum possible number of miles of road, 66 miles.




We will now conclude this section by doing an example of Problem 3. Suppose that a family with seven members in different parts of the country has a relative serving overseas. The family wants to set up a cell phone calling network so everyone will know the latest news about the overseas relative, for the least total cost. The grid below shows the cost for a 15-minute phone call between each pair of family members.


	Alice	Faith	Hillard	Kristy	Owen	Peter	Rachel
Alice	-	\$3.50	\$4.75	-	\$4.10	-	\$5.10
Faith	\$3.50	-	-	\$2.50	-	\$4.10	\$3.40
Hillard	\$4.75	-	-	\$2.95	-	\$4.40	-
Kristy	-	\$2.50	\$2.95	-	\$4.25	-	\$3.40
Owen	\$4.10	-	-	\$4.25	-	-	\$3.20
Peter	-	\$4.10	\$4.40	-	-	-	-
Rachel	\$5.10	\$3.40	-	\$3.40	-	-	-

What is the total cost of the least expensive calling network they can set up?

In this given real-world situation we need to represent our problem with a graph model. Let us have the vertices of our graph represent a family member using the first letter of their name, where two vertices are connected by an edge if the family members have a direct connection. The weight attached to each edge is the amount of money between the two callers on the grid.




Let us use Kruskal's algorithm to find a minimum spanning tree. Using this algorithm, a possible spanning tree is KF, KH, FR, FA, FP , and the AO edge with a minimum cost of \$20.55. If we interpret our solution in terms of our real world problem, we can see that Kristy calls Faith and Hillard, Faith calls Rachel, Alice, and Peter, and Alice calls Owen for a total minimum cost of \$20.55.




5.5 Homework Exercises

Mathematical Computations


- Use the breadth-first algorithm to find a spanning tree for this graph. Begin at vertex C .



- For the following graph, first use Prim's Algorithm to find a minimum spanning tree, then use Kruskal's Algorithm to find a minimum spanning tree. What is the minimum weight in each case?



3. Find a minimum spanning tree for this weighted graph using your favorite method.



Application Computations

Note: In the following problems, remember to use the Process of Mathematical Modeling when possible!


1. A local restaurant has opened an outdoor patio for the summer. The owner wants to hang nine festive light fixtures at designated locations on the overhead latticework. Because of the layout of the patio and the latticework, it is not possible to install wiring between every pair of lights. The grid below shows the distances in feet between lights that can be linked directly. The owner wants to use the minimum amount of wire to get all nine lights connected.

	A	B	C	D	E	F	G	H	I
A	-	16	-	-	15	15	-	-	-
B	16	-	16	12	-	-	-	-	-
C	-	16	-	-	-	-	12	-	-
D	-	12	-	-	-	-	10	-	-
E	15	-	-	-	-	7	-	-	-
F	15	-	-	-	7	-	-	-	-
G	-	-	12	10	-	-	-	18	-
H	-	-	-	-	-	-	18	-	8
I	-	-	-	-	-	-	-	8	-

What is the minimum amount of wire needed to connect all nine lights?


2. There are seven small towns in Madison County that are connected to each other by gravel roads, as in the following diagram. The distances are given in miles. The county wants to pave some of the roads so that people can get from town to town on paved roads, either directly or indirectly. However, Madison

County is on a tight budget so the total number of miles paved needs to be minimum.




Find and draw a network of paved roads that will fulfill the county's requirements.

3. The computers in each of the offices at Earl of March High School need to be linked by cable. The map below shows the cost of each link in hundreds of dollars. What is the minimum cost of linking the 14 offices?



Problem Solving Computations

- Suppose that for the graph below the edges represent possible highways that may be built to join pairs of cities. Let us assume that these weights are the projected costs in millions of dollars for the highways. The idea is to build the cheapest highway network to keep the state's budget down.












Now suppose that the governor lives in town k . He expects to make a lot of trips to towns c and l and has used his political clout to force the construction of the direct routes ck and kl even though those highways will be rather expensive. Find a minimum spanning tree for the graph subject to the restriction that edges ck and kl are in the tree. Explain why there can be no spanning tree containing ck and kl that has smaller cost than the tree that you have found.

5.6 Answer Key

In Class Exercises (see page 61)

The remaining 9 spanning trees for the graph in Figure 2.







Homework Exercises

Mathematical Computations (see page 75)


1. A possible tree beginning at vertex C .



2. A possible minimum spanning tree using Prim's algorithm.




A possible minimum spanning tree using Kruskal's algorithm.




The minimum weight using both algorithms is 17.

3. A possible spanning tree using Prim's algorithm.




A possible minimum spanning tree using Kruskal's algorithm.




The minimum weight using both algorithms is 35.

Application Computations (see page 77)


1. Let us represent our problem with a graph model. Let us have the vertices represent our festive light fixtures where two vertices are connected if the lights can be linked directly. The weight attached to each edge is the amount of wire in feet.




Let us use Prim's algorithm to find a minimum spanning tree. Using this algorithm, a possible minimum spanning tree is $FE, FA, AB, BD, DG, GC, GH$, and the HI edge with a minimum weight of 98. If we interpret our solution in terms of our real world problem, through our minimum spanning tree we have discovered a plan that connects all nine lights with the minimum amount of wire, 98 feet.



2. For our graph model, we can simply use the given map. Using Kruskal's algorithm, a possible minimum spanning tree consists of $AB, BD, EF, FB, FG,$ and $DC,$ with a minimal weight of 88. Now let us interpret our solution in terms of our real world problem. Through our minimum spanning tree we have discovered a network of paved roads that connects the towns with the minimum possible number of miles of road, 88 miles.




3. For our graph model we can simply use the given map. Using Kruskal's algorithm a possible minimum spanning tree consists of $MJ, JG, LI, IF, KL, EF, AB, NH, FB, HD, GC, CD,$ and $IJ,$ with a minimal weight of 55. Now let us interpret our solution in terms of our real world problem. Through our minimum spanning tree, we have discovered a plan that connects the 14 offices with the minimum cost of \$5,500.




Problem Solving Computations (see page 79)

1. For our model we can use the map below.



Using Kruskal's algorithm, after including the edges ck and cl first, a possible minimum spanning tree consists of $ck, kl, fb, ba, fc, ae, cg, dh, ei, ad, gm, jk$, and mn , with a minimum weight of 152. Through our minimum spanning tree we have discovered a plan to build the highway network for \$152 million dollars.



There can be no spanning tree containing ck and kl with a smaller cost because we included ck and kl first, and then took the smallest weighted edges out of the remaining edges.

6 DOMINATION UNIT

6.1 Introduction

In this unit we will examine a concept in Graph Theory called **domination**. This concept can be very useful in real life applications. For example, we will later see how domination techniques can be applied in placing a minimum number of security stations in a county, determining the fewest number of stops a bus driver needs to make, and constructing the least amount of radio stations in an area. A set of vertices S in graph G is a **dominating set** of G if every vertex of G is either in S or adjacent to a vertex in S . The **domination number** of G , denoted by $\gamma(G)$, of a graph G is the smallest number of vertices in any dominating set of G . Let us look at Figure 13 and determine the minimum dominating sets and the domination number.





Figure 13: Domination Example

First, we can clearly notice that all the vertices form a dominating set, but we want to find the least number. Notice that we could choose $\{b, d\}$ as a dominating set, since b dominates itself and the vertices adjacent to it, a, c , and e . Next d

dominates itself and its neighbors c and e . Other minimum dominating sets are $\{a, e\}$, $\{a, d\}$, $\{c, e\}$, $\{b, e\}$, $\{b, c\}$, and $\{c, d\}$. So we know that we need at most two vertices to dominate the graph, $\gamma(G) \leq 2$. To see that $\gamma(G) = 2$, we must show that one vertex can not dominate the graph. To see this, note that no one vertex is adjacent to every vertex in the graph. Hence $\gamma(G) = 2$.

6.2 In Class Exercises

Find the domination number of the graphs below.



6.3 Problem Solving

History of Domination

The concept of domination appears to have originated while playing a game of chess, where the idea is to cover or dominate squares of a chessboard with certain chess pieces. In the game of chess, a queen can move horizontally, vertically, or diagonally over any unoccupied squares. For example in Figure 14, the queen can move to (or attack, or dominate) all of the squares marked with an “X”. In 1862, de Jaenisch tried to figure out all of the minimum number of queens that can be placed on a chessboard so that every square is either occupied by a queen or being attacked by at least one queen. His question is now commonly known as **The Five Queens Problem**, since it has been proven that the answer is 5 [10, 5]. One possible solution is shown in Figure 14.




Figure 14: The Five Queens Graph

You might be wondering what the connection is between the above queens problem

and dominating sets in graphs. The connection can be easily seen through a model of this problem. We can let the 64 squares of a chessboard be the vertices of our graph G where two vertices (squares) are adjacent in G if each square can be reached by a queen on the other square in a single move. This graph G , which we have just constructed, is generally referred to as the **Queen's Graph**. Hence, the smallest number of queens that dominate all the squares of a chessboard is the domination number of G , $\gamma(G)$.

6.4 Applications

Now that we have an understanding of the domination concept, how can we use domination in day-to-day situations?

Potential Problems


1. Placing the smallest number of security stations in a county so that every high school is protected.
2. Determining the fewest number of stops a bus driver needs to make.
3. Constructing the least amount of radio stations in an area.

Let us do an example of Problem 1. Suppose that a county contains eight high schools connected by roads. The table below tells us which high schools are connected by a road, where a Y is given if the distance between the two schools is less than five miles. The Board of Education in this county wants to upgrade their security so that each high school either has a security station within their high school or is within 5 miles of a high school that does have one, so that if an emergency arises security can get to the scene quickly. Due to budgetary constraints only a minimum number of security stations can be built.


	A	B	C	D	E	F	G	H
A	-	Y	Y	-	Y	-	-	-
B	Y	-	-	Y	-	Y	-	-
C	Y	-	-	Y	Y	-	-	Y
D	-	Y	Y	-	-	-	-	-
E	Y	-	Y	-	-	Y	-	Y
F	-	Y	-	-	Y	-	Y	-
G	-	-	-	-	-	Y	-	Y
H	-	-	Y	-	Y	-	Y	-

What is the smallest number of stations that must be built? Give a set where these could be placed. In addition, in which high schools should security stations be placed if there is already one at B, the largest high school in the county?


Through our process of mathematical modeling, we begin with a real-world situation. Now we need to translate the above grid into a graph model. Let us have the vertices of our graph represent the different high school, where two vertices are adjacent if the high schools they represent are less than five miles apart.



To answer the first question, we need to find the smallest dominating set of the graph. The dominating set $\{C, F\}$ will dominate G . Vertex C dominates itself, A , D , E , and H . Vertex F dominates B, G, E and itself. Thus $\gamma(G) \leq 2$. To see that $\gamma(G) = 2$, we must show that one vertex can not dominate the graph. Since we do not have a vertex that is adjacent to every vertex in the graph, we know that $\gamma(G) = 2$. So, now let us interpret our solution in terms of our real-world problem. We can see that two security stations, located at the high schools C and F , are needed so that each of the remaining high schools has a security station nearby.




To answer the second question, we are still looking for the smallest dominating set, only this time we must include the vertex B in our dominating set. Subsequently, B , will dominate itself, A, F , and D . To dominate the remaining vertices C, E, H , and G , we can include H in our dominating set. Thus, we have found a second smallest dominating set that contains $\{B, H\}$. Now, let us interpret our solution in terms of our real-world problem. We can see that two security stations, located at the largest high school, B , and one at H , are needed to that each of the remaining high schools have a security station nearby.




Notice that the first smallest dominating set that we found, $\{C, F\}$, is not the only smallest dominating set. The only way to have an algorithm to determine the smallest dominating number of a graph is to try every possible set of vertices to see if it dominates. However, if you have a large graph this will be impossible in a reasonable amount of time, even using the world's fastest computers. Also, please note that if you were forced to include a different vertex from the beginning, say vertex D , the domination set will not be a minimum dominating set. A possible dominating set could be $\{D, E, G\}$ and the domination number is 3, which is not the smallest.

Now let us do an example of Problem 2. Suppose that a school district passed a rule that no child in elementary school shall have to walk more than one block to school or to a school bus stop. Due to increase of the price of gas, the school buses need to minimize their number of stops. Determine the fewest number of stops the bus driver would have to make so that for each intersection in the graph below there is either a bus stop or there is a bus stop (or school) no more than one block away.



We begin with our real-world situation. For our graph model, we can simply use the given map above. We need to find a dominating set that consists of the smallest number of intersections for school bus stops. The domination number is 5 and minimum dominating set consists of the vertices $\{(1,1), (1,5), (2,3), (3,1), (3,5), \text{ and } (4,3)\}$. Now, let us interpret our solution in terms our real-world problem. We can see that five bus stops need to be installed at intersections (1.1), (1.5), (3,1), and (3,5), so that the students are either one block away from the school or one block away from a bus stop.




Let us conclude this section by doing an example of Problem 3. Suppose that we have a collection of small towns along the Appalachian Mountains in the areas of the White Mts., Green Mts., Bershire Hills, Catskill Mts., Blue Ridge Mts., and

Cumberland Plateau. We would like to establish radio stations in some of the towns so that messages can be broadcast to all of the towns in the area. Since each radio station has a limited broadcasting range, fifty miles, we need to use several stations to reach all towns. But, radio stations are costly, so we need to construct as few as possible. The locations of the thirteen towns are given in the grid below with the distances between the town in miles.


	A	B	C	D	E	F	G	H	I	J	K	L	M
A	-	39	60					54	27	54			24
B	39	-	28	32	32				38		44		
C	60	28	-	25						54	25		
D		32	25	-	52							45	
E		32		52	-	15			23			37	
F					15	-	53		26			42	
G						53	-	42	54				
H	54						42	-	53				65
I	27	38			23	26	54	53	-				
J	54		54							-	33		36
K		44	25							33	-		
L				45	37	42						-	
M	24							65		36			-

What is the fewest number of stations that need to be constructed?

Through our process of mathematical modeling, we begin with a real-world situation. Now we need to translate the above grid into a graph model. Since we know that that a radio station has a broadcast range of only fifty miles, we can disregard towns that are more than fifty miles apart. Therefore, we can now have the vertices represent towns and connect two vertices by an edge whenever the towns they represent are 50 miles or less apart. This gives us the graph below.





We want to find a set of the least number of stations which dominate all other vertices. The domination number is 4 and a minimum dominating set consists of $\{B, F, H, J\}$. Now let us interpret our solution in terms of our real-world problem. We only need to construct four radio stations in the towns of $B, F, H,$ and $J,$ so that all of the other towns can be reached.





6.5 Homework Exercises

Mathematical Computations

1. Find the domination number of the graphs below.






Application Computations

Note: In the following problems, remember to use the Process of Mathematical Modeling when possible!


1. Suppose that a company contains eleven offices connected by hallways as indicated in the graph below. The manager of the company wants to install top of the line photocopy machines so that each office has a copier within their office or is near an office that has one. Unfortunately, the company is new and funds are limited, thus only a minimum number of photocopiers can be installed.



Tell the manager the minimum number of photocopier machines that need to be purchased and in which offices to place them.

2. Suppose that a contractor is building a new subdivision. The last decision that the contractor has to make is where to place the waste receptacles. Regrettably, the contractor went over budget building the community center, so not every intersection can have a waste receptacle. The contractor would like for you to determine the number of receptacles that are needed so that, for each

intersection, there is either a receptacle or there is one at an intersection one block away. The following figure is a street grid of the city.



Street grid of the city

3. Suppose that we have a collection of small villages in Alaska. We would like to locate radio stations in some of these villages so that messages can be broadcast to all of the villages in the region. Since each radio station has a limited broadcasting range, fifty miles, we need to use several stations to reach all the villages. The locations of the ten villages are given in the grid below with the distances between the villages in miles.

	A	B	C	D	E	F	G	H	I	J
A		40	45	40					65	
B	40		40		30	30				
C	45	40		40					50	
D	40		40				60			
E		30				30				70
F		30			30		45	45		
G				60		45		45		
H						45	45		50	50
I	65		50					50		50
J					70			50	50	

What is the fewest number of stations that need to be constructed?


Problem Solving Computations

1. Find a dominating set of five queens, all in the same row, for an 8x8 chessboard.
2. Find a dominating set of five queens, where no two queens attack each other, for an 8x8 chessboard.
3. Find a dominating set of five queens, all on the main diagonal, for an 8x8 chessboard.


6.6 Answer Key

In Class Exercises (see page 91)

The domination number of the graphs.




For the above graph (a), the domination number is 3. A possible dominating set is $\{a, c, e\}$, so $\gamma(G) \leq 3$. We know that there is a C_4 subgraph contained in the graph, so automatically you need at least two vertices to dominate it. No matter which two vertices of the cycle you choose, they cannot dominate the entire graph, so $\gamma(G) \geq 3$. Hence, $\gamma(G) = 3$. For the above graph (b), the domination number is 3. A possible dominating set is $\{e, b, j\}$, so $\gamma(G) \leq 3$. Note that there are two copies of the cycle C_5 in the graph. No matter which two vertices are used to dominate one copy of C_5 , they cannot dominate the entire graph, so at least three vertices are needed. Hence $\gamma(G) = 3$.




For the above graph (c), the domination number is 3. A possible dominating set is $\{a, b, c\}$, so $\gamma(G) \leq 3$. We know that G is just a C_8 , so at least three vertices are needed. Hence, $\gamma(G) = 3$. For the above graph (d), the domination number is 4. A possible dominating set is $\{a, f, c, k\}$, so $\gamma(G) \leq 4$. We know that G contains a subgraph of a K_4 , two P_3 s, and a $K_{1,5}$. We know that the domination number of any complete graph is 1. Likewise, it is easy to see that the domination number of any star is 1. However, the placement of the P_3 s in G make it impossible to dominate the entire graph G with only one more additional vertex. Thus, $\gamma(G) \geq 4$. Hence, $\gamma(G) = 4$.

Mathematical Computations

1. Find the domination number of the graphs below.





For the above graph (a), the domination number is 2 and a possible dominating set is $\{a, h\}$. For the above graph (b), the domination number is 2 and a possible dominating set is $\{b, g\}$.




For the above graph (c), the domination number is 4 and a possible dominating set is $\{a, g, l, k\}$. For the above graph (d), the domination number is 2 and a

possible dominating set is $\{g, f\}$.




For the above graph (e), the domination number is 3 and a possible dominating set is $\{d, f, l\}$. For the above graph (f), the domination number is 3 and a possible dominating set is $\{c, e, h\}$.

- Through our process of mathematical modeling, we begin with a real-world situation. Let us have the vertices of our graph represent the different offices, where two vertices are adjacent if the office they represent are connected by a hallway. We need to find the smallest dominating set of the graph. The dominating number for the graph is 4 and a dominating set consists of $\{A, D, F, I\}$. So, now let us interpret our solution in terms of our real-world problem. We can see that four photocopier machines need to be purchased and placed in office A, office D, office F, and office I.




- We begin with our real-world situation. For our graph model, we can simply use the given map below. We need to find a dominating set that consists of the smallest number of intersections for waste receptacles. Our smallest dominating set consists of the vertices $\{(1,1), (1,5), (2,3), (3,1), (3,5), (4,3), (5,1), (5,5), (6,3), (7,1), (7,5)\}$. Now, let us interpret our solution in terms of our real-world problem. We can see that eleven waste receptacles need to be placed at intersections $(1,1), (1,5), (2,3), (3,1), (3,5), (4,3), (5,1), (5,5), (6,3), (7,1), (7,5)$.

(7,1), and (7,5) so that the residents in the subdivision have a receptacle at their intersection or have a receptacle one block away.




- Through our process of mathematical modeling, we begin with a real-world situation. Now we need to translate the given grid into a graph model. Since we know that a radio station has a broadcast range of only fifty miles, we can disregard towns that are more than fifty miles apart. Therefore we can now have the vertices represent towns and connect two vertices by an edge whenever the towns they represent are 50 miles or less apart. This gives us the graph below. We want to find a set of the least number of stations which dominate all other vertices. A possible smallest dominating set consists of $\{B, D, H\}$. Now let us interpret our solution in terms of our real-world problem. We only need to construct three radio stations in the towns of $B, D,$ and $H,$ so that all of the other towns can be reached.




Problem Solving Computations (see page 106)

1. A dominating set of five queens all in the same row.



2. A dominating set of five queens where no two queens attack each other.



3. A dominating set of five queens all on the main diagonal.

Q							
		Q					
				Q			
					Q		
						Q	

7 HAMILTONIAN PATHS AND CYCLES UNIT

7.1 Introduction

In this unit, we will examine a concept in Graph Theory called **Hamiltonian paths and cycles**. This concept can be very useful in real life applications, such as how to solve transportation problems. For example, we will later see how Hamiltonian paths and cycles can be applied to determine tournament rankings and to model a well-known problem called the traveling salesperson problem. First, we need to give definitions for a Hamiltonian path and cycle. A **Hamiltonian path** is a path that visits each vertex of a graph *exactly once*. A **Hamiltonian cycle** is a Hamiltonian path that starts and ends at the same vertex. Try to find a Hamiltonian cycle in each of the graphs below in Figure 15.




Figure 15: Hamiltonian Paths and Cycles


We can see that the graph in Figure 15a has a Hamiltonian cycle. One such cycle can be listed as a, b, e, c, d, f, a , another as d, f, e, c, b, a, d . Notice that it is not required that every edge of the graph be used when visiting each vertex exactly once. The graph in Figure 15b also has a Hamiltonian cycle. One such cycle can be listed as a, b, c, d, e, f, a . The graph in Figure 15c does not have a Hamiltonian cycle.

We can easily observe that both Figure 15a and Figure 15b also have a Hamiltonian

path, because by definition a Hamiltonian cycle is a Hamiltonian path that starts and ends at the same vertex. In the above examples, drop the last vertex listed in the given cycles and a path is then shown, for example Figure 15a has the path, a, d, e, c, d, f . Thus, every graph that has a Hamiltonian cycle has a Hamiltonian path. However, sometimes there are graphs that have a Hamiltonian path but do not have a Hamiltonian cycle. For example, Figure 15c has a Hamiltonian path but no cycle. One such path can be listed as c, a, b, d, e, f .

7.2 In Class Exercises

Determine if the following graphs below contain a Hamiltonian cycle. If the graph does not contain a Hamiltonian cycle, does it contain a Hamiltonian path?



7.3 Problem Solving

Hamiltonian cycles are interesting to mathematicians because there is no straightforward test for determining if a graph has a Hamiltonian cycle. Later we will examine three different approaches for trying to determine if a graph has a Hamiltonian cycle. For now, the different approaches are the best that we can do when trying to find a Hamiltonian cycle. Mathematicians have concluded that finding a general approach that will work for any graph may be impossible.

However, mathematicians have been able to come up with several conditions that guarantee that a given graph has a Hamiltonian cycle. The following theorem guarantees the existence of a Hamiltonian cycle for certain types of graphs.

If a connected graph, G , has n vertices, where $n \geq 3$ and every vertex in G has a degree of at least $\frac{n}{2}$, then G has a Hamiltonian cycle [9].

Let us now go back to Figure 15a and check the degree of each vertex. Since each of the six vertices of the graph has a degree of at least $\frac{6}{2} = 3$, the graph has a Hamiltonian cycle. Thus, before we even tried to find a Hamiltonian cycle on our own, this theorem guarantees that we will find at least one. Unfortunately, this theorem does not tell us how to find an actual cycle in the graph.

If a graph has some vertices with a degree less than $\frac{n}{2}$, then the theorem does not apply. This does not automatically mean that the graph does not have a Hamiltonian cycle; the graph may or may not have a Hamiltonian cycle. Both Figure 15b and

Figure 15c has vertices with a degree less than 3, thus so far no conclusions can be drawn. However, after further inspection we did discover that Figure 15b did have a Hamiltonian cycle and Figure 15c did not.

History of Hamiltonian graphs

In 1857, Sir William Rowan Hamiltonian, a well-known Irish mathematician, invented a game consisting of a regular solid dodecahedron made out of wood and some string. Note that a dodecahedron has 20 vertices, 30 edges and 12 faces, where each face is shaped like a regular pentagon. Hamilton had a peg placed at each vertex to represent a famous city of the time. The object of the game was for the player to travel to each city exactly once; the player starts and ends at the same vertex and can only travel from one city to the next if an edge exists between the vertices. The string was used to visually show the route that the player traveled by having the player wind the string around the pegs as he/she was playing the game. Unfortunately, since the wooden dodecahedron was difficult to travel with, the game was not very popular [4, 9].

Subsequently, Hamilton decided to make another version of the game called the Icosian Game of the Around the World game. He “flattened” the dodecahedron into a vertex-edge graph, and had holes represent the vertices, so that a peg could be placed in the holes and moved around the graph in search of the tour that started and ended at the same vertex. Unfortunately, the game was still not a huge success, probably because this game is not that difficult to solve [4, 9].

Below, in Figure 16, is a graph of the dodecahedron. Play Hamilton’s Icosian game

by finding an around the world tour. Note that the around the world tour is simply a Hamiltonian cycle.




Figure 16: Hamilton's Icosian Game

7.4 Application

Now that we have an understanding of the concept of Hamiltonian paths and cycles, how can we use Hamiltonian paths and cycles in day-to-day situations?

Potential Problems

1. Determine the ranks of a team in a competition where each team plays every other team.
2. The Traveling Salesperson Problem.

Before we jump into looking at an example of Problem 1, we need to first define another graph theory term. We know that it is often useful for the edges of a graph to have direction. Think about a competition where each player plays every other player. We can use graph theory to illustrate this idea. A complete graph, where the vertices represent the players and a directed edge from vertex A to vertex B shows that player A defeats player B , is a type of digraph known as a **tournament**. A remarkable fact about this kind of digraph is that every tournament contains a Hamiltonian path. This means that, at the end of a tournament, it is possible to rank the teams in order from winner to loser.


Now let us do an example of Problem 1. Suppose four soccer teams play in the high school round robin tournament. The matrix below shows the results of the tournament. Give the rankings of the teams in the soccer round robin tournament.

High School Tournament Results

	S	Do	Da	U
Science Hill (S)	0	0	0	1
Dobyns Bennett (Do)	1	0	0	1
Daniel Boone (Da)	1	1	0	1
Unicoi Co (U)	0	0	0	0

Remember that the matrix is read from row to column, with a “1” representing a win. For example, the “1” in the Daniel Boone-Dobyns Bennett entry means that Daniel Boone beat Dobyns Bennett.

Through our process of mathematical modeling, we begin with a given real-world situation. Now we need to translate the above grid into a graph model. Let us have the vertices of our graph represent the different high schools. Using the information of which team won in each meet, a tournament digraph can be constructed.



After examining the digraph, we can see that the only Hamiltonian path is Da-Do-S-U. So, let us interpret our solution in terms of our real-world problem. The Hamiltonian path gives a sequence of teams where each team beats the next. Thus, the Hamiltonian path Da-Do-S-U also serves as a ranking. Daniel Boone High School


placed first, Dobyns Bennett High School placed second, Science Hill third, and Unicoi Co High School last.

Now we will do an example of Problem 2. Suppose that you are a salesperson who lives in Johnson City. You want to travel to several different cities, say Chicago, Atlanta, Washington D.C., exactly once and then return home to Johnson City. The list below represents the trips that are available to you and the costs of making a trip between the cities.


<u>Cities</u>	<u>Cost</u>
Atlanta - Chicago	\$400
Atlanta - Johnson City	\$550
Atlanta - Washington D.C.	\$1,260
Chicago - Johnson City	\$270
Chicago - Washington D.C.	\$910
Johnson City - Washington D.C.	\$670

Since you own your business, it is important to use the least amount of money for your trip. To save money, try to find the least expensive route that begins in Johnson City, visits each of the other cities exactly once, and returns to Johnson City.

Through our process of mathematical modeling, we begin a given real-world situation. Now we need to translate the above list into a graph model. Let us have the vertices of our graph represent the different cities. Two vertices are connected by an edge if a trip can be made between the two cities. The weight attached to each edge is the amount of money it will cost to travel between the two cities.



The first way that we will try to solve the problem of finding minimum total weight is to list every possible cycle, along with its cost. Using a tree diagram like the one below will help us sort out all of the possible cycles.



After looking at the diagram it is easy to see that out of all the possible routes, the

best solution comes from the cycle that consists of JC, A, C, W, JC or the cycle in the reverse order, JC, W, C, A, JC. Therefore the least expensive route that begins in Johnson City, visits each of the other cities exactly once, and returns to Johnson City costs \$2,530, where the trip itinerary goes as follows: Johnson City to Washington D.C. to Chicago to Atlanta back to Johnson City, or the reverse.

The method that we just used is called the **brute-force method**. You might think that, since this method guarantees a solution, we have found a general solution to the Traveling Salesperson Problem. However, we can quickly see that as the number of vertices increases, checking all the possible routes becomes almost impossible! Even with the world's fastest computer, it would take millions of years to compute the weights of every cycle for a graph with 25 vertices. [9] Thus, we have found a method that guarantees a solution but is not very efficient because it can only be used on graphs with a small number of vertices.


Another way that we will try to solve the problem of finding minimum total weight is to begin at a vertex, look for the nearest vertex, move to it, and so on until you complete the cycle. This method is called the **nearest-neighbor method**. Looking back, let us start at JC, and then move to the nearest neighboring vertex, then to the nearest vertex not yet visited, and return to JC when all of the other cities have been visited. In this case the nearest vertex is the vertex that costs the least amount of money. Therefore, the cycle would start at Johnson City, then Chicago, Atlanta, Washington D.C., and then back to Johnson City. So in this case the minimum weight is $270 + 400 + 1,260 + 670 = 2,600$. Thus, the trip would cost your business \$2,600.

But, we already discovered that the cheapest round trip would cost \$2,530. So, even though the nearest-neighbor method gives a very quick solution it will not always give the correct solution. Now you can understand why there is not a general solution for the traveling salesperson problem that will work in all situations. Either the brute-force method is chosen, which guarantees the best route but is prohibitively slow for large graphs or the nearest-neighbor method, which is quick but does not guarantee the best solution.



7.5 Homework Exercises

Mathematical Computations

1. Determine if the following graphs below contain a Hamiltonian cycle. If the graph does not contain a Hamiltonian cycle, does it contain a Hamiltonian path? (Do not forget to use the theorem that we learned about some Hamiltonian graphs!)



2. Find a minimum-weight Hamiltonian cycle that begins and ends at K . First find the minimum-weight Hamiltonian cycle by using the brute-force method, and then use the nearest neighbor method. Does the nearest-neighbor method give the shortest cycle possible?



Application Computations

Note: In the following problems, remember to use the Process of Mathematical Modeling when possible!

1. Suppose four girls play in the Elida High School round robin girls' tennis tournament. The matrix below shows the results of the tournament. Give the rankings of the girls in the round robin tennis tournament.

Elida High School Tournament Results

	A	B	C	D	E
Amy(A)	0	1	1	0	1
Beth(B)	0	0	1	0	0
Cathy(C)	0	0	0	0	0
Dana(D)	1	1	1	0	1
Emily(E)	0	1	1	0	0

2. Suppose that you are a salesperson who lives in Johnson City. You need to travel to several different cities in Tennessee, Knoxville, Chattanooga, and Memphis exactly once and then return home to Johnson City. The list below represents the trips that are available to you and the miles between the cities.

<u>Cities</u>	<u>Miles</u>
Chattanooga - Johnson City	217
Chattanooga - Knoxville	112
Chattanooga - Memphis	344
Johnson City - Knoxville	107
Johnson City - Memphis	497
Knoxville - Memphis	392

Since you are taking the company's car, it is important to travel the minimum amount of miles on your trip. To save mileage, try to find the minimum route

that begins in Johnson City, visits each of the other cities exactly once, and returns to Johnson City.



First find the minimum-weight Hamiltonian route by using the brute-force method, and then use the nearest-neighbor method. Does the nearest-neighbor method give the shortest possible route?

Problem Solving Computations



1. Draw a connected Hamiltonian graph that has five vertices. Label your vertices and provide the Hamiltonian cycle in your graph.
2. Draw a tournament with five players, where player A beats everyone, C beats everyone except A , B is beaten by everyone, and D beats E .
3. Draw a tournament with five players, A, B, C, D , and E , where there is a three-way tie between A, D , and E for first place.

7.6 Answer Key

In Class Exercises (see page 119)




Yes, the graph in (a) contains a Hamiltonian cycle. A possible Hamiltonian cycle is c, a, b, e, d, c . Yes, the graph in (b) contains a Hamiltonian cycle. A possible Hamiltonian cycle is e, d, b, c, f, a, e .



No, the graph in (c) does not contain a Hamiltonian cycle, however it does contain a Hamiltonian path. A possible Hamiltonian path is d, e, b, a, c . Yes, the graph in (d) contains a Hamiltonian cycle. A possible Hamiltonian cycle is b, a, g, f, e, d, c, b .

Hamilton's Icosian Game (see page 122)




A possible around the world tour \ Hamiltonian cycle is

$a, i, g, e, c, d, o, f, q, h, s, j, k, l, t, r, p, n, m, b, a.$


Homework Exercises

Mathematical Computations (see page 130)

1.





For the above graph (a), our theorem tells us that a Hamiltonian cycle exists because there are five vertices and $\frac{5}{2} \approx 3$ (Note that we can only have whole numbers for an answer since it is impossible to have half of a vertex!), and every vertex in graph (a) has a degree of at least three. A possible Hamiltonian cycle is a, c, d, b, e, a . For the above graph (b), we can not apply the theorem that we learned. The graph (b) does not contain a Hamiltonian cycle but does contain a Hamiltonian path. A possible Hamiltonian path is $b, f, j, d, h, g, e, c, a, i$.




For the above graph (c), we cannot apply the theorem that we learned. The graph (c) does not contain a Hamiltonian cycle but does contain a Hamiltonian path. A possible Hamiltonian path is c, g, e, f, d, b, a . For the above graph (d), we cannot apply the theorem that we learned. However the graph (d) does contain a Hamiltonian cycle. A possible Hamiltonian cycle is $d, e, i, h, f, g, a, b, c, j, k, d$.

2.



Using the brute-force method in (a) produces the following tree diagram.




After looking at the tree diagram, it is easy to see that a minimum-weight Hamiltonian cycle consists of K, N, M, L, K, or the reverse order, with a minimum weight of 60.

Using the nearest-neighbor method in (a), the minimum-weight Hamiltonian cycle consists of K, N, M, L, K, with a minimum weight of 60. In graph (a) the nearest-

neighbor method does give the shortest cycle possible.


Using the brute-force method in (b) produces the following tree diagram.



After looking at the tree diagram, it is easy to see that a minimum-weight Hamiltonian cycle consists of K, M, L, N, K, or the reverse order, with a minimum weight of 262.

Using the nearest-neighbor method in (b), the minimum-weight Hamiltonian cycle consists of K, N, L, M, K, with a minimum weight of 318. In graph (b) the nearest-neighbor method does not give the shortest cycle possible.


1. Through our process of mathematical modeling, we begin with a given real-world situation. Now we need to translate the given grid into a graph model. Let us have the vertices of our graph represent the girls playing in the tournament. Using the information of which girl won in each match, a tournament digraph can be constructed.




After examining the digraph, we can see that the only Hamiltonian path is D-A-E-B-C. So, let us interpret our solution in terms of our real-world problem. The Hamiltonian path gives a sequence of girls where each girl beats the next. Thus, the Hamiltonian path D-A-E-B-C also serves as a ranking. Dana placed first, Amy placed second, Emily third, Beth fourth, and Cathy last.

2. Through our process of mathematical modeling, we begin a given real-world situation. Now we need to translate the given list into a graph model. Let us

have the vertices of our graph represent the different cities. Two vertices are connected by an edge if a trip can be made between the two cities. The weight attached to each edge is the distance in miles between the two cities.




The first way that we will try to solve the problem of finding minimum-weight is to list every possible cycle, along with its milage. Using a tree diagram like the one below will help us sort out all of the possible hamiltonian cycles.




After looking at the diagram, it is easy to see that out of all the possible routes, the best solution comes from two different cycles that consists of JC, C, M, K, JC, its reverse order, and the cycle JC, K, C, M, JC plus its reverse order. Therefore, the route with the least amount of milage that begins in Johnson City, visits each of the other cities exactly once, and returns to Johnson City takes 1,060 miles. Trip itinerary goes as follows: Johnson City to Chattanooga to Memphis to Knoxville back to Johnson City, the reverse, or Johnson City to Knoxville to Chattanooga to Memphis back to Johnson City and its reverse.

Problem Solving Computations (see page 134)


1. A connected Hamiltonian graph with five vertices. A Hamiltonian cycle consists of A, E, C, D, B, A .



2. A tournament with five players, where player A beats everyone, C beats everyone but A, B is beaten by everyone, and D beats E .



3. A tournament with five players, A, B, C, D , and E , where there is a three-way ties between A, D , and E for first place. Notice that A, D , and E all need to win three games a piece. A possible tournament to represent this is below. Again, this is not the only answer.



8 SUMMARY

Since the Graph Theory Units were designed to introduce high school students to Graph Theory, the author and her chair thought that it would be appropriate to test the units with a high school class. At the end of the Spring 2004 semester, a local high school teacher had an extra week available at the end of the year and was interested in teaching a unit. (Note that this particular teacher did have a Master's degree with a concentration in Graph Theory. The author and her chair felt that the initial testing of the material would go more smoothly with a teacher who was already familiar with the subject.) The high school teacher decided to teach the Graph Theory Introduction Unit and the Vertex Coloring Unit.

Feedback from the teacher indicated that the class really enjoyed the material. The teacher spent one day on the Introduction Unit, the next day on the computations and applications of vertex coloring, and a third day on the algorithm. (For a teacher not familiar with Graph Theory the author does recommend spending a week on each unit if time permits.) The teacher felt that the unit was easy to follow which allowed for everything to run smoothly. The teacher asked the students to write a written response concerning the units. The students were asked the following questions: 1) What they liked best?, 2) What they liked least?, and 3) Any suggestions to make it better? The majority of the students replied that they liked the actual coloring part of the unit the best. One student replied, "I liked Graph Theory and Vertex Coloring because it takes away the adding and subtracting of math. It's fun and easy and it also can be used to solve real situations that we may come across." Another stated, "I enjoyed the coloring part. It was fun to try different methods to get the

chromatic number. It was also neat to put the graph problems to real life problems.” The majority of the class did not have a part that they did not like. A few replied that the terminology at first was a little confusing but, after working through some examples, it was ok. However one student did say, “I guess didn’t like the fact that there was no set way to do things. I’m used to algebra, where there is 1 way to solve a problem. I didn’t like that you could do everything right and still get the answer wrong. But it didn’t bother me that bad.” Finally, the students said that they would not change anything.

Seeing the positive response that the high school teacher received compelled the author to do more testing of the units. During the Fall 2004 semester, the author was given permission to teach the Vertex Coloring Unit and the Minimum Spanning Tree Unit to her chair’s MATH 5340 Graph Theory and its Applications class. The author wanted to be able to test her material for herself. The MATH 5340 class was informed at the beginning of the class that they were to view this information through the eyes of a high school student. (Due to the fact that the class already knew the basic definitions in Graph Theory, the Introduction Unit was not presented and the author spent one class period on each unit.) After the lecture, the students were instructed to answer the same questions that were given to the high school students. Overall the MATH 5340 students really liked the units. The author received many suggestions and incorporated some of the ideas into the revised units.

From the successful testing of the units with the high school class and the “pretend” high school class, the author hopes that other high school teachers in the local area will be open to the idea of expanding their students’ minds by introducing them

to the wonderful subject known as Graph Theory.

BIBLIOGRAPHY

- [1] A. Coxford, J. Fey, C. Hirsch, H. Schoen, G. Burrill, E. Hart, A. Watkins, M. Messenger, and B. Ritsema, *Course 1 Part A Contemporary Mathematics in Context*, Everyday Learning, Chicago, 1998.
- [2] A. Coxford, J. Fey, C. Hirsch, H. Schoen, G. Burrill, E. Hart, A. Watkins, M. Messenger, and B. Ritsema, *Course 2 Part B Contemporary Mathematics in Context*, Everyday Learning, Chicago, 1998.
- [3] F. Buckley and M. Lewinter, *A Friendly Introduction to Graph Theory*, Pearson Education, New Jersey, 2003.
- [4] G. Chartrand, *Intoductory Graph Theory*, Dover Publications, New York, 1985.
- [5] L. Lesniak and G. Chartrand, *Graphs and Digraphs*, 3rd ed, Chapman and Hall, New York, 1996.
- [6] *Mathematics Curriculum Standards*, Discrete Mathematics with Statistics and Probability, Tennessee Department of Education, viewed 15 January 2004, <<http://www.state.tn.us/education/ci/cimathhighschool/ciescretemath.htm> >.
- [7] M. Blomhoj and T. Hojgaard Jensen, Developing mathematical modeling competence: Conceptual clarificaton and educational planning. *Teaching Mathematics and its applications 22 3* (2003), 123-139.

- [8] M. Niss, Aims and scope of applications and modeling in mathematics curricula. In W. Blum, et al, editors, *Applications and modeling in learning and teaching mathematics*, pages 22-31. Chicester, UK: Horwood Publishing, 1989.
- [9] N. Crisler, P. Fisher, and G. Froelich, *Discrete Mathematics Through Applications*, W.H. Freeman and Company, New York, 1994.
- [10] T.W. Haynes, S.T. Hedetniemi, and P.J. Slater, *Fundamentals of Domination in Graphs*, Marcel Dekker, New York, 1998.
- [11] W. Blum et al., *ICMI-Study 14. Application and modeling in mathematics education* Discussion Document. Special issue of ICMI-Bulletin (2003).

VITA

Dayna Brown Smithers

1810 Lone Oak Road

Johnson City, TN 37604

zdrb20@imail.etsu.edu

Education

- Masters of Science Degree in Mathematics, East Tennessee State University,
May 2005.
- Bachelors of Science Degree in Mathematics, Auburn University,
May 2003.