



GRADUATE SCHOOL
EAST TENNESSEE STATE UNIVERSITY

East Tennessee State University
Digital Commons @ East
Tennessee State University

Electronic Theses and Dissertations

Student Works

8-2019

Barbell Trajectory and Kinematics during Two International Weightlifting Championships

Aaron J. Cunanan
East Tennessee State University

Follow this and additional works at: <https://dc.etsu.edu/etd>



Part of the [Biomechanics Commons](#), and the [Sports Sciences Commons](#)

Recommended Citation

Cunanan, Aaron J., "Barbell Trajectory and Kinematics during Two International Weightlifting Championships" (2019). *Electronic Theses and Dissertations*. Paper 3635. <https://dc.etsu.edu/etd/3635>

This Dissertation - unrestricted is brought to you for free and open access by the Student Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

Barbell Trajectory and Kinematics during Two International Weightlifting Championships

A dissertation

presented to

the faculty of the Department of Sport, Exercise, Recreation, and Kinesiology

East Tennessee State University

In partial fulfillment

of the requirements for the degree

Doctor of Philosophy in Sport Physiology and Performance

by

Aaron J. Cunanan

August 2019

Michael H. Stone, PhD, Chair

Kyle C. Pierce, EdD

Kimitake Sato, PhD

Satoshi Mizuguchi, PhD

Keywords: Biomechanics, Technique Analysis, Snatch

ABSTRACT

Barbell Trajectory and Kinematics during Two International Weightlifting Championships

by

Aaron J. Cunanan

Several methods have been used in the scientific literature to study the weightlifting pull. Broadly, these methods are used to measure kinematic or kinetic variables exhibited by the lifter, the barbell, or the lifter-barbell system. However, there is an apparent disconnect between weightlifting research and coaching practice that may reduce the perceived benefits of technique analysis among coaches and present some challenges for coaches who seek to incorporate technique analysis into their coaching practice. Differences and trends in the technique of competitive weightlifting performances are apparent from the available literature. However, there are also gaps in the literature due to infrequent analyses that are limited to narrow subgroups of the weightlifting population. Therefore, the purposes of this dissertation were to 1) update to the scientific knowledge of weightlifting technique and performance, 2) improve coaches' ability to conduct and interpret technique analysis, and 3) enhance transferability of weightlifting in training to improve sport performance.

A review of methods used to evaluate the weightlifting pull provides some practical guidance for coaches on the application and interpretation of weightlifting technique analysis. Video analysis is recommended as the most practicable method for coaches to implement technique analysis themselves. Methods used to study 319 lifts by women and men from two major international competitions demonstrate the feasibility and usefulness of video analysis as an inexpensive,

time-efficient, and user-friendly method for coaches to conduct reliable technique analysis. The results of this dissertation suggest that a variety of techniques can be used to achieve international weightlifting success and provide some evidence of changes in weightlifting technique since at least the mid-1980's. These results also indicate that a stereotypical technique profile among elite international weightlifters does not exist, which further support the notion that strength is a primary determinant of weightlifting ability.

Copyright 2019 Aaron J. Cunanan

All Rights Reserved

DEDICATION

To my mother, Ellyn, for your fierce compassion.

To my father, Nelson, for your quiet wisdom.

To my wife, Sarah, for your unwavering love, inspiration, and belief.

To my children, Harper, Samson, and Esther, for your past, present, and future.

ACKNOWLEDGEMENTS

I am humbled by my committee's confidence in me to continue their legacy.

Dr. Stone, thank you for allowing us all to stand on your shoulders.

Dr. Pierce, thank you for embodying the spirit of Olympism.

Dr. Sato, thank you for your unfiltered pursuit of excellence.

Dr. Mizuguchi, thank you for your fidelity to doing things the right way.

I would like to especially thank Doc and Coach Stone for their pioneering vision and steadfast commitment to excellence.

Thank you to Grant Gardis for teaching me what it means to be a coach.

I would also like to thank the athletes and coaches of the Olympic Training Site and the ETSU Weightlifting team for allowing me the privilege to be part of their journey.

TABLE OF CONTENTS

	Page
ABSTRACT.....	2
DEDICATION.....	5
ACKNOWLEDGEMENTS.....	6
LIST OF TABLES	9
LIST OF FIGURES	11
Chapter	
1. INTRODUCTION	12
2. METHODS TO EVALUATE THE WEIGHTLIFTING PULL: A PRACTICAL REVIEW	16
Abstract.....	18
Introduction.....	19
Technique Analysis.....	20
Basic Technique of the Weightlifting Pull.....	21
Methods of Analysis	25
Kinematic Analysis.....	25
Three-dimensional Motion Capture	28
Film/Video Analysis	28
Linear Position Transducer	31
Transceiver Systems	32
Accelerometer/Inertial Measurement Unit	32
Kinetic Analysis.....	33
Force Plate	36
Inverse Dynamics.....	36
Electromyography.....	37
Artificial Neural Networks	38

Practical Applications	38
Conclusion	40
References.....	43
3. SURVEY OF BARBELL TRAJECTORY AND KINEMATICS OF THE SNATCH LIFT FROM THE 2015 WORLD AND 2017 PAN-AMERICAN WEIGHTLIFTING CHAMPIONSHIPS	53
Abstract.....	54
Introduction.....	54
Materials and Methods.....	55
Participants.....	55
Data Collection	56
Video Analysis.....	56
Calibration Variability and Inter-Rater Reliability	57
Outcome Variables.....	57
Statistical Analysis.....	59
Results.....	59
Descriptive Analysis	59
Barbell Trajectory	59
Kinematic Variables.....	63
Comparative Analysis.....	70
Women.....	70
Men	72
Discussion.....	74
Conclusions.....	77
References.....	78
4. SUMMARY AND FUTURE DIRECTIONS	82
REFERENCES	84

VITA..... 100

LIST OF TABLES

Table	Page
2.1 Methods of Kinematic Analysis.....	27
2.2 Methods of Kinetic Analysis	35
3.1 Athlete Characteristics for 2015 World Weightlifting Championship A Sessions.....	56
3.2 Athlete Characteristics for 2017 Pan-American Weightlifting Championship A Sessions	56
3.3 Distribution of Barbell Trajectory Type of Each Athlete’s Heaviest Successful Snatch Attempt in A Sessions at 2015 World Weightlifting Championships	60
3.4 Distribution of Barbell Brajectory Type of Each Athlete’s Heaviest Successful Snatch Attempt in A Sessions at 2017 Pan-American Weightlifting Championships	61
3.5 Distribution of Barbell Trajectory Type of Each Top-Three Finisher’s Heaviest Successful Snatch Attempt at 2015 World Weightlifting Championship	62
3.6 Distribution of Barbell Trajectory Type of Each Top-Three Finisher’s Heaviest Successful Snatch Attempt at 2017 Pan-American Weightlifting Championship.....	63
3.7 Load and Kinematic Variables for Heaviest Successful Snatch Attempt for A Session Lifters at 2015 World Weightlifting Championship.....	64
3.8 Load and Kinematic Variables for Heaviest Successful Snatch Attempt for A Session Lifters at 2017 Pan-American Weightlifting Championship.....	65
3.9 Load and Kinematic Variables for Heaviest Successful Snatch Attempt for Top-Three Women in Each Weight Category at 2015 World Weightlifting Championship	66
3.10 Load and Kinematic Variables for Heaviest Successful Snatch Attempt for Top-Three Men in Each Weight Category at 2015 World Weightlifting Championship.....	67

3.11 Load and Kinematic Variables for Heaviest Successful Snatch Attempt for Top-Three Women in Each Weight Category at 2017 Pan-American Weightlifting Championship	68
3.12 Load and Kinematic Variables for Heaviest Successful Snatch Attempt for Top-Three Men in Each Weight Category at 2017 Pan-American Weightlifting Championship..	69
3.13 Results of Omnibus 2x3 Between-Subjects ANOVAs for Women	70
3.14 Results of Omnibus 2x3 Between-Subjects ANOVAs for Men	72

LIST OF FIGURES

Figure	Page
2.1 Barbell velocity-time profiles	22
2.2. Barbell trajectory types	23
2.3 Center of pressure changes during the weightlifting pull	25
2.4 Phases of the snatch lift.....	26
2.5 Potential barbell kinematic variables of interest.....	31
3.1 Barbell trajectory types	58
3.2 Examined barbell kinematic variables	58
3.3 Cohen's d effect size with 95% confidence interval for main effects of competition and placement among top-three women	71
3.4 Cohen's d effect size with 95% confidence interval for main effects of competition and placement among top-three men	73

CHAPTER 1

INTRODUCTION

Weightlifting is a popular international sport consisting of the snatch and the clean and jerk. There is also wide professional interest in the weightlifting movements among coaches due to the demonstrated efficacy of the implementation of weightlifting movements in training to improve sport performance (Chaouachi et al., 2014; Hackett, Davies, Soomro, & Halaki, 2016; Seitz, Trajano, & Haff, 2014; Tricoli, Lamas, Carnevale, & Ugrinowitsch, 2005). While there is an extensive body of research on the technical and biomechanical aspects of weightlifting, it is important to bear in mind that the purpose of technique analysis should be to improve sport performance (Lees, 2002). Indeed, from a practical standpoint, technique analysis can be useful for coaches to devise teaching and coaching frameworks and may also provide rationale to implement the weightlifting movements in training to improve sport performance, weightlifting or otherwise.

However, coaches have largely been restricted from conducting technique analysis due to financial, spatial, and temporal constraints. These barriers may reduce the perceived benefit of technique analysis among coaches and reduce the likelihood of them consulting the scientific literature. Thus, the translational impact of formal technique analysis and research may be diminished. Emerging commercial technologies that seek to leverage the capabilities of personal electronic devices show some promise to enable coaches to conduct extensive technique analysis themselves. However, the historic disconnect between research and practice may present some challenges for coaches who are inexperienced or untrained in conducting technique analysis. These challenges namely include issues of validity, reliability, and data interpretation.

While these issues do exist, one cannot negate the invaluable knowledge owed to the scientific literature. For example, the kinematic and kinetic structure of the lifter, the barbell, and the lifter-barbell system are well characterized. Such analyses often delineate the weightlifting pull into discrete phases based on the position and movement of the barbell and lifter (Akkuş, 2012; Gourgoulis et al., 2002), which can be useful to identify key moments during the lift and to understand relationships between components of the lifter-barbell system throughout the lift. Furthermore, detailed study of barbell trajectory has given rise to a classification scheme for patterns of barbell displacement that aid in the assessment of an individual's technical proficiency (Hiskia, 1997; Vorobyev, 1978).

Several notable differences and trends in these aspects are also apparent from the existing literature. For example, the rate and magnitude of knee joint rotation during different phases differ based on strength and sex (Gourgoulis et al., 2002; Gourgoulis, Aggeloussis, Kalivas, Antoniou, & Mavromatis, 2004; Harbili, 2012; Kauhanen, Häkkinen, & Komi, 1984), and some variants of the velocity-time profile are more likely based on weight category (Hiskia, 1997). Additionally, differences in the relative frequencies of barbell trajectory types are apparent between different periods of competitive weightlifting history. For example, past investigations found the type 1 and 2 trajectories were most prevalent at international competitions during the 1980's with limited observations of the type 3 trajectory (Baumann, Gross, Quade, Galbierz, & Schwirtz, 1988; Garhammer, 1989, 1990). However, Hiskia observed the type 3 trajectory to be most prevalent at several international competitions in the early 1990's (Hiskia, 1997). Subsequent studies of the 2010 World Weightlifting Championship have also found the type 3 trajectory to be more prevalent, even among gold medalists, compared to past studies (Akkuş, 2012; Harbili, 2012). Thus, it appears that weightlifting technique as assessed by barbell

trajectory has changed over time. Given these apparent changes in barbell trajectory, it is important to evaluate if similar trends are present in other technical and biomechanical aspects of weightlifting performance.

Despite several notable studies, analysis of performances during international competitions are not frequently reported and have usually been limited to select groups of competitors (Akkuş, 2012; Baumann et al., 1988; Garhammer, 1981; Garhammer, 1991; Harbili, 2012; Hiskia, 1997; Musser, Garhammer, Rozenek, Crussemeyer, & Vargas, 2014; Stone, O'Bryant, Williams, Johnson, & Pierce, 1998). Changes in technique over time may also occur due to changes in contested weight categories since anthropometric profiles differ based on weight category (Ford, Detterline, Ho, & Cao, 2000; Musser et al., 2014) and differences in anthropometrics contribute to differences in barbell and lifter kinematics (Musser et al., 2014). Thus, the infrequent and narrow analysis of competitive performances presents a gap in the scientific knowledge that may reduce the accuracy of existing technical models of effective weightlifting technique and the generalizability of such models to athletes of different skill, size, sex, or ability.

Existing research provides a rich body of knowledge on weightlifting technique. However, several practical barriers may impede coaches from directly implementing or accessing technique analysis. Furthermore, the cross-sectional nature of most studies on weightlifting technique presents gaps in the literature that may reduce the applicability of research findings over time. These issues highlight that further efforts to bridge the gap between research and practice are warranted and ongoing study of weightlifting technique among a variety of athletes is necessary. It is also apparent that coaches may benefit from additional guidance on incorporating technique analysis into their everyday practice—whether applying

research findings or conducting technique analysis themselves. Therefore, the purposes of this dissertation were to 1) contribute to the scientific knowledge of weightlifting technique and performance, 2) improve coaches' ability to conduct and interpret technique analysis, and 3) enhance transferability of weightlifting in training to improve sport performance.

CHAPTER 2

METHODS TO EVALUATE THE WEIGHTLIFTING PULL: A PRACTICAL REVIEW

Prepared for submission to *Professional Strength and Conditioning*

Methods to evaluate the weightlifting pull: a practical review

Aaron J. Cunanan¹, Satoshi Mizuguchi¹, Kimitake Sato¹, Kyle C. Pierce², and Michael H. Stone¹

¹ Center of Excellence for Sport Science and Coach Education,

Department of Sport, Exercise, Recreation, and Kinesiology

East Tennessee State University

Johnson City, TN 37614

² LSU Shreveport Weightlifting Center for High Performance and Development

Department of Kinesiology and Health Science

Louisiana State University Shreveport

Shreveport, LA 71115

Keywords: biomechanics; kinetics; kinematics; snatch; clean and jerk; technique analysis;
barbell trajectory

Corresponding Author: Aaron J. Cunanan

Email: aaron.cunanan@gmail.com

Phone: +1-317-529-6063

Abstract

Several methods have been used to study the weightlifting pull. Broadly, these methods are used to measure kinematic or kinetic variables exhibited by the lifter, the barbell, or the lifter-barbell system. Coaches should understand the utility of existing methods used to analyze and delineate the WL pull in order to improve their ability to effectively implement weightlifting technique analysis, integrate and apply findings across various methods, and promote evidence-driven practices. Therefore, the objectives of this review are 1) to summarize methods used to analyze the weightlifting pull and its phases and 2) to provide practical guidance for coaches to interpret and apply weightlifting technique analysis.

Introduction

The weightlifting (WL) movements and their derivatives form the basis of training for competitive weightlifters¹ and are commonly implemented in the resistance training of other athletes². Considerable research has elucidated the kinematic and kinetic structure of the WL movements with particular focus directed toward the WL pull and examining various phases within it. The WL pull refers to the portion of the snatch or the clean in which the barbell is raised from the ground to approximately waist height³. Interest in the WL pull derives from application of the theoretical concepts of transfer of training⁴⁻⁶ and specificity^{7, 8} through utilization of WL pulling movements to improve sport performance.

Several methods have been used to study the WL pull. Broadly, these methods are used to measure kinematic or kinetic variables exhibited by the lifter, the barbell, or the lifter-barbell system. Criteria dependent on the selected method are used to delineate phases of the pull, which provide insight on qualitative and quantitative characteristics of WL technique and performance (e.g. joint displacement-time profiles, joint angular velocities). Such analyses can be used to guide exercise prescription in accordance with the principles of specificity and transfer of training.

While an extensive body of research examining WL technique exists, coaches have historically been restricted from conducting instrumented technique analysis of their athletes largely due to financial, temporal, and spatial constraints. This apparent gap between research and practice among coaches may reduce the perceived benefits of formal technique analysis and research. Conversely, burgeoning consumer-based technologies may promote the incorporation of technique analysis into one's coaching practice. However, the validity and reliability of many

of these emerging technologies are not well reported⁹, and lack of training or experience may lead well-intentioned coaches to waste their efforts using errant methods.

Coaches should understand the utility of existing methods used to analyze and delineate the WL pull in order to increase their ability to effectively implement WL technique analysis, integrate and apply findings across various methods, and promote evidence-driven practices. Therefore, the purpose of this review is to improve coaches' understanding of various methods to analyze WL technique and the delineation of phases during the WL pull. The objectives of this review are 1) to summarize methods used to analyze the WL pull and its phases and 2) to provide practical guidance for coaches to interpret and apply WL technique analysis.

Technique Analysis

'Technique analysis' refers specifically to the systematic evaluation of sport skill performance and encompasses both descriptive and analytic goals (e.g. how movements are made, determining a movement's effectiveness or its effect on performance outcomes) ultimately to improve sport performance¹⁰. The kinematic and temporal features of a movement are of primary practical importance for coaches because these are visibly observable and provide a basis for instruction and immediate feedback¹¹. Arend & Higgins¹² further suggest careful examination to identify 'critical features', which they define as the 'parts or phases of the movement [that] can be least modified by the performer in order to achieve the goal'. Additionally, knowledge of internal and external forces and patterns of muscle activation contribute to understanding the demands of a given technique and can inform exercise prescription. Indeed, considerable research has examined all these aspects of the WL pull, and some general features of basic technique of the WL pull are apparent from this literature.

Basic Technique of the Weightlifting Pull

The WL pull consists of two main phases: the first pull and the second pull. The first pull refers to the initial raising of the barbell from the platform to approximately knee height¹³. During this phase, the lifter's knees extend to a first maximum joint angle while maintaining relatively constant hip and torso angles. There is a 'transition phase' between the first and second pulls during which the lifter orients his or her torso vertically while the knees re-bend (the 'double knee bend') prior to rapid, forceful extension of the hips, knees, and ankles during the second pull. The double knee bend positions the lifter to effectively produce vertical force against the ground, increases the mechanical advantage of the hip extensors, positions the knee extensors closer to their optimal length, increases stored elastic energy, and may initiate a myotatic reflex in the knee extensors^{3, 14-16}.

The barbell typically maintains a constantly increasing velocity during the first pull and reaches peak velocity during the second pull (Figure 1); although, peak velocity may occur during the first pull in rare instances^{17, 18}. Barbell velocity may plateau or slightly decrease during the transition phase, which may occur more commonly in taller lifters¹⁷. A velocity decrease during the transition phase is also more likely to occur when there is greater velocity at the end of the first pull relative to peak velocity¹⁸⁻²⁰ and may be influenced by other factors such as strength²¹ and skill^{14, 19, 20}. Thus, while informative, the velocity-time profile alone does not predict successful performance, and optimal profiles may differ individually.




Figure 1. The barbell velocity-time profile may exhibit two maxima, each occurring during the first and second pull respectively, interceded by a plateau or slight decrease in velocity during the transition phase. Alternatively, velocity may rise continuously to a single peak.

Barbell trajectory is one of the greatest indicators of WL pull technique. Vorobyev¹³ identified three basic trajectory types based on the pattern of horizontal displacement relative to the lifter and crossing of a vertical reference line drawn to intersect the center of the barbell at the start of the pull (Figure 2). Both the type 1 and 2 trajectories consist of a ‘toward-away-toward’ pattern with neither crossing the vertical reference line during the first ‘toward’ phase. The type 1 trajectory crosses the vertical reference line during the ‘away’ phase and may or may not cross again during the final ‘toward’ phase. The type 2 trajectory does not cross the vertical reference line at any point during the lift. The type 3 trajectory exhibits an ‘away-toward-away-toward’ pattern. The barbell typically crosses the vertical reference line during the first ‘toward’ phase. If the barbell crosses again during the second ‘away’ phase, there is a possible subsequent third crossing during the final ‘toward’ phase. Notably, Hiskia¹⁷ identified a fourth trajectory type in which the barbell exhibits a ‘toward-away-toward-away-toward’ pattern that does not

necessarily cross the vertical reference line at any instant. However, two variants of the type 4 trajectory may be observed following initial horizontal movement either toward, as during the type 1 or 2 trajectory, or away, as during the type 3 trajectory, from the lifter. As such, the type 4 trajectory may more generally be characterized as exhibiting an interceding ‘away-toward’ phase between the first ‘toward’ phase and final ‘away-toward’ phase.




Figure 2. Barbell trajectory type is determined by the pattern of horizontal displacement relative to the lifter during the first pull and number of crossings during the second pull of a vertical reference line drawn through the center of the barbell prior to lift-off. Redrawn from Vorobyev¹³ and Hiskia¹⁷.

Observations by Hiskia¹⁷ and Stone et al.¹⁸ have revealed that only the type 1 trajectory necessarily crosses the vertical reference line. Vorobyev¹³ initially theorized the type 1 trajectory to be the ideal pattern. However, several investigations have since found the type 2 trajectory to be more common among elite weightlifters during international competition^{14, 17, 20}, leading some authors to suggest the type 2 trajectory to be preferable^{18, 22}. There is commonly a net rearward barbell displacement regardless of trajectory type, which is usually accompanied by the lifter

jumping backward to accommodate the barbell's movement^{14, 18, 20, 23-25}. Patterns of horizontal displacement are likely related to anthropometry²⁶ and influenced by absolute and relative strength levels^{14, 18, 27}; thus, the optimal trajectory pattern may differ individually^{28, 29}.

The lifter experiences shifts in balance due to changes in the lifter-barbell system center of mass (COM), which is influenced by the separate movements of the lifter and barbell COMs. The center of pressure (COP) through the foot begins near the ball or arch of the foot and moves rearward during the first pull, begins to shift anteriorly toward the ball of the foot during the transition, and finally ends in front of the vertical reference line at the end of the second pull³⁰ (Figure 3). Changes in COP location may reflect an athlete's ability to effectively exert force against the ground throughout the pull³¹. It should be noted that COP movement during the first pull is correlated directly to the magnitude and time history of barbell horizontal displacement³⁰. The anterior shift in COP after the first pull may influence the magnitude of the barbell's forward horizontal displacement during the transition phase and second pull³⁰. Furthermore, barbell horizontal displacement affects muscular effort during key positions, such as at the end of the first pull and transition phase^{3, 14, 30}.

While these general features provide a basic template of WL technique, a survey of methods used to study the WL pull may provide additional insights to inform coaching practices. Refer to Bartonietz¹⁹, Baumann et al.¹⁴, Enoka³, Hiskia¹⁷, and Vorobyev¹³ for detailed reviews of technique of the WL pull.




Figure 3. Center of pressure through the foot shifts anteroposteriorly during the weightlifting pull. A, balance location at lift-off; B, balance location at end of first pull; C, balance location at end of second pull. Balance locations may vary between lifts, load, and athlete. Mediolateral position not represented in figure. Redrawn and adapted from Garhammer & Taylor³⁰.

Methods of Analysis

Kinematic analysis

Kinematics is the study of motion without consideration of the forces producing or resulting from motion. Kinematic analysis involves both qualitative and quantitative approaches. These methods are used to measure or calculate variables related to the displacement, velocity, and acceleration of components of the lifter-barbell system (Table 1). Some methods of kinematic analysis have led to the delineation of several phases in addition to the first and second pull, which can be applied to both the snatch and clean. Although nomenclature and definitions may vary slightly across studies, typical phases derived from kinematic analyses are identified by changes in barbell height and direction and magnitude of knee joint rotation^{13, 22, 27, 32} (Figure 4), with some studies also identifying features from the velocity- or acceleration-time profiles that

coincide with the start and end points of some phases¹⁷. Additionally, many studies examine the snatch or clean in their entirety and so include phases that extend beyond the lifter's final extension during the second pull.




Figure 4. Although the snatch is depicted here, phases of both the snatch and clean can be determined by changes in barbell height and magnitude and direction of knee joint rotation. First pull, from lift-off until the first maximum knee extension; transition phase, from the end of the first pull until the second minimum knee angle; second pull, from the end of the transition phase until the second maximum knee angle; turnover phase, from the end of the second pull until maximum barbell height; and catch phase, from maximum barbell height until stabilization in the catch position³².

Table 1

Methods of Kinematic Analysis

Method	Component evaluated	Measurement	Primary measured or calculated variable(s) of interest
3D MOCAP	Lifter, barbell, lifter-barbell system	Lifter: segment linear displacement; joint angular displacement Barbell: linear displacement	Lifter: joint range of motion; segment displacement; segment/joint angular velocity; change in segment angle; center of mass location Barbell: vertical and horizontal displacement; vertical velocity; vertical and resultant acceleration; center of mass location Lifter-barbell system: center of mass location
Film/video analysis	Lifter, barbell	Lifter: joint angle; segment angle relative to vertical or horizontal Barbell: linear displacement	Lifter: joint range of motion; segment displacement; segment/joint angular velocity; change in segment angle Barbell: vertical and horizontal displacement; vertical velocity; vertical and resultant acceleration
LPT	Barbell	Linear displacement	Single LPT: vertical displacement Dual LPT: vertical and horizontal displacement; vertical axis rotation
Accelerometer/IMU	Barbell	Linear acceleration	Vertical and resultant acceleration; vertical velocity

3D MOCAP, three-dimensional motion capture; IMU, inertial measurement unit; LPT, linear position transducer

Three-dimensional motion capture. Three-dimensional (3D) motion capture (MOCAP) typically uses an arrangement of multiple infrared cameras to record movement of reflective markers within an established 3D reference frame. Markers are placed to measure displacement of the barbell and joints or body segments of interest. 3D MOCAP is considered the gold standard of motion analysis with accuracy of some systems reported to be 0.058 to 0.068 mm³³ (95% confidence interval), mean absolute error < 0.5 mm³⁴, and negligible system variability given proper calibration and procedures³⁵. However, 3D MOCAP is uncommon outside of laboratory settings due to the exorbitant cost of equipment and complexity of data collection, processing, and analysis. Despite these practical limitations, the impeccable accuracy of 3D MOCAP in measuring displacement allows for extremely precise calculation of the (linear or angular) velocity or acceleration of the object(s) of interest through the single or double differentiation of the displacement-time data, respectively. Thus, 3D MOCAP allows comprehensive profiling of the displacement, velocity, and acceleration histories of various components of the lifter-barbell system. Additionally, when coupled with direct measurements of force, advanced processing of 3D MOCAP data enables complex biomechanical analysis such as inverse dynamics, which is discussed further in the section ‘Kinetic Analysis’. Markerless systems have recently entered the market but currently lack the accuracy of traditional marker-based systems. Refer to Robertson et al.³⁶ and van der Kruk and Reijne³³ for a more detailed explanation of 3D MOCAP methods and analysis.

Film/video analysis. Film analysis contributed greatly to early insight on WL technique due to its relative low cost and ease of data collection. A significant drawback of film analysis is manual frame-by-frame digitization of the object(s) of interest. The advent of digital video recorders eventually led to the replacement of film with digital video to record lifts. Current

methods of video analysis involve specialized software that can analyze a variety of video file types and perform automatic or semi-automatic object tracking, automatic calculation of variables of interest, and export of data to a spreadsheet for further analysis. Software range in cost from free, open-source computer programs (e.g. Kinovea) and free or low-cost mobile applications (e.g. BarSense, Coach's Eye, Ubersense, WeightLifting Motion) to several thousand USD for a single license for some programs (e.g. Dartfish). Common program features include graphical overlay and annotation to highlight objects or events of interest. In addition, some programs utilize online or cloud-based systems that allow remote recording, analysis, and sharing of video files between athletes and coaches.

These technological advances and the ubiquity of video recorders make video analysis a pragmatic method of technique analysis for both research and coaching purposes. However, camera specifications and arrangement will affect the accuracy, precision, and reliability of measurements and calculations. High frame rates (≥ 100 fps) are required for accurate determination of kinematic variables such as barbell displacement, velocity, and acceleration³⁷. Observation of some component(s) of the lifter-barbell system may be obstructed depending on viewing angle. For example, in a side view to observe barbell trajectory, the plates will obstruct viewing of parts of the lifter's body during portions of the lift. Similarly, viewing angle can influence the accuracy of measurements such as distance or joint angle due to perspective error. Detailed treatment of these and other considerations are presented by Garhammer and Newton³⁷ and Payton³⁸.

The use of a single camera only provides one vantage point, which impacts which component(s) of the lifter-barbell system can be analyzed and permits only two-dimensional (2D) analysis. 2D analysis of linear displacement requires calibration of the software's distance

scale using an object of known length in the same plane of the object(s) of interest. For example, the diameter of the largest plate (45 cm) nearest to the camera can be used to calibrate the distance scale when analyzing barbell displacement from a side-view.

A two-camera setup alleviates many of the limitations of single-camera methods and allows for 3D analysis. Cameras can be arranged to allow unobstructed viewing of the entire lifter-barbell system, and calibration of a 3D reference frame allows for the accurate calculation of linear and angular measurements outside the observation planes of the cameras. Both single and two-camera methods permit detailed analysis of barbell trajectory (Figure 5) and the identification of phases and subphases of the WL pull, such as those in Figure 4, to closely examine the relationships and interactions between components of the lifter-barbell system throughout the lift. However, synchronization of footage from multiple cameras requires additional equipment or data processing. Calibration procedures and data analysis are also more complex compared to typical 2D methods.




Figure 5. Several potential variables of interest can be quantified through video analysis. X_{net} , net horizontal displacement from start position to catch; X_1 , horizontal displacement from start position to end of first pull; X_2 , horizontal displacement from end of first pull to most forward position during second pull; Y_{max} , maximum barbell height; Y_{catch} , height of barbell at catch; Y_{drop} , difference between ($Y_{max} - Y_{catch}$). Modified from Stone et al.¹⁸

Linear position transducer. Linear position transducers (LPTs) determine barbell displacement through movement of a wire that is connected to the sensor and tethered to the barbell. Measurement using a single LPT allows for one-dimensional (1D) analysis of displacement-time data and calculation of velocity and acceleration, which are both vector quantities, through the single and double differentiation of the displacement-time data, respectively. Because an LPT is only capable of measuring in one dimension, measurement error is introduced when the movement includes more than one dimension (e.g. includes both

horizontal and vertical components) or the object's trajectory is not directly in line with the LPT axis. A dual-LPT setup where both LPTs are tethered to the same point on the barbell provides the ability to derive values in two dimensions through triangulation. Additionally, a system using two pairs of LPTs, with one pair attached to each side of the barbell, permits separate 2D analysis of the left and right sides of the barbell, which can differ with respect to vertical and horizontal displacement³⁹⁻⁴¹ or due to rotation around the vertical axis³⁹. Refer to Westenburg et al.⁴² for more information about dual-LPT setup and analysis.

Transceiver systems. This technology was a predecessor to modern microsensors and consumer-based software programs. One example of these systems uses an array of transceivers to transmit infrared signals to and receive ultrasonic signals from a sensor attached to the end of the barbell. Some investigators have used multiple units to analyze barbell motion bilaterally⁴¹. Software then computes the 3D kinematics of the end of the barbell to which the sensor is attached. This technology produces accurate data that allows the determination of barbell linear kinematic variables and the calculation of force and power. Some notable studies using this technology have contributed insight on barbell trajectory and kinematics¹⁸, especially of lifts during international competition¹⁷. However, this type of transceiver system has been supplanted largely by the development of microsensors (e.g. accelerometers, inertial measurement units) and accompanying software and mobile applications. Refer to Hiskia¹⁷ for an example of analysis conducted using a transceiver system.

Accelerometer/inertial measurement unit. An accelerometer is a type of microsensor that is attached or fixed to the barbell and measures changes in acceleration along a given axis. Many available devices are triaxial (i.e. 3D). However, barbell rotation during the lift changes the orientation of the sensor, and the data no longer reflect the linear movement of the barbell.

Inertial measurement units (IMUs) are accelerometers that incorporate gyroscopes to detect and correct for changes in sensor orientation making them more suitable to study the WL pull. Refer to Flores⁴³, Sato⁴⁴, and Wagner⁴⁵ for more information and examples on the use of accelerometers and IMUs to study the WL pull.

These devices yield acceleration-time data that can be analyzed directly and are often used to calculate velocity by single integration. Phases of positive acceleration indicate increasing velocity, while negative acceleration corresponds to decreasing velocity. Additionally, transitions in which the acceleration-time curve crosses zero indicate deflections in velocity. Positive-to-negative transitions of acceleration coincide with moving from a phase of increasing velocity to one of decreasing velocity and vice versa with negative-to-positive transitions of acceleration. Periods of zero acceleration indicate constant velocity. Furthermore, evaluation of component and resultant acceleration-time profiles may provide useful information about the nature of the lifter's application of force to the barbell⁴⁶.

Kinetic analysis

Kinetics is the branch of mechanics dealing with the forces and torques involved in producing or resulting from motion. Kinetic analysis includes both qualitative and quantitative approaches. In addition to the magnitude of involved forces or torques, kinetic variables of interest include impulse, power (expressed as the product of either force or torque multiplied by velocity), and rate of force or torque development (Table 2).

Although kinematics does not involve forces or torques, some kinematic measurements can be used to calculate variables such as force or power. For example, barbell displacement and velocity data can be used to calculate changes in the barbell's potential and kinetic energy (i.e. work done on the barbell), which allows the calculation of instantaneous power or average power

over a given time interval such as during the first or second pull^{19, 47}. Similarly, barbell acceleration can be used to calculate the barbell's force—which is also a vector quantity—using the equation for Newton's second law of motion ($\vec{F} = m \cdot \vec{a}$). This calculated force is reflective of the force imparted by the lifter onto the barbell³. Some of the remaining methods in this section also involve the calculation of force or torque; however, they do primarily include direct measurement of force as part of their calculations.

Table 2

Methods of Kinetic Analysis

Method	Component evaluated	Measurement	Primary measured or calculated variable(s) of interest
Force plate	Lifter-barbell system	Ground reaction force	Vertical and resultant ground reaction force; rate of force development; impulse; center of pressure location
Inverse dynamics	Lifter, barbell, lifter-barbell system	Lifter: segment linear displacement; joint angular displacement Barbell: linear displacement Lifter-barbell system: vertical ground reaction force	Net and component muscle or joint forces or torques; joint power; rate of torque development
EMG	Lifter	Muscle activation (i.e. motor unit action potentials)	EMG signal frequency, amplitude, and rate of rise
ANNs	Predictive model of lifter, barbell, or lifter-barbell system	Model output of kinematic or kinetic data	Model sensitivity or accuracy (typically to assess or predict joint torque, joint angular velocity, joint range of motion)

ANN, artificial neural network; EMG, electromyography

Force plate. The WL pull requires the lifter to produce vertical ground reaction force (GRF_Z) to perform vertical work to lift the barbell. Force plates are used to measure GRF_Z using single or dual-platform arrangements. Most modern force plates contain multiple load cells, making it possible to also determine the COP. Additionally, triaxial force plates allow measurement of the resultant and component force vectors. Refer to Baumann et al.¹⁴, Enoka³, and Robertson et al.³⁶ for more information and examples of force plate analysis of the WL pull.

Unlike phases determined through kinematic analysis, force plate measurements have been used to identify phases based on changes in GRF_Z relative to the system weight (i.e. total weight of the lifter and barbell load). There are two periods during which GRF_Z is greater than the system weight, 'Weighting I' and 'Weighting II', and one 'Unweighting' phase during which GRF_Z is less than the system weight³. Weighting I commences at barbell lift-off and lasts through nearly the entire first pull. Unweighting follows, beginning and ending slightly before the start and finish of the transition phase. Weighting II is the last phase and represents the final positive impulse to accelerate the barbell upward³. The barbell force- and acceleration-time profiles exhibit concordance with these phases³. Furthermore, both the barbell and GRF_Z force-time profiles support the notion that the double knee bend enhances force production during the second pull³. Measurement of the WL pull using force plates reveals kinetic characteristics including the maximum force^{3, 14, 48-50}, rate of force development^{14, 51-53}, and impulse^{3, 48-50, 53} during the WL pull. This information is useful in understanding the kinetic demands of the WL pull and the potential transfer of the WL pull to sport performance.

Inverse dynamics. Inverse dynamics differs from the calculation of force or power using only kinematic data. While inverse dynamics can be used to calculate 2D or 3D muscle forces or joint torques, it requires combined measurement of both kinematic (i.e.

displacement-time) and kinetic data (i.e. GRF) for these calculations. These data are then used to solve for the variable(s) of interest. Potential variables of interest in addition to muscle force⁵⁴ and joint torque^{14, 55-57} include joint power^{55, 58}, rate of torque development⁵⁵, and whole body power^{47, 59-61}. Refer to Kipp et al.⁵⁶ and Robertson et al.³⁶ for further discussion and examples of analysis using inverse dynamics.

Inverse dynamics is valuable to gain insight on the internal forces involved during the WL pull that would otherwise be unobservable. Practically, such information provides knowledge of the nature of muscular effort and power output during the WL pull. For example, different loads are required to produce maximum joint power at the hips, knees, and ankles^{55, 58}. Additionally, inverse dynamics has been used to identify asymmetries between left and right side internal joint loads during the snatch, which coaches may be interested in to inform training and evaluate movement performance⁶².

Electromyography. Although electromyography (EMG) measures muscle activation (i.e. motor unit action potentials), it is often combined with other techniques that directly measure force. Some researchers have demonstrated the ability to predict muscle forces using EMG during a weightlifting movement⁶³. However, while EMG amplitude alone does not necessarily indicate the level of muscle force, the pattern of the EMG signal may indicate changes in activation and motor unit recruitment in muscles of interest throughout the WL pull^{48, 49, 64}. EMG pattern may also indicate differences in skill level⁶⁵. Thus, EMG studies may be used to guide exercise prescription based on patterns of muscle activation and, potentially, muscle force. Additionally, some researchers have reported differences in muscle activation patterns⁶⁶ and EMG signal^{67, 68} as a result of WL training. Refer to Enoka⁶⁹, Häkkinen et al.⁴⁸, and Robertson et al.³⁶ for further details and examples of analysis using EMG.

Artificial neural networks. Artificial neural networks (ANNs) are a type of machine learning used to predict some output(s) of interest. ANNs consist of multiple nodes arranged in an input layer, one or more hidden layers, and an output layer. Information typically flows sequentially from the input layer through nodes in subsequent layers. Connections between nodes in neighboring layers comprise algorithms that use the outputs from preceding nodes to influence the interaction of subsequent nodes within the model. ‘Training’ of ANNs involves running multiple iterations of the model to manually tune each node or algorithm or to allow them to self-tune. Initial efforts to apply these approaches to the WL pull include qualitative assessment of lifter kinematics⁷⁰ and predicting joint torques based on lifter kinematics⁷¹ or barbell mass and displacement data⁷². The development of robust models may allow for the accurate prediction of required muscle forces and joint torques and may contribute to the optimization of barbell and lifter kinematics. Refer to Schöllhorn⁷³ for a review of biomechanical applications of ANNs.

Practical Applications

Personal electronics (e.g. smartphones, tablets, laptops) offer capable platforms for portable or microsensor technologies, mobile applications, and computer programs that broaden the coach’s ability to conduct technique analysis in everyday practice. Online and cloud-based systems provide additional flexibility across multiple devices and platforms. These technologies have the potential to democratize many methods of technique analysis that were previously restricted to laboratory settings. However, coaches must be aware of the potential tradeoff between convenience and a device or software’s validity and reliability^{9, 74, 75}. Furthermore, some devices and software may restrict users from accessing information about how data are processed or variables are calculated, which may complicate interpretation of results. Nevertheless, some methods, such as video analysis, can be implemented immediately with personal electronics many coaches are likely to already

possess and have clear guidelines to improve reliability and minimize error^{37, 38}. Although coach-friendly options have practical benefits, the scientific literature remains indispensable to gain comprehensive knowledge of WL technique. Thus, coaches should have explicit knowledge of how to interpret information collected from technique analysis whether they are conducting technique analysis or consulting the literature.

Kinematic analysis has provided insight into characteristic displacement-, velocity-, and acceleration-time profiles of both the lifter and barbell during the WL pull. Notable differences in these profiles have been observed due to skill level and sex, which further supports the role of kinematic analysis in the qualitative assessment of WL pull technique and to guide coaching practice. For example, women and less-skilled athletes exhibit a slower and shallower double knee bend^{21, 22, 27, 32, 49, 58, 76}, which some authors have attributed to weakness in the hip and thigh musculature^{14, 22, 27, 32, 76}. Thus, coaches may seek to emphasize strength development of the hip and thigh musculature in athletes from these groups. Furthermore, the delineation of phases in the WL pull also allows careful examination of the interactions that occur between components of the lifter-barbell system, especially at key positions or moments during the lift. For example, the pattern and magnitude of horizontal barbell displacement during the first pull may influence the amount of loop during the second pull¹⁸, so coaches may choose to focus on these parameters in an effort to optimize barbell trajectory. It is also apparent that horizontal barbell displacement affects the moment arms at each joint and has a profound effect on muscular demands, especially at the knee and hip joints, and that the lengths of these moment arms should be minimized^{14, 30}.

Kinetic analysis provides important information on the nature of the forces required to perform the WL pull, which can also provide rationale for the implementation of the WL pull in training. Specifically, kinetic analysis can identify the contributions of individual muscle forces and joint torques to produce GRF_Z and barbell or whole body power. Such information

can be used to target joint specific adaptations to maximize joint torque, increase GRFz, or improve power output. Muscle activation patterns from EMG may also help to identify muscles that require special emphasis in training or that may be developed by performing the WL pull. For example, WL training produced different muscle coactivation adaptations compared to traditional resistance training and was associated with greater improvements in jump performance⁶⁶. Furthermore, kinetic analysis in relation to specific kinematic patterns or arrangements may also reveal differences in their effectiveness, which may influence a coach's representation of idealized technique for each athlete or in general. Such analysis may contribute to optimization of barbell and lifter kinematics^{71, 72, 77, 78}, with the development of ANNs being potentially useful in this regard. Coaches may also wish to consider the influence of anthropometry on kinematic and kinetic aspects of WL technique^{26, 79}. Furthermore, the monitoring of both kinematic and kinetic variables may be potentially useful for coaches to identify changes in skill^{21, 23, 58, 62, 65, 70, 78, 80-82}, fatigue^{52, 70, 83}, and adaptation^{19, 21, 49, 66, 84, 85}.

Conclusion

Many methods have been used to conduct technique analysis of the WL pull and have contributed to an extensive knowledge of WL technique to help coaches in the conceptualization of idealized technique, the process of error correction, and the programming of training. However, coaches must consult the scientific literature to attain insight generated by several of these methods because they are generally inaccessible to coaches otherwise. Video analysis and portable or wearable devices are likely the most viable methods for the modern coach to conduct advanced technique analysis. While consumer-based technologies are aimed at providing coaches with detailed kinematic data, some of these devices may be cost prohibitive in many cases, in addition to issues of validity, reliability, and durability.

The ubiquity of digital video cameras and the availability of free or low-cost video analysis software make video analysis a readily accessible method for coaches to employ without the need to purchase additional equipment. Most current personal electronic devices have built-in high-definition video cameras with many capable of recording at 120 fps or greater, making them suitable for quantitative analysis³⁷. Automated enhancement of video frame rate may improve quantitative analysis of videos recorded at < 100 fps^{86, 87}. Common video analysis software features include automated object tracking, graphical overlay, and automated calculation of kinematic variables such as displacement and velocity. These features enable the coach to provide augmented visual feedback and rapid determination of kinematic variables of interest. The use of spreadsheet templates or computer scripts (e.g. R or Python programming languages) can increase the efficiency of more extensive data analysis and visualization. Thus, video analysis is a practicable method for coaches to conduct WL technique analysis.

Coaches must remain cognizant that technique is only one factor that contributes to performance outcomes. In fact, it is not unusual for the highest performers to display apparently 'suboptimal' technique. For example, only 3 out of 7 women snatch gold medalists from the 2010 World Weightlifting Championship displayed either of the 'toward-away-toward' barbell trajectory patterns²². Such observations highlight that coaches must be discerning about which kinematic features of the WL pull they idealize and what aspects of technique they choose to focus on when teaching or coaching. Technique analysis may also be useful to identify kinetic factors associated with producing or resulting from a given technique or movement pattern. Kinematic and kinetic analyses thus provide extensive information to characterize various components of WL pull technique. This information can be beneficial for coaches to determine effective technique, target specific movement or performance outcomes, and, ultimately, improve sport performance.

Acknowledgements

The authors would like to thank the artist Danny Schlag for his illustration of the snatch phases.

References

1. Storey, A and Smith, H K. Unique aspects of competitive weightlifting. *Sports Medicine*, 42 (9): 769-790. 2012.
2. Suchomel, T J, Comfort, P and Lake, J P. Enhancing the force-velocity profile of athletes using weightlifting derivatives. *Strength and Conditioning Journal*, 39 (1): 10-20. 2017.
3. Enoka, R M. The pull in Olympic weightlifting. *Medicine and Science in Sports*, 11 (2): 131-137. 1979.
4. Issurin, V. Training transfer: scientific background and insights for practical application. *Sports Medicine*, 43 (8): 675-694. 2013.
5. Suarez, D G, Wagle, J P, Cunanan, A J, Sausaman, R W and Stone, M H. Dynamic correspondence of resistance training to sport: a brief review. *Strength and Conditioning Journal*, 2018. DOI: 10.1519/SSC.0000000000000458.
6. Verkhoshansky, Y V and Siff, M C. *Supertraining*. Denver, CO: Supertraining Institute. 2009.
7. Bosch, F. *Strength Training and Coordination: an Integrative Approach*. Rotterdam, Netherlands: 2010Publishers. 2015.
8. Stone, M H, Stone, M and Sands, W A. *Principles and Practice of Resistance Training*. Champaign, IL: Human Kinetics. 2007.
9. Peake, J M, Kerr, G and Sullivan, J P. A critical review of consumer wearables, mobile applications, and equipment for providing biofeedback, monitoring stress, and sleep in physically active populations. *Frontiers in Physiology*, 9 1-19. 2018. DOI: 10.3389/fphys.2018.00743.
10. Lees, A. Technique analysis in sports: a critical review. *Journal of Sports Sciences*, 20 (10): 813-828. 2002.

11. Bober, T. Biomechanical aspects of sports techniques. Proceedings of the Seventh International Congress of Biomechanics (A Morecki). VIIB. Warsaw, Poland: University Park Press. 1979. pp 501-510.
12. Arend, S and Higgins, J R. A strategy for the classification, subjective analysis, and observation of human movement. *Journal of Human Movement Studies*, 2 36-52. 1976.
13. Vorobyev, A N. *A Textbook on Weightlifting*. Budapest, Hungary: International Weightlifting Federation. 1978.
14. Baumann, W, Gross, V, Quade, K, Galbierz, P and Schwirtz, A. The snatch technique of world class weightlifters at the 1985 world championships. *International Journal of Sport Biomechanics*, 4 (1): 68-89. 1988.
15. Garhammer, J. Energy flow during Olympic weight lifting. *Medicine and Science in Sports and Exercise*, 14 (5): 353-360. 1982.
16. Smidt, G L. Biomechanical analysis of knee flexion and extension. *Journal of Biomechanics*, 6 79-92. 1973.
17. Hiskia, G. Biomechanical analysis of performance of World and Olympic champions. Proceedings of the Proceedings of Weightlifting Symposium Ancient Olimpia, Greece: International Weightlifting Federation. 1997. pp 137-163.
18. Stone, M H, O'Bryant, H S, Williams, F E, Johnson, R L and Pierce, K C. Analysis of bar paths during the snatch in elite male weightlifters. *Strength and Conditioning Journal*, 20 (4): 30-38. 1998.
19. Bartonietz, K E. Biomechanics of the snatch: Toward a higher training efficiency. *Strength and Conditioning Journal*, 18 (3): 24-31. 1996.

20. Isaka, T, Okada, J and Funato, K. Kinematic analysis of the barbell during the snatch movement of elite Asian weight lifters. *Journal of Applied Biomechanics*, 12 (4): 508-516. 1996.
21. Gourgoulis, V, Aggeloussis, N, Kalivas, V, Antoniou, P and Mavromatis, G. Snatch lift kinematics and bar energetics in male adolescent and adult weightlifters. *Journal of Sports Medicine and Physical Fitness*, 44 (2): 126. 2004.
22. Akkuş, H. Kinematic analysis of the snatch lift with elite female weightlifters during the 2010 World Weightlifting Championship. *Journal of Strength and Conditioning Research*, 26 (4): 897-905. 2012.
23. Garhammer, J. Biomechanical characteristics of the 1978 world weightlifting champions. *International Series on Biomechanics*, 3 300-304. 1981.
24. Gourgoulis, V, Aggeloussis, N, Mavromatis, G and Garas, A. Three-dimensional kinematic analysis of the snatch of elite Greek weightlifters. *Journal of Sports Sciences*, 18 (8): 643-652. 2000.
25. Whitehead, P N, Schilling, B K, Stone, M H, Kilgore, J L and Chiu, L Z. Snatch technique of United States national level weightlifters. *Journal of Strength and Conditioning Research*, 28 (3): 587-591. 2014.
26. Musser, L J, Garhammer, J, Rozenek, R, Crussemeyer, J A and Vargas, E M. Anthropometry and barbell trajectory in the snatch lift for elite women weightlifters. *Journal of Strength and Conditioning Research*, 28 (6): 1636-1648. 2014.
27. Gourgoulis, V, Aggeloussis, N, Antoniou, P, Christoforidis, C, Mavromatis, G and Garas, A. Comparative 3-dimensional kinematic analysis of the snatch technique in elite male and female Greek weightlifters. *Journal of Strength and Conditioning Research*, 16 (3): 359-366. 2002.

28. Garhammer, J. Biomechanical profiles of Olympic weightlifters. *International Journal of Sport Biomechanics*, 1 (2): 122-130. 1985.
29. Garhammer, J. Bar trajectories of world champion male and female weightlifters: coaching applications. Part 1. *International Olympic Lifter*, 10 (5): 7-8. 1989.
30. Garhammer, J and Taylor, L. Center of pressure movements during weightlifting. *Proceedings of the 2nd International Symposium on Biomechanics in Sports* (J Terauds, K Barthels, E Kreighbaum, R Mann and J Crakes). Colorado Springs, USA: ISBS. 1984. pp 279-291.
31. Liu, Y and Chen, W. Foot pressure study during pulling phase of snatch lifting. *Proceedings of the 19th International Symposium on Biomechanics in Sports* (J R Blackwell and R H Sanders). San Francisco, USA: ISBS. 2001. pp 143-146.
32. Harbili, E. A gender-based kinematic and kinetic analysis of the snatch lift in elite weightlifters in 69-kg category. *Journal of Sports Science & Medicine*, 11 (1): 162. 2012.
33. van der Kruk, E and Reijne, M M. Accuracy of human motion captures systems for sport applications; state-of-the-art review. *European Journal of Sport Science*, 18 (6): 806-819. 2018. DOI: 10.1080/17461391.2018.1463397.
34. Merriaux, P, Dupuis, Y, Boutteau, R, Vasseur, P and Savatier, X. A study of Vicon system positioning performance. *Sensors*, 17 1-18. 2017. DOI: 10.3390/s17071591.
35. Gorton III, G E, Herbert, D A and Gannotti, M. Assessment of the kinematic variability among 12 motion analysis laboratories. *Gait & Posture*, 29 398-402. 2009.
36. Robertson, D G E, Caldwell, G E, Hamill, J, Kamen, G and Whittlesey, S N. *Research Methods in Biomechanics*. Champaign, IL: Human Kinetics. 2004.
37. Garhammer, J and Newton, H. Applied video analysis for coaches: Weightlifting examples. *International Journal of Sports Science & Coaching*, 8 (3): 581-594. 2013.

39. Bruenger, A J, Smith, S L, Sands, W A and Leigh, M R. Validation of instrumentation to monitor dynamic performance of Olympic weightlifters. *Journal of Strength and Conditioning Research*, 21 (2): 492-499. 2007.
40. Chiu, L Z F, Schilling, B K, Fry, A C and Salem, G J. The influence of deformation on barbell mechanics during the clean pull. *Sports Biomechanics*, 7 (2): 260-273. 2008.
41. Rossi, S J, Buford, T W, Smith, D B, Kennel, R, Haff, E E and Haff, G G. Bilateral comparison of barbell kinetics and kinematics during a weightlifting competition. *International Journal of Sports Physiology and Performance*, 2 (2): 150-158. 2007.
43. Flores, F J, Sedano, S, de Benito, A M and Redondo, J C. Validity and reliability of a 3-axis accelerometer for measuring weightlifting movements. *International Journal of Sports Science & Coaching*, 11 (6): 872-879. 2016.
44. Sato, K, Smith, S L and Sands, W A. Validation of an accelerometer for measuring sport performance. *Journal of Strength and Conditioning Research*, 23 (1): 341-347. 2009.
45. Wagner, J F. About motion measurement in sports based on gyroscopes and accelerometers - an engineering point of view. *Gyroscopy and Navigation*, 9 (1): 1-18. 2018.
46. Gourgoulis, V, Aggeloussis, N, Garas, A and Mavromatis, G. Unsuccessful vs. successful performance in snatch lifts: a kinematic approach. *Journal of Strength and Conditioning Research*, 23 (2): 486-494. 2009.
47. Garhammer, I. A Review of Power Output Studies of Olympic and Powerlifting: Methodology, Performance. *Journal of Strength and Conditioning Research*, 7 (2): 76-89. 1993.
48. Hakkinen, K and Kauhanen, H. A biomechanical analysis of selected assistant exercises of weightlifting. *Journal of Human Movement Studies*, 12 (6): 271-288. 1986.

49. Kauhanen, H, Häkkinen, K and Komi, P V. A biomechanical analysis of the snatch and clean & jerk techniques of Finish elite and district level weightlifters. *Scandinavian Journal of Sports Sciences*, 6 47-56. 1984.
50. Grieve, D W. The defeat of gravity in weightlifting. *British Journal of Sports Medicine*, 5 37-41. 1970.
51. Haff, G G, Carlock, J M, Hartman, M J and Kilgore, J L. Force-time curve characteristics of dynamic and isometric muscle actions of elite women olympic weightlifters. *Journal of Strength and Conditioning Research*, 19 (4): 741. 2005.
52. Haff, G G, Jackson, J R, Kawamori, N, Carlock, J M, Hartman, M J, Kilgore, J L, Morris, R T, Ramsey, M W, Sands, W A and Stone, M H. Force-time curve characteristics and hormonal alterations during an eleven-week training period in elite women weightlifters. *Journal of Strength and Conditioning Research*, 22 (2): 433-446. 2008.
53. Garhammer, J and Gregor, R. Propulsion forces as a function of intensity for weightlifting and vertical jumping. *Journal of Applied Sport Science Research*, 6 (3): 129-34. 1992.
54. Kipp, K and Harris, C. Muscle-specific effective mechanical advantage and joint impulse in weightlifting. *Journal of Strength and Conditioning Research*, 31 (7): 1905-1910. 2017.
55. Kipp, K, Harris, C and Sabick, M B. Lower extremity biomechanics during weightlifting exercise vary across joint and load. *Journal of Strength and Conditioning Research*, 25 (5): 1229-1234. 2011.
56. Kipp, K, Redden, J, Sabick, M and Harris, C. Kinematic and kinetic synergies of the lower extremities during the pull in Olympic weightlifting. *Journal of Applied Biomechanics*, 28 (3): 271-278. 2012.

57. Kipp, K, Redden, J, Sabick, M B and Harris, C. Weightlifting performance is related to kinematic and kinetic patterns of the hip and knee joints. *Journal of Strength and Conditioning Research*, 26 (7): 1838-1844. 2012.
58. Enoka, R M. Load-and skill-related changes in segmental contributions to a weightlifting movement. *Medicine and Science in Sports and Exercise*, 20 (2): 178-187. 1988.
59. Garhammer, J. Power production by Olympic weightlifters. *Medicine and Science in Sports and Exercise*, 12 (1): 54-60. 1980.
60. Lee, S, DeRosia, K, Lamie, L and Levine, N. Correlation profiles between lower extremity net joint torques and whole body power during the power clean. *International Journal of Sports Science*, 7 (2): 66-72. 2017.
61. Lee, S, DeRosia, K D and Lamie, L M. Evaluating the contribution of lower extremity kinetics to whole body power output during the power snatch. *Sports Biomechanics*, 17 (4): 554-556. 2017.
62. Czaplicki, A, Dziewiecki, K, Mazur, Z and Blajer, W. Assessment of internal loads in the joints of the lower extremities during the snatch in young weightlifters. *International Journal of Structural Stability and Dynamics*, 19 (5): 1-19. 2019. DOI: 10.1142/S0219455419410116.
63. Wei, G, Tian, F, Tang, G and Wang, C. A wavelet-based method to predict muscle forces from surface electromyography signals in weightlifting. *Journal of Bionic Engineering*, 9 48-58. 2012.
64. Chen, S K, Wu, M T, Huang, C H, Wu, J H, Guo, L Y and Wu, W L. The analysis of upper limb movement and EMG activation during the snatch under various loading conditions. *Journal of Mechanics in Medicine and Biology*, 13 (1): 1-13. 2013. DOI: 10.1142/S0219519413500103.

65. Enoka, R M. Muscular control of a learned movement: The speed control system hypothesis. *Experimental Brain Research*, 51 (1): 135-145. 1983.
66. Arabatzi, F and Kellis, E. Olympic Weightlifting Training Causes Different Knee Muscle–Coactivation Adaptations Compared with Traditional Weight Training. *Journal of Strength and Conditioning Research*, 26 (8): 2192-2201. 2012.
67. Felici, F, Rosponi, A, Sbriccoli, P, Filligoi, G C, Fattorini, L and Marchetti, M. Linear and non-linear analysis of surface electromyograms in weightlifters. *European Journal of Applied Physiology*, 84 337-342. 2001.
68. Häkkinen, K, Komi, P V, Alén, M and Kauhanen, H. EMG, muscle fibre and force production characteristics during a 1 year training period in elite weight-lifters. *European Journal of Applied Physiology*, 56 419-427. 1987.
69. Enoka, R M. *Neuromechanics of Human Movement*. Champaign, IL: Human Kinetics. 2015.
70. Ibarra, F J R, Encinas, M T S, Gómez, C E R and Garcia, O A P. Design of a biomechanical model and a set of neural networks for monitoring weightlifting. *Research in Computing Science*, 80 31-42. 2014.
71. Salaami, F, Jamshidi, N, Rostami, M and Najarian, S. Power enhancement of weightlifters during snatch through reducing torque on joints by particle swarm optimization. *American Journal of Applied Sciences*, 5 (12): 1670-1675. 2008.
72. Kipp, K, Giordanelli, M and Geiser, C. Predicting net joint moments during a weightlifting exercise with a neural network model. *Journal of Biomechanics*, 74 225-229. 2018.
73. Schöllhorn, W I. Applications of artificial neural nets in clinical biomechanics. *Clinical Biomechanics*, 19 876-898. 2004.

74. Courel-Ibáñez, J, Martínez-Cava, A, Morán-Navarro, R, Escribano-Peñas, P, Chavarren-Cabrero, J, González-Badillo, J J and Pallarés, J G. Reproducibility and repeatability of five different technologies for bar velocity measurement in resistance training. *Annals of Biomedical Engineering*, 1-19. 2019. DOI: 10.1007/s10439-019-02265-6.
75. Adesida, Y, Papi, E and McGregor, A H. Exploring the role of wearable technology in sport kinematics and kinetics: a systematic review. *Sensors*, 19 (7): 1597. 2019. DOI: 10.3390/s19071597.
76. Liu, G, Fekete, G, Yang, H, Ma, J, Sun, D, Mei, Q and Gu, Y. Comparative 3-dimensional kinematic analysis of snatch technique between top-elite and sub-elite male weightlifters in 69-kg category. *Heliyon*, 4 (7): 2018.
77. Ho, L K, Lorenzen, C, Wilson, C J, Saunders, J E and Williams, M D. Reviewing current knowledge in snatch performance and technique: the need for future directions in applied research. *Journal of Strength and Conditioning Research*, 28 (2): 574-586. 2014.
79. Ford, L E, Detterline, A J, Ho, K K and Cao, W. Gender-and height-related limits of muscle strength in world weightlifting champions. *Journal of Applied Physiology*, 89 (3): 1061-1064. 2000.
80. Winchester, J B, Erickson, T M, Blaak, J B and McBride, J M. Changes in bar-path kinematics and kinetics after power-clean training. *Journal of Strength and Conditioning Research*, 19 (1): 177. 2005.
81. Burdett, R G. Biomechanics of the snatch technique of highly skilled and skilled weightlifters. *Research Quarterly for Exercise and Sport*, 53 (3): 193-197. 1982.
82. Carson, H J, Collins, D and Jones, B. A case study of technical change and rehabilitation: intervention design and interdisciplinary team interaction. *International Journal of Sport Psychology*, 45 (1): 57-78. 2014.

84. Bazyler, C D, Mizuguchi, S, Zourdos, M C, Sato, K, Kavanaugh, A A, DeWeese, B H, Breuel, K F and Stone, M H. Characteristics of a National Level Female Weightlifter Peaking for Competition: A Case Study. *Journal of Strength and Conditioning Research*, 32 (1): 3029-3038. 2017.
85. Flores, F J, Sedano, S and Redondo, J C. Optimal load and power spectrum during snatch and clean: differences between international and national weightlifters. *International Journal of Performance Analysis in Sport*, 17 (4): 521-533. 2017.
86. Jiang, H, Sun, D, Jampani, V, Yang, M H, Learned-Miller, E and Kautz, J. Super slomo: high quality estimation of multiple intermediate frames for video interpolation. *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition* Salt Lake City, UT: Computer Vision Foundation. 2018. pp 9000-9008.
87. Srisen, P, Auephanwiriyakul, S, Theera-Umpon, N and Chamnongkich, S. Kinect joints correction using optical flow for weightlifting videos. *Proceedings of the 7th International Conference on Computational Intelligence, Modelling and Simulation* Kuantan, Malaysia: IEEE. 2015. pp 26-31.

CHAPTER 3
SURVEY OF BARBELL TRAJECTORY AND KINEMATICS OF THE SNATCH LIFT
FROM THE 2015 WORLD AND 2017 PAN-AMERICAN WEIGHTLIFTING
CHAMPIONSHIPS

Prepared for submission to *Sports*

Survey of barbell trajectory and kinematics of the snatch lift from the 2015 World and 2017 Pan-American Weightlifting Championships

Aaron J. Cunanan ^{1,*}, Mark A. South ¹, W. Guy Hornsby, III ², Kristina P. Ushakova ¹, Satoshi Mizuguchi ¹, Kimitake Sato ¹, Kyle C. Pierce ³, and Michael H. Stone ¹

¹ Center of Excellence for Sport Science and Coach Education; Department of Sport, Exercise, Recreation, and Kinesiology; East Tennessee State University; 1276 Gilbreath Drive, Johnson City, TN 37614, USA; Affiliation 1; southma@etsu.edu (M.A.S.); ushakovak@etsu.edu (K.P.U.); harahara10@hotmail.com (S.M.); jpnsatok@hotmail.com (K.S.); stonem@etsu.edu (M.H.S.)

² Department of Coaching and Teaching Studies, West Virginia University, 375 Birch Street, Morgantown, WV 26505, USA; william.hornsby@mail.wvu.edu (W.G.H.)

³ LSU Shreveport Weightlifting Center for High Performance and Development; Department of Kinesiology and Health Science; Louisiana State University Shreveport; One University Place, Shreveport, LA 71115, USA; kyle.pierce@lsus.edu (K.C.P.)

* Correspondence: aaron.cunanan@gmail.com (A.J.C.); Tel.: +1-432-439-4265

Received: date; Accepted: date; Published: date

Abstract: Analysis of elite performances are important to elucidate the characteristics of effective weightlifting technique at the highest level of achievement. The general technique of the weightlifting movements is well-established. However, it is also apparent that weightlifting technique can differ based on athlete characteristics. Thus, existing technical models may not accurately reflect current technique of top performers or be generalizable to athletes of different skill, size, sex, or ability. Therefore, the purpose of this study was to update the scientific knowledge of snatch technique of top international weightlifters. This study used video analysis to determine barbell trajectory and kinematics of successful snatch attempts from two major international competitions. Relative frequencies of barbell trajectory types differed based on competition, sex, category, and ranking. No statistical differences were observed among top-three performers of either sex for most kinematic variables, and there were no overall discernible patterns of effect size differences for individual or clusters of kinematic variables. Weightlifting success can be achieved with a variety of technique profiles. Strength is likely a primary determinant of weightlifting performance and ability that may also secondarily allow individuals to overcome suboptimal technique.

Keywords: biomechanics; technique analysis; Olympic-style lifting

1. Introduction

Attempts completed during major international weightlifting competitions comprise maximum or near-maximum performances by the most-skilled performers of the weightlifting movements (the snatch, the clean, and the jerk), and these performances are instructive for the idealization of effective weightlifting technique. The general technique of the weightlifting movements is well-established [1-5]. However, it is also apparent that weightlifting technique can differ based on athlete characteristics. For example, differences in the relative frequencies of barbell trajectory types among lifters in A versus B sessions during international competitions have been observed [4,6]. Several authors have also reported differences in lifter and barbell kinematics and kinetics based on skill level [6-8]. Other investigations of competitive performances provide evidence that weightlifting technique may also

be influenced by other factors such as anthropometry, weight category, and sex [4,6,9-12]. Such findings may provide some guidance to individualize teaching and coaching to best suit each athlete.

Technique differences may also be suggestive of a need for different training objectives or emphases to address deficiencies that may influence weightlifting success or sport performance. For example, some authors have suggested that the ability to execute a stretch-shortening cycle during the transition phase may be improved by increasing knee flexor concentric strength [6,8] or knee extensor eccentric strength [9,11,13]. Additionally, coaches may employ physical training to increase an athlete's speed of moving under the barbell after completing the second pull [7].

It is worth noting that the best performers may exhibit apparently suboptimal technique [9]. However, it is unclear whether top performers with suboptimal technique achieve success because of or despite their technique. Furthermore, the observed technique differences in many of the aforementioned studies are confounded by differences in weightlifting ability (i.e. absolute load lifted), and the technique an individual exhibits is partly dependent on her or his absolute and relative strength. Thus, strength is likely to be a primary determinant of weightlifting success [14].

These technique and strength differences notwithstanding, analyses of performances during major international competitions are infrequent [4,6,9,11,12,15-17] or limited to select rankings or weight categories [9,11]. Thus, existing technical models may not accurately reflect current technique of top performers or be generalizable to athletes of different skill, size, sex, or ability. Accurate technical models are important for coaches to devise frameworks for teaching and coaching the weightlifting movements. Technical and biomechanical analyses can also provide rationale for coaches to implement the weightlifting movements in training on the basis of specificity [18-20] and transference of training [21,22].

Cross-sectional analyses contribute to the cumulative scientific knowledge on weightlifting technique with the potential to inform coaching practices. Analysis of elite performances is important to elucidate the characteristics of effective weightlifting technique at the highest levels of achievement. Additionally, observed technique differences indicate the need for serial investigations encompassing multiple subgroups of performers. Therefore, the purpose of this study was to update the scientific knowledge of snatch technique of top international weightlifters. The primary aim of this study was to elucidate technical and biomechanical parameters of successful snatch attempts by lifters at the 2015 World Weightlifting Championship (WWC) and 2017 Pan-American Weightlifting Championship (PAWC).

2. Materials and Methods

2.1. Participants

This study was an investigation of the heaviest successful snatch attempt for all athletes who lifted in the A sessions of WWC and PAWC. The heaviest successful snatch attempt by each of the top three finishers at the time of competition in each women's and men's weight category for WWC and PAWC was also identified for separate analyses. Seven women's and 8 men's weight categories were contested at WWC, and 8 weight categories each for both women and men were contested at PAWC. Three eligible attempts each from both WWC and PAWC were not recorded due to software/hardware error, so those lifters and their results were excluded from analysis. Excluded lifters included the 3rd place finisher in the women's 75 kg category from WWC and the 3rd place finisher in the women's 48 kg category from PAWC. Seven lifters from WWC and 4 lifters from PAWC did not complete any successful attempts in the snatch and were also excluded from analysis. In total, 77 women (24.1 ± 3.1 y) and 82 men (25.0 ± 3.3 y) from WWC and 75 women (25.2 ± 5.1 y) and 85 men (23.8 ± 4.2 y) from PAWC were included in this study (Tables 1 and 2). A total of 159 lifts from WWC and 160 lifts from PAWC were analyzed.

Table 1. Athlete characteristics for 2015 World Weightlifting Championship.

	n	Age (y)	BW (kg)
Women			
48	11	23.5 ± 2.7	47.65 ± 0.23
53	12	25.3 ± 2.8	52.53 ± 0.71
58	11	25.8 ± 2.5	57.49 ± 0.51
63	12	23.0 ± 3.2	62.45 ± 0.59
69	10	24.5 ± 2.7	68.24 ± 0.81
75	10	24.3 ± 2.3	74.53 ± 0.40
+75	11	22.7 ± 4.1	114.06 ± 16.69
	77	24.1 ± 3.1	67.77 ± 21.64
Men			
56	7	24.1 ± 4.0	55.79 ± 0.13
62	11	25.4 ± 3.0	61.82 ± 0.14
69	11	24.4 ± 2.9	68.78 ± 0.15
77	11	24.7 ± 4.1	76.60 ± 0.30
85	10	24.2 ± 3.0	84.36 ± 0.38
94	10	25.0 ± 2.4	93.46 ± 0.48
105	10	24.9 ± 3.4	104.54 ± 0.45
+105	12	26.5 ± 3.7	148.74 ± 6.96
	82	25.0 ± 3.3	88.76 ± 29.02

Values are mean ± SD; BW, bodyweight.

Table 2. Athlete characteristics for 2017 Pan-American Weightlifting Championship.

	n	Age (y)	BW (kg)
Women			
48	11	24.9 ± 4.8	47.67 ± 0.31
53	9	25.5 ± 6.0	52.35 ± 1.10
58	9	27.2 ± 5.2	57.44 ± 0.68
63	9	27.1 ± 5.1	62.70 ± 0.28
69	9	21.9 ± 3.3	67.89 ± 0.96
75	10	24.8 ± 4.3	74.58 ± 0.38
90	9	25.7 ± 6.4	85.58 ± 4.22
+90	9	24.4 ± 5.2	110.01 ± 18.24
	75	25.2 ± 5.1	69.25 ± 20.07
Men			
56	10	23.2 ± 6.4	55.78 ± 0.23
62	15	22.5 ± 3.1	61.40 ± 0.60
69	14	23.8 ± 4.1	68.31 ± 0.62
77	10	21.4 ± 3.3	76.54 ± 0.56
85	8	24.3 ± 5.6	84.42 ± 0.43
94	10	25.4 ± 4.3	92.29 ± 2.62
105	9	24.7 ± 2.0	103.88 ± 1.22
+105	9	25.8 ± 3.7	135.76 ± 18.17
	85	23.8 ± 4.2	81.83 ± 24.47

Values are mean ± SD; PAWC; BW, bodyweight.

It should also be noted that the International Weightlifting Federation changed the official women's weight categories in 2016. Thus, the women's 90 kg category from PAWC was excluded from comparative analysis, and the women's +75 category from WWC and +90 category from PAWC were considered equivalent during comparative analysis.

The ethics committee of East Tennessee State University determined this study to not be human subjects research. Therefore, subjects were not required to provide informed consent for inclusion in this study.

2.2. Data Collection

Lifts were recorded at both competitions using a GoPro HERO4 Black digital video camera (San Mateo, CA, USA) in 720p resolution (1280 × 720 pixels [px]) at 240 fps. Camera setup conformed to recommendations to minimize measurement error of sagittal plane barbell kinematics [23]. The camera was arranged on a tripod 15 m and 11 m from the left edge of the competition platform at WWC and PAWC, respectively, to avoid interference with competition proceedings. The camera lens was centered and leveled 0.71 m above the competition platform surface facing in line and parallel with the platform center for both competitions.

2.3. Video Analysis

Video analysis was conducted using Kinovea software (version 0.8.27). The software's working area was set to 400% zoom to improve accuracy of digitized marker placement and distance

calibration. Kinovea's automatic tracking feature was used to determine barbell displacement, recorded in px, by placing a digitized marker on the center of the visible end of the barbell prior to lift-off. Raw displacement data was exported to a spreadsheet for subsequent analysis using a custom Labview program. Raw data was converted from px to cm using a respective scaling coefficient (WWC: $0.70740 \text{ cm}\cdot\text{px}^{-1}$; PAWC: $0.67497 \text{ cm}\cdot\text{px}^{-1}$) and smoothed using a 20-point moving average. Outcome variables were determined from the converted, smoothed data.

2.3.1. Calibration variability and inter-rater reliability

Scaling coefficients for WWC and PAWC were determined from 20 lifts from the respective competition using the following procedures. Ten lifts each for both women and men with at least one lift from each weight category were randomly selected. Kinovea's line tool was used to draw a line on the image along the vertical diameter of the largest plate (45 cm) nearest to the camera prior to lift-off. The length of the line was recorded in px and converted to a scaling factor ($\text{cm}\cdot\text{px}$) for each file. The mean scaling factor for each set of 20 lifts were also determined. Outcome variables were determined for each file using both the original scaling factor associated with each file and the respective mean scaling factor.

The mean \pm SD of calibration factors was $0.70740 \pm 0.007 \text{ cm}\cdot\text{px}^{-1}$ for WWC and $0.67497 \pm 0.008 \text{ cm}\cdot\text{px}^{-1}$ for PAWC (95% confidence intervals: 0.70440 to $0.71039 \text{ cm}\cdot\text{px}^{-1}$ and 0.67146 to $0.67847 \text{ cm}\cdot\text{px}^{-1}$, respectively). The 95% confidence intervals for % coefficient of variation for each set of original calibration scales was 0.6 to 1.3% for WWC and 0.8 to 1.6% for PAWC.

Repeated measures ANOVAs for each outcome variable revealed no statistical differences between methods ($F_{(1,19)} = 0.00040$ to 0.23 ; $p = .6$ to $> .9$ and $F_{(1,19)} = 0.0024$ to 0.14 ; $p = .7$ to $> .9$ for WWC and PAWC, respectively). 95% confidence intervals for Pearson's correlation coefficients for pairs of outcome variables determined from both methods were $r = 0.988$ to 0.999 and $r = 0.982$ to 0.999 for WWC and PAWC respectively ($p < .001$). 95% confidence limits of the standard error of measurement for all displacement variables ranged from a lower limit of -0.7 cm to an upper limit of 2.1 cm for WWC. For PAWC, 95% confidence limits of the standard error of the measurement for all displacement variables ranged from a lower limit of -0.8 cm to an upper limit of 2.5 cm . 95% confidence intervals for the standard error of measurement for peak vertical velocity ranged from -0.01 to $0.04 \text{ m}\cdot\text{s}^{-1}$ for WWC and -0.02 to $0.05 \text{ m}\cdot\text{s}^{-1}$ for PAWC. 95% confidence intervals for ICC reliability coefficients for the determination of all outcome variables between both methods were 0.977 to 0.999 and 0.969 to 0.999 for WWC and PAWC, respectively, based on a single measure, absolute agreement, 2-way random effects model ($p < .001$). Thus, the likely amount of error between the two methods was deemed to be negligible.

2.4. Outcome Variables

Barbell trajectory type was classified using definitions from Cunanan et al. [24], which are based on work by Vorobyev [5] and Hiskia [4] (Figure X). Barbell kinematic variables were modified from Stone et al. [17] and included peak vertical velocity (V_{max}), maximum barbell height (Y_{max}), height at catch (Y_{catch}), difference between Y_{max} and Y_{catch} (Y_{drop}), ratio of Y_{catch} divided by Y_{max} ($\text{Catch}_{\text{rel}}$), angle relative to vertical reference line from start position to position at X_1 (θ_1), net horizontal displacement from start position to most rearward position during first phase of displacement toward the lifter (X_1), horizontal distance from X_1 to most anterior position between X_1 and Y_{max} (X_2), horizontal distance from position at X_2 to position at catch (X_{loop}), net horizontal displacement from start position to Y_{catch} (X_{net}) (Figure X). For determination of Y_{drop} and X_{loop} , the catch was defined as the first instance after the phase of negative vertical velocity following Y_{max} that the barbell reached a vertical velocity of $0 \text{ m}\cdot\text{s}^{-1}$.




Figure X. Barbell trajectory types determined by pattern of horizontal displacement and crossing of vertical reference line. Figure redrawn and adapted from Vorobyev [5] and Hiskia [4].




Figure X. Barbell kinematic variables of displacement. Y_{max} , maximum height; Y_{catch} , height at catch; Y_{drop} , difference between Y_{max} and Y_{catch} ; θ_1 , angle relative to vertical reference line from start position to X_1 ; X_1 , net horizontal displacement from start position to most rearward position during first phase of displacement toward the lifter; X_2 , horizontal distance from X_1 to most anterior position between X_1 and Y_{max} ; X_{loop} , horizontal distance from X_2 to Y_{catch} ; X_{net} , net horizontal displacement from start position to Y_{catch} . Modified and adapted from Stone et al. [17].

2.5. Statistical Analysis

The count and relative frequency of each barbell trajectory type was categorized by competition, sex, weight category, and continent. Descriptive statistics of each kinematic variable from all A session lifts, separated by competition, sex, and weight category, were calculated.

Separate omnibus 2x3 between-subjects ANOVAs for women and men were conducted to compare the main effects of competition (WWC, PAWC) and placement (1, 2, 3) and the interaction effect of competition and placement on each of the kinematic variables from the top-three women and men finishers from both WWC and PAWC. Assumptions of normality and homoscedasticity were assessed using Shapiro-Wilk test and Levene's test, respectively. No outliers were present in the data. In cases where assumptions of normality or homoscedasticity were violated, a robust ANOVA procedure was conducted [25]. Critical alpha was set at $\alpha = .05$. Cohen's d effect size was calculated to evaluate the magnitude of all cell and marginal mean differences between groups.

3. Results

3.1. Descriptive Analysis

3.1.1. Barbell Trajectory

Relative frequencies of each trajectory type were similar between WWC and PAWC. Top-three finishers at WWC and PAWC exhibited some differences in relative frequencies of trajectory types compared to A session lifters within and between competitions (Tables X-X).

The type 3 trajectory was the most prevalent type among all A session lifters at both WWC and PAWC (53% and 59%, respectively), with heavier men's categories exhibiting a greater relative frequency than other categories. The type 3 trajectory was also the most common type among top-three finishers at both meets (43% and 49%, respectively); although, it was less prevalent among women compared to men at WWC (30% vs. 54%) and PAWC (43% vs. 54%). The type 2 trajectory was exhibited by approximately 30% of women and men at both WWC and PAWC. The type 1 trajectory accounted for 13% and 8% of lifts at WWC and PAWC, respectively. The type 4 trajectory occurred least frequently at both WWC and PAWC (6% and 3%, respectively).

A greater proportion of women's top-three finishers at both WWC and PAWC exhibited the type 2 trajectory compared to their A session counterparts (50% vs. 29% and 39% vs. 29%, respectively). The top-three finishers in the men's 105 and +105 categories at both WWC and PAWC exhibited the type 3 trajectory exclusively.

European top-three finishers from WWC most frequently exhibited the type 3 trajectory (57%), while Asian top-three finishers from WWC most frequently exhibited the type 2 trajectory (43%). There were no top-three finishers from North America, South America, or Africa at WWC. The type 4 trajectory accounted for 7% of the lifts among top-three finishers at WWC.

South American top-three finishers at PAWC most commonly exhibited the type 3 trajectory (55%) followed by the type 2 trajectory (38%). North American top-three finishers at PAWC had an equal distribution of type 2 and 3 trajectories (39% each) with the remainder being the type 1 trajectory (22%). No top-three finishers at PAWC exhibited the type 4 trajectory.

Table X. Distribution of barbell trajectory type of each athlete's heaviest successful snatch attempt in A sessions at 2015 World Weightlifting Championship.

	Type 1	Type 2	Type 3	Type 4
Women				
48	-	7 (64)	4 (36)	-
53	1 (8)	1 (8)	8 (67)	2 (17)
58	1 (9)	5 (45)	5 (45)	-
63	2 (17)	3 (25)	7 (58)	-
69	1 (10)	4 (40)	5 (50)	-
75	4 (40)	1 (10)	5 (50)	-
+75	1 (9)	1 (9)	9 (82)	-
	10 (13)	22 (29)	43 (56)	2 (3)
Men				
56	2 (29)	3 (43)	1 (14)	1 (14)
62	3 (27)	6 (55)	1 (9)	1 (9)
69	1 (9)	3 (27)	5 (45)	2 (18)
77	1 (9)	5 (45)	4 (36)	1 (9)
85	2 (20)	1 (10)	7 (70)	-
94	-	2 (20)	8 (80)	-
105	1 (10)	-	8 (80)	1 (10)
+105	-	3 (25)	8 (67)	1 (8)
	10 (12)	23 (28)	42 (51)	7 (9)
Continent				
North America	1 (17)	2 (33)	3 (50)	-
South America	3 (25)	4 (33)	4 (33)	1 (8)
Asia	6 (9)	26 (39)	31 (46)	4 (6)
Europe	9 (14)	10 (15)	43 (65)	4 (6)
Africa	1 (13)	3 (38)	4 (50)	-
Grand Total	20 (13)	45 (28)	85 (53)	9 (6)

Values are count (% relative frequency).

Table X. Distribution of barbell trajectory type of each athlete's heaviest successful snatch attempt in A sessions at 2017 Pan-American Weightlifting Championship.

	Type 1	Type 2	Type 3	Type 4
Women				
48	2 (18)	6 (55)	2 (18)	1 (9)
53	-	6 (67)	3 (33)	-
58	2 (22)	2 (22)	5 (56)	-
63	1 (11)	3 (33)	5 (56)	-
69	1 (11)	1 (11)	6 (67)	1 (11)
75	-	2 (20)	8 (80)	-
90	-	1 (11)	8 (89)	-
+90	3 (33)	1 (11)	4 (44)	1 (11)
	9 (12)	22 (29)	41 (55)	3 (4)
Men				
56	1 (10)	2 (20)	7 (70)	-
62	1 (7)	11 (73)	3 (20)	-
69	-	4 (29)	9 (64)	1 (7)
77	-	4 (40)	6 (60)	-
85	1 (13)	2 (25)	5 (63)	-
94	1 (10)	1 (10)	8 (80)	-
105	-	1 (11)	8 (89)	-
+105	-	-	8 (89)	1 (11)
	4 (5)	25 (29)	54 (64)	2 (2)
Continent				
North America	9 (11)	26 (32)	46 (56)	1 (1)
South America	4 (5)	21 (27)	49 (63)	4 (5)
Grand Total	13 (8)	47 (29)	95 (59)	5 (3)

Values are count (% relative frequency).

Table X. Distribution of barbell trajectory type of each top-three finisher's heaviest successful snatch attempt at 2015 World Weightlifting Championship.

	Type 1	Type 2	Type 3	Type 4
Women				
48	-	2	1	-
53	-	-	1	2
58	-	2	1	-
63	-	2	1	-
69	1	2	-	-
75*	1	1	-	-
+75	-	1	2	-
	2 (10)	10 (50)	6 (30)	2 (10)
Men				
56	1	1	1	-
62	2	1	-	-
69	1	-	1	1
77	-	1	2	-
85	1	1	1	-
94	-	1	2	-
105	-	-	3	-
+105	-	-	3	-
	5 (21)	5 (21)	13 (54)	1 (4)
Continent				
North America	-	-	-	-
South America	-	-	-	-
Asia	4 (13)	13 (43)	11 (37)	2 (7)
Europe	3 (21)	2 (14)	8 (57)	1 (7)
Africa	-	-	-	-
Grand Total	7 (16)	15 (34)	19 (43)	3 (7)

Values are count (% relative frequency); * one lift not recorded due to hardware/software error.

Table X. Distribution of barbell trajectory type of each top-three finisher's heaviest successful snatch attempt at 2017 Pan-American Weightlifting Championship.

	Type 1	Type 2	Type 3	Type 4
Women				
48*	1	1	-	-
53	-	2	1	-
58	-	2	1	-
63	1	1	1	-
69	1	-	2	-
75	-	2	1	-
90	-	-	3	-
+90	1	1	1	-
	4 (17)	9 (39)	10 (43)	-
Men				
56	-	2	1	-
62	-	2	1	-
69	-	2	1	-
77	-	2	1	-
85	1	-	2	-
94	1	1	1	-
105	-	-	3	-
+105	-	-	3	-
	2 (8)	9 (38)	13 (54)	-
Continent				
North America	4 (22)	7 (39)	7 (39)	-
South America	2 (7)	11 (38)	16 (55)	-
Grand Total	6 (13)	18 (38)	23 (49)	-

Values are count (% relative frequency); * one lift not recorded due to hardware/software error.

3.1.2. Kinematic Variables

Overall, the direction of the cell mean difference of most kinematic variables from WWC and PAWC was inconsistent over weight categories for both sexes (Tables X and X). Heavier lifters tended to exhibit greater V_{max} , Y_{max} , and Y_{catch} , which is partly due to heavier lifters tending to also be taller [2,4,10,12]. Despite differences in Y_{max} and Y_{catch} , Y_{drop} did not exhibit an increasing or decreasing trend over weight categories for either sex. Heavier lifters also tended to exhibit greater X_2 , which is likely partly attributable to differences in anthropometric variables among weight categories [12]. There were no discernible trends in barbell kinematics within weight categories among top-three finishers for either sex at WWC or PAWC (Tables X and X).

Table X. Load and kinematic variables for heaviest successful snatch attempt for A session lifters at 2015 World Weightlifting Championship.

	Load	V_{max}	Y_{max}	Y_{catch}	Y_{drop}	Catch_{rel}	θ₁	X₁	X₂	X_{loop}	X_{net}
	(kg)	(m·s⁻¹)	(m)	(m)	(m)	(%)	(°)	(m)	(m)	(m)	(m)
Women											
48	82 ± 4	1.82 ± 0.10	0.89 ± 0.05	0.73 ± 0.04	0.16 ± 0.04	82 ± 4	7 ± 3	0.05 ± 0.02	0.05 ± 0.02	0.09 ± 0.04	0.09 ± 0.07
53	90 ± 6	1.81 ± 0.11	0.93 ± 0.05	0.77 ± 0.06	0.16 ± 0.04	83 ± 4	5 ± 3	0.04 ± 0.02	0.05 ± 0.02	0.10 ± 0.04	0.09 ± 0.06
58	100 ± 7	1.84 ± 0.13	0.95 ± 0.04	0.79 ± 0.05	0.17 ± 0.04	83 ± 4	7 ± 4	0.06 ± 0.03	0.05 ± 0.02	0.11 ± 0.04	0.12 ± 0.09
63	104 ± 7	1.90 ± 0.11	0.97 ± 0.06	0.81 ± 0.05	0.16 ± 0.03	83 ± 3	6 ± 4	0.06 ± 0.03	0.06 ± 0.03	0.10 ± 0.06	0.11 ± 0.11
69	112 ± 6	1.78 ± 0.11	0.98 ± 0.06	0.82 ± 0.06	0.16 ± 0.04	84 ± 4	6 ± 3	0.06 ± 0.03	0.06 ± 0.02	0.11 ± 0.04	0.12 ± 0.08
75	114 ± 9	1.88 ± 0.10	1.02 ± 0.05	0.86 ± 0.06	0.16 ± 0.03	84 ± 3	5 ± 2	0.05 ± 0.02	0.08 ± 0.03	0.09 ± 0.05	0.06 ± 0.10
+75	126 ± 12	1.99 ± 0.11	1.10 ± 0.06	0.96 ± 0.06	0.15 ± 0.03	87 ± 2	7 ± 2	0.07 ± 0.02	0.07 ± 0.02	0.15 ± 0.05	0.15 ± 0.08
	104 ± 16	1.86 ± 0.13	0.98 ± 0.08	0.82 ± 0.09	0.16 ± 0.04	84 ± 4	6 ± 3	0.06 ± 0.03	0.06 ± 0.03	0.11 ± 0.05	0.10 ± 0.09
Men											
56	126 ± 9	1.79 ± 0.11	0.92 ± 0.07	0.77 ± 0.06	0.15 ± 0.03	84 ± 3	10 ± 5	0.06 ± 0.03	0.05 ± 0.02	0.08 ± 0.03	0.09 ± 0.05
62	138 ± 7	1.77 ± 0.07	0.92 ± 0.01	0.77 ± 0.04	0.15 ± 0.04	84 ± 4	7 ± 4	0.06 ± 0.04	0.05 ± 0.03	0.09 ± 0.07	0.10 ± 0.14
69	149 ± 8	1.73 ± 0.09	0.95 ± 0.04	0.79 ± 0.06	0.16 ± 0.04	83 ± 4	6 ± 2	0.05 ± 0.02	0.05 ± 0.02	0.10 ± 0.03	0.11 ± 0.06
77	162 ± 7	1.82 ± 0.09	0.99 ± 0.04	0.86 ± 0.04	0.13 ± 0.04	87 ± 4	7 ± 3	0.06 ± 0.02	0.05 ± 0.02	0.09 ± 0.04	0.10 ± 0.07
85	168 ± 7	1.83 ± 0.12	1.02 ± 0.05	0.89 ± 0.03	0.13 ± 0.04	87 ± 4	5 ± 1	0.04 ± 0.01	0.06 ± 0.02	0.09 ± 0.04	0.07 ± 0.07
94	176 ± 4	1.84 ± 0.10	1.07 ± 0.03	0.90 ± 0.06	0.17 ± 0.04	84 ± 4	5 ± 3	0.05 ± 0.04	0.07 ± 0.02	0.09 ± 0.03	0.08 ± 0.09
105	182 ± 6	1.88 ± 0.08	1.10 ± 0.03	0.93 ± 0.05	0.17 ± 0.03	84 ± 3	5 ± 2	0.05 ± 0.02	0.08 ± 0.02	0.11 ± 0.03	0.09 ± 0.05
+105	196 ± 8	1.88 ± 0.14	1.17 ± 0.08	1.02 ± 0.07	0.15 ± 0.04	87 ± 3	6 ± 3	0.07 ± 0.03	0.08 ± 0.02	0.13 ± 0.05	0.12 ± 0.09
	164 ± 22	1.82 ± 0.11	1.02 ± 0.10	0.87 ± 0.10	0.15 ± 0.04	85 ± 4	6 ± 3	0.06 ± 0.03	0.06 ± 0.03	0.10 ± 0.05	0.10 ± 0.08

Values are mean ± SD.

Table X. Load and kinematic variables for heaviest successful snatch attempt for A session lifters at 2017 Pan-American Weightlifting Championship.

	Load	V_{max}	Y_{max}	Y_{catch}	Y_{drop}	Catch_{rel}	θ₁	X₁	X₂	X_{loop}	X_{net}
	(kg)	(m·s⁻¹)	(m)	(m)	(m)	(%)	(°)	(m)	(m)	(m)	(m)
Women											
48	68 ± 10	1.78 ± 0.10	0.92 ± 0.03	0.77 ± 0.05	0.16 ± 0.03	83 ± 3	8 ± 2	0.06 ± 0.01	0.05 ± 0.02	0.09 ± 0.03	0.10 ± 0.04
53	83 ± 3	1.77 ± 0.09	0.93 ± 0.05	0.76 ± 0.06	0.17 ± 0.03	81 ± 4	8 ± 2	0.08 ± 0.02	0.04 ± 0.02	0.12 ± 0.04	0.16 ± 0.06
58	89 ± 5	1.89 ± 0.08	0.99 ± 0.02	0.81 ± 0.03	0.18 ± 0.04	82 ± 4	5 ± 2	0.05 ± 0.02	0.05 ± 0.01	0.10 ± 0.02	0.09 ± 0.04
63	89 ± 5	1.88 ± 0.08	1.00 ± 0.04	0.86 ± 0.05	0.14 ± 0.03	86 ± 3	7 ± 2	0.06 ± 0.01	0.05 ± 0.03	0.13 ± 0.06	0.14 ± 0.09
69	93 ± 9	1.94 ± 0.10	1.05 ± 0.04	0.86 ± 0.04	0.19 ± 0.04	82 ± 3	5 ± 3	0.05 ± 0.03	0.07 ± 0.03	0.10 ± 0.04	0.08 ± 0.10
75	99 ± 6	1.85 ± 0.06	1.05 ± 0.04	0.87 ± 0.07	0.17 ± 0.04	83 ± 4	0 ± 18	0.05 ± 0.03	0.07 ± 0.03	0.11 ± 0.04	0.09 ± 0.08
90	101 ± 13	1.83 ± 0.08	1.04 ± 0.03	0.89 ± 0.04	0.15 ± 0.04	86 ± 4	6 ± 5	0.06 ± 0.05	0.07 ± 0.02	0.12 ± 0.04	0.11 ± 0.10
+90	99 ± 22	1.99 ± 0.12	1.12 ± 0.07	0.98 ± 0.09	0.14 ± 0.04	87 ± 4	8 ± 4	0.08 ± 0.04	0.08 ± 0.03	0.13 ± 0.05	0.13 ± 0.10
	90 ± 15	1.86 ± 0.11	1.01 ± 0.07	0.85 ± 0.09	0.16 ± 0.04	84 ± 4	6 ± 7	0.06 ± 0.03	0.06 ± 0.03	0.11 ± 0.04	0.11 ± 0.08
Men											
56	102 ± 10	1.77 ± 0.11	0.95 ± 0.07	0.81 ± 0.05	0.14 ± 0.04	85 ± 3	6 ± 3	0.05 ± 0.03	0.04 ± 0.03	0.11 ± 0.04	0.11 ± 0.09
62	113 ± 11	1.84 ± 0.16	0.97 ± 0.06	0.82 ± 0.07	0.14 ± 0.04	85 ± 4	9 ± 3	0.08 ± 0.03	0.03 ± 0.02	0.12 ± 0.06	0.17 ± 0.10
69	120 ± 16	1.79 ± 0.10	0.99 ± 0.05	0.85 ± 0.05	0.14 ± 0.04	86 ± 4	7 ± 3	0.07 ± 0.04	0.06 ± 0.04	0.10 ± 0.05	0.11 ± 0.12
77	145 ± 8	1.81 ± 0.10	1.02 ± 0.04	0.86 ± 0.03	0.16 ± 0.03	84 ± 3	6 ± 3	0.06 ± 0.04	0.03 ± 0.02	0.13 ± 0.04	0.16 ± 0.09
85	155 ± 5	1.78 ± 0.12	1.03 ± 0.04	0.89 ± 0.06	0.14 ± 0.04	86 ± 4	5 ± 2	0.05 ± 0.02	0.04 ± 0.02	0.13 ± 0.04	0.15 ± 0.06
94	153 ± 12	1.85 ± 0.09	1.08 ± 0.06	0.91 ± 0.05	0.17 ± 0.05	84 ± 4	5 ± 3	0.05 ± 0.03	0.05 ± 0.03	0.11 ± 0.05	0.11 ± 0.10
105	163 ± 11	1.84 ± 0.15	1.12 ± 0.06	0.97 ± 0.08	0.15 ± 0.03	86 ± 3	4 ± 4	0.04 ± 0.04	0.07 ± 0.02	0.09 ± 0.05	0.06 ± 0.10
+105	165 ± 11	1.96 ± 0.12	1.20 ± 0.07	1.05 ± 0.06	0.15 ± 0.05	87 ± 4	5 ± 3	0.06 ± 0.03	0.08 ± 0.02	0.11 ± 0.04	0.09 ± 0.05
	136 ± 25	1.83 ± 0.13	1.03 ± 0.09	0.88 ± 0.09	0.15 ± 0.04	85 ± 4	6 ± 3	0.06 ± 0.03	0.05 ± 0.03	0.11 ± 0.05	0.12 ± 0.10

Values are mean ± SD.

Table X. Load and kinematic variables for heaviest successful snatch attempt for top-three women in each weight category at 2015 World Weightlifting Championship.

	Load	V_{max}	Y_{max}	Y_{catch}	Y_{drop}	Catch_{rel}	θ₁	X₁	X₂	X_{loop}	X_{net}
	(kg)	(m·s⁻¹)	(m)	(m)	(m)	(%)	(°)	(m)	(m)	(m)	(m)
Women											
48											
1	88	1.80	0.86	0.67	0.19	78	9	0.07	0.03	0.13	0.18
2	85	1.93	0.94	0.74	0.21	78	0	0.00	0.06	0.06	0.00
3	85	1.73	0.81	0.70	0.12	86	5	0.04	0.03	0.12	0.12
53											
1	101	1.61	0.82	0.66	0.16	80	2	0.01	0.04	0.18	0.15
2	96	1.78	0.88	0.74	0.15	83	2	0.02	0.07	0.07	0.02
3	96	1.68	0.89	0.70	0.20	78	5	0.04	0.04	0.08	0.08
58											
1	112	1.94	0.94	0.78	0.16	83	8	0.07	0.04	0.09	0.12
2	108	1.71	0.88	0.71	0.17	80	11	0.11	0.00	0.18	0.29
3	106	2.10	1.02	0.88	0.15	86	7	0.06	0.06	0.13	0.13
63											
1	113	1.79	0.92	0.75	0.17	82	7	0.07	0.03	0.15	0.18
2	112	1.74	0.99	0.82	0.17	83	7	0.07	0.05	0.11	0.12
3	112	1.86	0.94	0.81	0.14	86	9	0.08	0.06	0.11	0.12
69											
1	120	1.79	0.95	0.76	0.19	80	6	0.06	0.03	0.20	0.22
2	116	1.69	0.97	0.77	0.20	80	8	0.08	0.04	0.09	0.14
3	116	1.94	1.11	0.89	0.22	80	5	0.05	0.07	0.07	0.05
75											
1	127	1.94	0.99	0.86	0.13	87	5	0.05	0.08	0.09	0.07
2	125	1.88	0.98	0.82	0.16	84	7	0.07	0.02	0.13	0.18
3*	-	-	-	-	-	-	-	-	-	-	-
+75											
1	148	1.95	1.07	0.96	0.10	90	7	0.08	0.04	0.11	0.15
2	145	1.92	1.06	0.94	0.12	88	5	0.06	0.04	0.13	0.15
3	136	2.18	1.14	1.02	0.12	90	8	0.08	0.05	0.14	0.17

* Lift not recorded due to hardware/software error.

Table X. Load and kinematic variables for heaviest successful snatch attempt for top-three men in each weight category at 2015 World Weightlifting Championship.

	Load (kg)	V_{max} (m·s⁻¹)	Y_{max} (m)	Y_{catch} (m)	Y_{drop} (m)	Catch_{rel} (%)	θ₁ (°)	X₁ (m)	X₂ (m)	X_{loop} (m)	X_{net} (m)
Men											
56											
1	139	1.67	0.93	0.73	0.20	78	11	0.07	0.07	0.06	0.06
2	132	1.87	0.85	0.76	0.09	89	2	0.01	0.03	0.05	0.03
3	131	1.62	0.82	0.67	0.15	82	8	0.05	0.05	0.07	0.06
62											
1	151	1.68	0.91	0.72	0.19	79	5	0.04	0.07	0.01	-0.02
2	150	1.87	0.93	0.76	0.17	82	12	0.11	0.01	0.22	0.32
3	141	1.72	0.92	0.81	0.11	88	6	0.05	0.07	0.05	0.03
69											
1	160	1.80	0.98	0.84	0.14	86	5	0.04	0.06	0.08	0.07
2	160	1.78	0.91	0.81	0.10	89	6	0.05	0.02	0.11	0.13
3	158	1.84	0.99	0.82	0.17	83	1	0.01	0.06	0.07	0.02
77											
1	175	1.87	1.03	0.89	0.15	86	8	0.07	0.05	0.09	0.11
2	171	1.85	1.00	0.86	0.14	86	6	0.05	0.05	0.08	0.09
3	167	1.69	0.94	0.83	0.11	88	3	0.04	0.03	0.12	0.13
85											
1	178	1.94	1.05	0.92	0.13	87	7	0.07	0.04	0.17	0.20
2	176	1.70	0.95	0.88	0.07	93	6	0.05	0.06	0.07	0.06
3	173	1.79	0.96	0.85	0.11	88	6	0.02	0.08	0.08	0.03
94											
1	181	2.02	1.05	0.96	0.09	91	4	0.04	0.05	0.08	0.07
2	180	1.70	1.01	0.80	0.21	79	5	0.05	0.05	0.09	0.08
3	178	1.95	1.09	0.89	0.20	82	11	0.12	0.05	0.12	0.19
105											
1	192	1.87	1.15	0.92	0.23	80	4	0.04	0.08	0.15	0.11
2	191	1.79	1.07	0.93	0.14	87	8	0.07	0.10	0.08	0.05
3	186	1.85	1.07	0.92	0.16	85	7	0.08	0.07	0.06	0.07
+105											
1	211	1.79	1.14	0.99	0.15	87	8	0.10	0.08	0.12	0.14
2	207	1.95	1.30	1.17	0.13	90	4	0.06	0.09	0.09	0.05
3	203	1.89	1.09	1.01	0.08	93	1	0.01	0.10	0.03	-0.05

Table X. Load and kinematic variables for heaviest successful snatch attempt for top-three women in each weight category at 2017 Pan-American Weightlifting Championship.

	Load (kg)	V_{max} (m·s⁻¹)	Y_{max} (m)	Y_{catch} (m)	Y_{drop} (m)	Catch_{rel} (%)	θ₁ (°)	X₁ (m)	X₂ (m)	X_{loop} (m)	X_{net} (m)
Women											
48											
1	77	1.69	0.92	0.77	0.15	84	6	0.08	0.00	0.10	0.17
2	77	1.63	0.94	0.79	0.15	84	5	0.05	0.06	0.04	0.02
3*	-	-	-	-	-	-	-	-	-	-	-
53											
1	88	1.85	1.03	0.84	0.19	82	9	0.09	0.04	0.08	0.13
2	86	1.70	0.88	0.71	0.18	80	8	0.06	0.04	0.11	0.13
3	86	1.78	0.95	0.76	0.18	81	9	0.09	0.04	0.12	0.16
58											
1	94	1.84	0.97	0.80	0.17	83	9	0.07	0.06	0.09	0.10
2	94	1.85	1.00	0.82	0.18	82	2	0.02	0.05	0.12	0.09
3	93	1.85	0.98	0.82	0.15	85	7	0.06	0.05	0.10	0.11
63											
1	97	2.01	1.03	0.90	0.14	87	7	0.06	0.10	0.05	0.02
2	93	1.86	0.98	0.81	0.17	82	6	0.06	0.03	0.19	0.22
3	93	2.01	1.02	0.93	0.09	91	7	0.06	0.07	0.09	0.08
69											
1	107	1.85	0.99	0.78	0.20	80	8	0.08	0.02	0.18	0.24
2	100	1.94	1.04	0.87	0.17	84	3	0.03	0.08	0.08	0.03
3	99	2.00	1.05	0.88	0.17	84	4	0.03	0.07	0.08	0.04
75											
1	110	1.85	1.00	0.84	0.16	84	-39	0.00	0.07	0.08	0.01
2	103	1.78	1.07	0.88	0.19	82	7	0.08	0.04	0.16	0.20
3	101	1.78	1.01	0.82	0.19	81	7	0.07	0.03	0.13	0.17
+90											
1	120	2.16	1.24	1.14	0.10	92	7	0.07	0.10	0.11	0.09
2	119	1.86	1.17	1.02	0.15	87	11	0.15	0.05	0.24	0.34
3	118	1.92	1.11	0.99	0.11	90	6	0.07	0.08	0.12	0.11

* Lift not recorded due to hardware/software error.

Table X. Load and kinematic variables for heaviest successful snatch attempt for top-three men in each weight category at 2017 Pan-American Weightlifting Championship.

	Load	V_{max}	Y_{max}	Y_{catch}	Y_{drop}	Catch_{rel}	θ₁	X₁	X₂	X_{loop}	X_{net}
	(kg)	(m·s⁻¹)	(m)	(m)	(m)	(%)	(°)	(m)	(m)	(m)	(m)
Men											
56											
1	119	1.61	0.86	0.75	0.11	87	6	0.05	0.03	0.12	0.14
2	118	1.70	0.87	0.80	0.07	92	6	0.05	0.00	0.14	0.19
3	108	1.85	0.95	0.79	0.16	83	6	0.06	0.01	0.17	0.22
62											
1	125	1.71	0.92	0.81	0.11	88	8	0.07	0.02	0.13	0.18
2	124	1.71	0.91	0.75	0.15	83	11	0.10	0.03	0.06	0.13
3	123	1.85	0.95	0.78	0.18	82	4	0.03	0.05	0.04	0.03
69											
1	145	1.93	0.99	0.90	0.09	91	10	0.11	0.01	0.16	0.25
2	143	1.65	0.96	0.79	0.17	82	5	0.05	0.04	0.13	0.14
3	137	1.71	0.98	0.82	0.16	84	5	0.06	0.03	0.12	0.15
77											
1	162	2.00	1.02	0.81	0.20	80	8	0.08	0.01	0.15	0.22
2	152	1.69	0.99	0.84	0.15	85	10	0.11	0.02	0.14	0.23
3	151	1.77	0.95	0.82	0.13	86	8	0.09	0.01	0.23	0.31
85											
1	160	1.78	1.05	0.88	0.17	84	7	0.08	0.01	0.19	0.25
2	159	1.77	1.07	0.90	0.17	84	3	0.03	0.03	0.17	0.17
3	158	2.00	1.09	0.99	0.10	91	4	0.04	0.06	0.08	0.06
94											
1	169	1.93	1.05	0.89	0.16	85	8	0.08	0.03	0.11	0.16
2	166	1.86	1.07	0.87	0.20	82	4	0.04	0.02	0.18	0.20
3	166	2.01	1.24	0.99	0.25	79	6	0.06	0.07	0.05	0.03
105											
1	182	1.78	1.13	0.97	0.16	86	2	0.02	0.05	0.10	0.07
2	175	1.93	1.12	0.98	0.13	88	4	0.05	0.09	0.06	0.02
3	167	1.63	1.15	0.94	0.21	82	2	0.02	0.04	0.08	0.07
+105											
1	178	2.08	1.21	1.15	0.06	95	2	0.02	0.07	0.14	0.09
2	176	1.78	1.16	0.99	0.16	86	8	0.09	0.06	0.07	0.10
3	175	1.86	1.13	0.98	0.15	87	5	0.05	0.10	0.09	0.04

3.2. Comparative Analysis

3.2.1. Women

Y_{catch} , θ_1 , and X_1 from PAC did not meet the assumption of normality ($p = .01$, $< .001$, and $.005$, respectively; all other $p = .1$ to $> .9$). θ_1 also did not satisfy the assumption of homoscedasticity ($p = .002$; all other $p = .1$ to $> .9$). A statistical main effect of competition was found only for Y_{max} ($\eta^2 = 0.106$; $F_{(1,34)} = 4.2$; $p = .049$). Y_{max} mean \pm SD was 0.96 ± 0.09 m for WWC and 1.02 ± 0.08 m for PAWC (Cohen's d 95% confidence interval = -1.3 to -0.02). No statistical main or interaction effects were found for any other variable ($p = .2$ to $> .9$) (Table X). Magnitude and directionality of effect sizes varied considerably among variables (Cohen's $d = -0.87$ to 0.77) (Figure X).

Table X. Results of omnibus 2x3 between-subjects ANOVAs for women.

	Competition			Placement			Competition by placement		
	η^2	F	p	η^2	F	p	η^2	F	p
V_{max}	0.0017	0.06	.8	0.092	1.8	.2	0.020	0.38	.7
Y_{max}	0.11	4.2	.049*	0.011	0.21	.8	0.019	0.37	.7
$Y_{\text{catch}}^\ddagger$	-	-	.3	-	-	.4	-	-	$> .9$
Y_{drop}	0.0011	0.041	.8	0.050	0.90	.4	0.0036	0.066	$> .9$
$\text{Catch}_{\text{rel}}$	0.024	0.89	.4	0.048	0.88	.4	0.0010	0.018	$> .9$
θ_1^\ddagger	-	-	$> .9$	-	-	.8	-	-	.9
X_1^\ddagger	-	-	$> .9$	-	-	.8	-	-	.9
X_2	0.043	1.6	.2	0.026	0.48	$> .9$	0.006	0.11	.9
X_{loop}	0.0059	0.23	.6	0.025	0.48	.7	0.090	1.7	.2
X_{net}	0.0045	0.16	.7	0.018	0.33	.8	0.032	0.58	.6

*, statistical effect at $p < .05$; \ddagger , η^2 and F statistic were not included in robust ANOVA procedure output.




Figure X. Cohen's d effect size with 95% confidence interval for main effects of competition (a-c) and placement (d-i) among top-three women: (a) 1st place at WWC vs. 1st place at PAWC; (b) 2nd place at WWC vs. 2nd place at PAWC; (c) 3rd place at WWC vs. 3rd place at PAWC; (d) 1st place at WWC vs. 2nd place at WWC; (e) 1st place at WWC vs. 3rd place at WWC; (f) 2nd place at WWC vs. 3rd place at WWC; (g) 1st place at PAWC vs. 2nd place at PAWC; (h) 1st place at PAWC vs. 3rd place at PAWC; (i) 2nd place at PAWC vs. 3rd place at PAWC; WWC, 2015 World Weightlifting Championship; PAWC, 2017 Pan-American Weightlifting Championship.

3.2.2. Men

No variable violated assumptions of normality or homoscedasticity ($p = .1$ to $> .9$). A statistical main effect of competition was found for X_2 ($\eta^2 = 0.17$; $F_{(1,42)} = 8.8$; $p = .005$), X_{loop} ($\eta^2 = 0.10$; $F_{(1,42)} = 5.6$; $p = .02$), and X_{net} ($\eta^2 = 0.13$; $F_{(1,42)} = 6.5$; $p = .01$). Mean \pm SD for X_2 was 0.06 ± 0.02 m for WWC and 0.04 ± 0.03 m for PAWC (Cohen's d 95% confidence interval = 0.24 to 1.48). Mean \pm SD for X_{loop} was 0.09 ± 0.04 m for WWC and 0.12 ± 0.05 m for PAWC (Cohen's d 95% confidence interval = -1.3 to -0.06). Mean \pm SD for X_{net} was 0.08 ± 0.08 for WWC and 0.14 ± 0.08 m for PAWC (Cohen's d 95% confidence interval = -1.3 to -0.13). No statistical main or interaction effects were present for any other variable ($p = .1$ to $> .9$) (Table X). Effect sizes showed considerable variation with no clear pattern within or across factor levels (Cohen's $d = -1.06$ to 0.68) (Figure X).

Table X. Results of omnibus 2x3 between-subjects ANOVAs for men.

	Competition			Placement			Competition by placement		
	η^2	F	p	η^2	F	p	η^2	F	p
V_{max}	0.00026	0.01	> .9	0.036	0.81	.5	0.031	0.69	.5
Y_{max}	0.020	0.87	.4	0.0059	0.13	.9	0.021	0.47	.6
Y_{catch}	0.0097	0.45	.5	0.010	0.24	.8	0.086	2.0	.1
Y_{drop}	0.010	0.45	.5	0.0046	0.10	> .9	0.0079	0.17	.8
$Catch_{rel}$	0.0017	0.08	.8	0.0087	0.20	.8	0.068	1.5	.2
θ_1	0.00029	0.01	> .9	0.027	0.59	.6	0.0049	0.11	> .9
X_1	0.015	0.66	.4	0.042	0.94	.4	0.0020	0.04	> .9
X_2	0.17	8.8	.005*	0.024	0.62	.5	0.016	0.41	.7
X_{loop}	0.10	5.1	.03*	0.045	1.1	.3	0.0090	0.22	.8
X_{net}	0.13	6.6	.01*	0.057	1.5	.2	0.0060	0.16	.9

*, statistical effect at $p < .05$.




Figure X. Cohen's d effect size with 95% confidence interval for main effects of competition (a-c) and placement (d-i) among top-three men: (a) 1st place at WWC vs. 1st place at PAWC; (b) 2nd place at WWC vs. 2nd place at PAWC; (c) 3rd place at WWC vs. 3rd place at PAWC; (d) 1st place at WWC vs. 2nd place at WWC; (e) 1st place at WWC vs. 3rd place at WWC; (f) 2nd place at WWC vs. 3rd place at WWC; (g) 1st place at PAWC vs. 2nd place at PAWC; (h) 1st place at PAWC vs. 3rd place at PAWC; (i) 2nd place at PAWC vs. 3rd place at PAWC; WWC, 2015 World Weightlifting Championship; PAWC, 2017 Pan-American Weightlifting Championship.

4. Discussion

The aim of this study was to identify technical and biomechanical factors that might characterize successful performances of the snatch lift for elite weightlifters. This study involved 1) descriptive analysis of barbell trajectory and kinematics of the snatch lift for A session lifters at two major international competitions and 2) comparative analysis of barbell kinematics between top-three performers in the snatch lift from each meet.

The pooled relative frequencies of type 2 and 3 trajectories for women at WWC and PAWC were similar to those observed for women at the 1993 and 1994 World and European Weightlifting Championships [4]. Pooled data show that women at WWC and PAWC exhibited the type 1 trajectory less frequently (~10%) than the women included in the report by Hiskia [4]. Antoniuk et al. observed from 137 women competing at several World and European Weightlifting Championships that the type 2 trajectory was most common [26]. It is unclear in what years 2 out of the 4 competitions included in that study took place. A subsequent report by this group presented data from a total of 304 attempts by 140 women presumably from the same competitions reported in their first study that the type 1 trajectory was most common with the remainder sharing an equal distribution of type 2 and 3 trajectories [15]. Both studies reported a greater prevalence of the type 3 trajectory among the +75 category [15,26]. Pooled data for men at WWC and PAWC show similar relative frequency of the type 1 trajectory, lower relative frequency for the type 2 trajectory (~14%), and higher relative frequency for the 3 trajectory (~9%) compared to men at the 1993 and 1994 World and European Championships [4]. To the authors' knowledge, the present study is the first to quantify the prevalence of the type 4 trajectory at any competitive level; it is thus unknown how reclassification of data from prior studies to include the type 4 trajectory would affect comparison to the results of the present study.

The present results corroborate the findings of Akkuş [9] who observed that female world champions at the 2010 World Weightlifting Championship exhibited a variety of trajectory types. Musser et al. [12] found that no women snatch medalists at the 2009 Pan-American Weightlifting Championship exhibited the type 1 trajectory. However, the results of the present study demonstrate that top-three women at the Pan-American championship level can also likely exhibit a variety of trajectory types including the type 1 trajectory. The differences in relative frequencies of trajectory types between this study and previous reports [4,12,15,26] are likely due to different athlete pools, and it is unclear whether the success of certain trajectory types are exclusive to or more likely to occur in particular weight categories. However, it is likely from these differences that the relative frequencies of trajectory types observed among women at different major international competitions do vary and that high placement is not exclusive to any existing trajectory type. Although, some trajectory types may potentially be more common among top performers in some weight categories partly due to anthropometry [10,12].

Although Baumann et al. [6] did not report counts or relative frequencies of barbell trajectory type, they did indicate that 'nearly all' men in the A sessions at the 1985 World Weightlifting Championship exhibited the type 2 trajectory, with the remainder presumably exhibiting the type 1 trajectory. Baumann et al. [6] did not report any occurrences of the type 3 trajectory. Limited data from Garhammer do indicate the presence of the type 3 trajectory for the snatch and the clean among women and men world and Olympic champions during the 1980's [27,28]. However, the type 3 trajectory does not appear to have been common among top international lifters during that period [6,27]. In fact, Garhammer suggested that technique assessed by barbell trajectory had not changed from the mid-1970's to the late-1980's [27]. Subsequently, Hiskia [4] found the type 3 trajectory to be the most common among both women and men in the A sessions of the 1993 and 1994 World and European Weightlifting Championships. Data from Akkuş [9] of all women snatch gold medalists and Harbili [11] of women and men in the A session of their respective 69 kg categories at the 2010 World Weightlifting Championship indicate increased prevalence of the type 3 trajectory among top international weightlifters at that competition. Collectively, these data seem to indicate that the relative frequencies of barbell trajectory types likely vary between

and within international competitions. Furthermore, the available evidence indicates a likely shift in technique among international weightlifters based on barbell trajectory.

The observed differences in the prevalence of trajectory types among top international weightlifters may reflect changes in coaching philosophy, teaching methods, and/or training methods to accommodate what have previously been considered 'suboptimal' technique (i.e. type 2 and 3 trajectories) [5,9,27]. Furthermore, the differences observed between women in this study and women in previous investigations may partially be due to the selective pressure of elite competition. Given the longer history of men's weightlifting, such selective pressure is less likely to be a factor for men currently. Kinematic differences due to body size and anthropometry [4,10,12] may also partly explain observed differences for both sexes due to periodic changes in weight categories. Serial investigations can help to delineate any apparent trends across extant cross-sectional analyses and future competitions. Furthermore, continued study of the relative frequency of trajectory types over a range of competitive levels and subgroups among them can help to identify which trajectory type(s), if any, is most characteristic of a given group.

The observations in this and recent studies [9,11] of high placing individuals at the world championship level who exhibit the type 3 trajectory is noteworthy. The type 1 trajectory was initially considered to be most favorable [5,27]. Some authors have more recently suggested the type 2 trajectory to be advantageous [17]. However, the type 3 trajectory is generally regarded to be non-beneficial and potentially detrimental [5,17] based on several biomechanical and theoretical bases. Garhammer and Taylor [29] found that anterior barbell displacement at lift-off, such as occurs with the type 3 trajectory, results in a forward shift of the lifter's center of pressure, or balance point. Such barbell displacement also increases moment arm length between the barbell center of mass and joint centers, thereby increasing joint moments and muscular forces required to lift the barbell [6]. Furthermore, anterior displacement during the first pull can increase mechanical work and decrease lift efficiency [30,31]. Thus, the prevalence of the type 3 trajectory among top-three finishers—especially among the men's 105 and +105 categories, who lifted the heaviest loads—at both WWC and PAWC emphasize the importance of high levels of absolute strength, possibly to overcome apparent technical deficiencies [17]. Greater lower limb length may also partly explain the increased prevalence of the type 3 trajectory among heavier categories [12], and anthropometric variables may partly explain differences in the prevalence of trajectory types more generally [12,32]. Nonetheless, numerous observations of the gamut of trajectory types among A session and top-three international weightlifters somewhat challenges the notion that barbell trajectory type is a useful criterion of effective weightlifting technique at this level.

Few statistical effects were observed in this study. The observed main effect of competition for Y_{\max} among top-three women in this study could be due to differences in stature [2,6,10,33], skill [7,8], or load lifted [33]. However, the lack of accompanying statistical or clear effect size differences for Y_{drop} , which has also been suggested to depend on skill [7,8], and observations of weightlifters who lifted heavier loads to greater absolute and relative vertical displacements compared to lower caliber athletes in the same weight category [8] suggest stature to be a more likely explanation of Y_{\max} differences observed in this study. Top-three men at WWC, who lifted the heaviest loads of any group in this study, exhibited greater X_2 and less X_{net} and X_{loop} than top-three men at PAWC. It is unlikely that greater X_2 itself is beneficial for performance, as greater X_2 would increase the overall work and energy required to complete the lift [6,34-36] and may increase instability during the catch [6]. These effects could be compounded by a potential subsequent increase of X_{loop} , which is likely to be less during successful attempts [17]. Greater X_2 among individuals of greater weightlifting ability may be consequent to the larger forces and accelerations associated with lifting heavier loads [3,6,8,14,37]. As such, it is not recommended that individuals attempt to deliberately increase X_2 such as by 'hipping' or swinging the bar away during the second pull [17,38]. The reduced X_{loop} and X_{net} among top-three men at WWC likely indicate that they jumped backward less than top-three men at PAWC. These results support the findings of Stone et al. that, while net rearward displacement is generally not detrimental or disadvantageous,

smaller relative X_{loop} and X_{net} are likely associated with greater weightlifting success and ability [17]. Several authors have suggested that the direction of force application is important especially during the second pull [17,38,39]. The observations of reduced X_{loop} and X_{net} among stronger weightlifters possibly reflect these individuals' ability to produce greater vertical force and acceleration [3,6,8,36,40]. Stronger individuals are also more likely to produce faster rates of force development [41], which may improve their ability to counteract greater X_2 through a more effective reversal of anterior horizontal barbell acceleration during the second pull and turnover phase thereby reducing X_{loop} and X_{net} [38,39]. Altogether, these results suggest that greater strength may improve energy flow, force application, and vertical acceleration to favorably influence horizontal barbell kinematics that affect weightlifting performance and ability.

Studies that have identified kinematic and kinetic differences based on skill or sex have consistently identified factors that partly depend on strength [8,9,11,37]. For example, greater strength improves the ability to perform stretch-shortening cycle tasks [42-44], such as occurs during the transition phase of the weightlifting pull. Thus, strength likely mediates the rate and magnitude of knee flexion during the transition phase [9,11,13]. Several studies have also made direct comparisons between different groups of weightlifters (e.g. women vs. men [45], adolescent vs. adult [37], district vs. national/international [14]) and found overall similarity in kinematic and kinetic structure. However, when considering differences in weightlifting ability, there are notable differences in maximum ground reaction force [3,14,46], rate of force development [14,47], and absolute and relative joint and whole body power [31,37,48-50], which are all dependent on maximum strength [41]. The lifters in this study generally lifted heavier loads compared to lifters in the same or similar weight categories in previous studies [6,9,11,12,36] while exhibiting no clear differences in technique, lending further credence to the notion that strength is the primary determinant of weightlifting ability.

Indeed, there exists strong relationships between measures of maximum strength and weightlifting ability among weightlifters of a variety of competitive level. For example, Stone et al. reported Pearson's $r = 0.79$ to 0.95 for the relationships between back squat one-repetition maximum (1RM) and snatch and clean 1RMs and Pearson's $r = 0.83$ to 0.84 for the relationships between isometric peak force assessed using the isometric mid-thigh pull and snatch and clean 1RMs in national and international level junior and senior weightlifters from the United States [51]. Lucero et al. reported Pearson's $r = 0.91$ to 0.94 for the relationships between self-reported back and front squat 1RMs and snatch and clean 1RMs among male competitive weightlifters in the United States [52]. Furthermore, greater strength levels may also be beneficial for the expression of other desirable physical characteristics, such as rate of force development, impulse, and power [41].

The observed variety of barbell trajectory types and overall lack of pattern among individual or clusters of kinematic variables in this study suggest no standard 'technique profile' is requisite for high achievement in weightlifting. In fact, several investigations have had limited success differentiating kinematic profiles of successful versus unsuccessful weightlifting attempts [38,39]. However, such findings should not be interpreted to suggest that technique is not an important determinant of weightlifting success at any level. Rather these findings more likely reflect the fact that the weightlifting movements are influenced by multiple degrees of freedom [17,53-55], and the varied results of this study may reflect individualized solutions to the degrees of freedom problem in weightlifting. The results of this study thus suggest the possibility of a variety of effective individualized technique profiles for weightlifting performance and ability.

It is purported that individual weightlifting technique stabilizes after only a few months of training [1]. However, the amount of intra-individual variation of lifter or barbell kinematics and kinetics during maximal attempts under competition conditions is yet to be elucidated. The results presented in two studies by Antoniuk et al., which analyzed different sets of snatch attempts from the same pool of athletes and competitions, substantiates that an individual can exhibit different barbell trajectories across attempts [15,26]. Nonetheless, the reliability of individual technique profiles is unclear. These

uncertainties notwithstanding, individual technique profiles consisting of any or some technical or biomechanical parameters may potentially be useful for evaluating, monitoring, or predicting weightlifting performance and ability.

There are a variety of laboratory and field-based technologies available to conduct weightlifting technique analysis [24], and both researchers and coaches should explore the development of individual technique profiles. The methods of this study provide support for the implementation of video analysis for determining barbell kinematics and developing technique profiles. The total time to analyze a single video file including calibration during this study was less than two minutes. Use of the Kinovea's built-in graphics and analysis functions could reduce this time to 15 to 30 seconds. Additionally, Carson et al. reported a case study that demonstrate another such approach that involved long-term monitoring of lifter kinematics during an intervention intended to change weightlifting technique [56]. This case study and the anecdotal experiences of athletes and coaches highlight the complexity and difficulty of instilling changes to technique that are robust enough to persist during attempts at high relative intensities [56]. Thus, the relatively short latency of weightlifting technique stabilization and the associated challenges of technique correction underscore the importance of establishing sound technique during the earliest stages of a weightlifter's career.

Instructional and coaching methods should generally be guided by principles and tenets from the fields of biomechanics, motor learning, and physiology. While the available evidence has not identified universal optimal technique, there are general guidelines for basic weightlifting technique apparent from the extensive body of scientific literature [2-6,24]. However, coaches should consider and make appropriate accommodations for individual differences that may manifest nuances or peculiarities in technique.

5. Conclusions

The methods used in this study demonstrate an inexpensive, time-efficient, and user-friendly method to conduct technique analysis. The error and reliability of the methods herein can be improved by optimizing camera setup and specifications [23,35,57], controlling conditions during data collection (e.g. barbell placement [58] and lighting [59]), and selecting appropriate data smoothing or filtering techniques [59-61]. Video analysis may aid coaches in determining barbell kinematics and developing technique profiles. Such technique profiles may be useful in the evaluation, monitoring, or prediction of weightlifting performance and ability.

A variety of barbell trajectories were observed with differences based on competition, sex, category, and ranking. There was no discernible pattern of statistical or effect size differences for most of the kinematic variables observed in this study. Incidentally, the variability of individual weightlifting technique at any level requires further determination. Thus, while practically relevant, barbell trajectory or any of the examined kinematic variables alone is unlikely to reliably indicate weightlifting ability at this level. Therefore, coaches may consider evaluating weightlifting technique within a more general framework.

It seems apparent from this study's results and the available evidence that strength may be the most important determinant of weightlifting performance and ability [2,11,14,17,37,47,51,52]. It is possible that greater strength may also aid weightlifting performance by compensating for apparent technical deficiencies [17]. Thus, once an individual's technique is established, weightlifting training should primarily emphasize the development of strength and other related physical characteristics using a periodized approach [62,63] while incorporating complementary methods to refine technique [56].

Author Contributions: Conceptualization, A.J.C. and M.H.S.; methodology, A.J.C., S.M., K.S., and M.H.S.; software, S.M.; investigation, A.J.C., W.G.H., M.A.S., and K.P.U.; data curation: A.J.C and K.P.U.; validation, A.J.C.; formal analysis, A.J.C. and S.M.; visualization, A.J.C.; writing—original draft preparation, A.J.C. and M.H.S.; writing—review and editing, A.J.C., W.G.H., M.A.S., S.M., K.C.P., K.S., and M.H.S.; project administration, A.J.C. and M.H.S.; supervision, S.M., K.C.P., K.S., M.H.S.

Funding: This research received no external funding.

Acknowledgments: The authors would like to thank the International Weightlifting Federation, the Pan-American Weightlifting Federation, and USA Weightlifting for their cooperation and support to conduct this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Aján, T.; Baroga, L. *Weightlifting: fitness for all sports*; International Weightlifting Federation: Budapest, Hungary, 1988.
2. Bartonietz, K.E. Biomechanics of the snatch: Toward a higher training efficiency. *Strength and Conditioning Journal* **1996**, *18*, 24-31.
3. Enoka, R.M. The pull in Olympic weightlifting. *Medicine and Science in Sports* **1979**, *11*, 131-137.
4. Hiskia, G. Biomechanical analysis of performance of World and Olympic champions. In Proceedings of Proceedings of Weightlifting Symposium, Ancient Olympia, Greece; pp. 137-163.
5. Vorobyev, A.N. *A Textbook on Weightlifting*; International Weightlifting Federation: Budapest, Hungary, 1978.
6. Baumann, W.; Gross, V.; Quade, K.; Galbierz, P.; Schwirtz, A. The snatch technique of world class weightlifters at the 1985 world championships. *International Journal of Sport Biomechanics* **1988**, *4*, 68-89.
7. Burdett, R.G. Biomechanics of the snatch technique of highly skilled and skilled weightlifters. *Research Quarterly for Exercise and Sport* **1982**, *53*, 193-197.
8. Liu, G.; Fekete, G.; Yang, H.; Ma, J.; Sun, D.; Mei, Q.; Gu, Y. Comparative 3-dimensional kinematic analysis of snatch technique between top-elite and sub-elite male weightlifters in 69-kg category. *Heliyon* **2018**, *4*.
9. Akkuş, H. Kinematic analysis of the snatch lift with elite female weightlifters during the 2010 World Weightlifting Championship. *Journal of Strength and Conditioning Research* **2012**, *26*, 897-905.
10. Ford, L.E.; Dettlerline, A.J.; Ho, K.K.; Cao, W. Gender-and height-related limits of muscle strength in world weightlifting champions. *Journal of Applied Physiology* **2000**, *89*, 1061-1064.
11. Harbili, E. A gender-based kinematic and kinetic analysis of the snatch lift in elite weightlifters in 69-kg category. *Journal of Sports Science & Medicine* **2012**, *11*, 162.
12. Musser, L.J.; Garhammer, J.; Rozenek, R.; Crusemeyer, J.A.; Vargas, E.M. Anthropometry and barbell trajectory in the snatch lift for elite women weightlifters. *Journal of Strength and Conditioning Research* **2014**, *28*, 1636-1648.
13. Gourgoulis, V.; Aggeloussis, N.; Antoniou, P.; Christoforidis, C.; Mavromatis, G.; Garas, A. Comparative 3-dimensional kinematic analysis of the snatch technique in elite male and female Greek weightlifters. *Journal of Strength and Conditioning Research* **2002**, *16*, 359-366.
14. Kauhanen, H.; Häkkinen, K.; Komi, P.V. A biomechanical analysis of the snatch and clean & jerk techniques of Finish elite and district level weightlifters. *Scandinavian Journal of Sports Sciences* **1984**, *6*, 47-56.
15. Antoniuk, O.; Vynogradskyi, B.; Pavlyuk, O.; Chopyk, T.; Pavlyuk, Y.; Soltyk, O. Peculiarities of barbell motion trajectory during snatch lift of elite female weightlifters. *Journal of Physical Education and Sport* **2017**, *17*, 402-406.
16. Garhammer, J. Biomechanical characteristics of the 1978 world weightlifting champions. *International Series on Biomechanics* **1981**, *3*, 300-304.
17. Stone, M.H.; O'Bryant, H.S.; Williams, F.E.; Johnson, R.L.; Pierce, K.C. Analysis of bar paths during the snatch in elite male weightlifters. *Strength and Conditioning Journal* **1998**, *20*, 30-38.
18. Bosch, F. *Strength Training and Coordination: an Integrative Approach*; 2010Publishers: Rotterdam, Netherlands, 2015.
19. Stone, M.H.; Stone, M.; Sands, W.A. *Principles and Practice of Resistance Training*; Human Kinetics: Champaign, IL, 2007.
20. Verkhoshansky, Y.V.; Siff, M.C. *Supertraining*, 6th ed.; Supertraining Institute: Denver, CO, 2009.
21. Issurin, V. Training transfer: scientific background and insights for practical application. *Sports Medicine* **2013**, *43*, 675-694.
22. Suarez, D.G.; Wagle, J.P.; Cunanan, A.J.; Sausaman, R.W.; Stone, M.H. Dynamic correspondence of resistance training to sport: a brief review. *Strength and Conditioning Journal* **2018**, DOI: 10.1519/SSC.0000000000000458, doi:DOI: 10.1519/SSC.0000000000000458.
23. Garhammer, J.; Newton, H. Applied video analysis for coaches: Weightlifting examples. *International Journal of Sports Science & Coaching* **2013**, *8*, 581-594.
24. Cunanan, A.J.; Mizuguchi, S.; Sato, K.; Pierce, K.C.; Stone, M.H. Methods to evaluate the weightlifting pull: a practical review. *Professional Strength and Conditioning* **In review**.

25. Mair, P.; Wilcox, R. WRS2: Wilcox robust estimation and testing. **2018**.
26. Antoniuk, O.V.; Pavlyuk, O.S.; Chopyk, T.V.; Pavlyuk, E.A. Characteristics of barbell trajectory in snatch, fulfilled by elite female weight-lifters. *Pedagogics, Psychology, Medical-biological Problems of Physical Training and Sports* **2016**, *20*, 4-8, doi:10.15561/18189172.2016.0601.
27. Garhammer, J. Bar trajectories of world champion male and female weightlifters: coaching applications. Part 1. *International Olympic Lifter* **1989**, *10*, 7-8.
28. Garhammer, J. Bar trajectories of world champion male and female weightlifters: coaching applications. Part 2. *International Olympic Lifter* **1990**, *10*, 12-13.
29. Garhammer, J.; Taylor, L. Center of pressure movements during weightlifting. In Proceedings of 2nd International Symposium on Biomechanics in Sports, Colorado Springs, USA; pp. 279-291.
30. Garhammer, J. Performance evaluation of Olympic weightlifters. *Medicine and Science in Sports* **1979**, *11*, 284-287.
31. Garhammer, J. Biomechanical profiles of Olympic weightlifters. *International Journal of Sport Biomechanics* **1985**, *1*, 122-130.
32. Schilling, B.K.; Stone, M.H.; O'bryant, H.S.; Fry, A.C.; Coglianesi, R.H.; Pierce, K.C. Snatch technique of collegiate national level weightlifters. *Journal of Strength and Conditioning Research* **2002**, *16*, 551-555.
33. Hoover, D.L.; Carlson, K.M.; Christensen, B.K.; Zebas, C.J. Biomechanical analysis of women weightlifters during the snatch. *Journal of Strength and Conditioning Research* **2006**, *20*, 627.
34. Garhammer, J. Energy flow during Olympic weight lifting. *Medicine and Science in Sports and Exercise* **1982**, *14*, 353-360.
35. Garhammer, J. A Review of Power Output Studies of Olympic and Powerlifting: Methodology, Performance. *Journal of Strength and Conditioning Research* **1993**, *7*, 76-89.
36. Isaka, T.; Okada, J.; Funato, K. Kinematic analysis of the barbell during the snatch movement of elite Asian weight lifters. *Journal of Applied Biomechanics* **1996**, *12*, 508-516.
37. Gourgoulis, V.; Aggeloussis, N.; Kalivas, V.; Antoniou, P.; Mavromatis, G. Snatch lift kinematics and bar energetics in male adolescent and adult weightlifters. *Journal of Sports Medicine and Physical Fitness* **2004**, *44*, 126.
38. Kipp, K.; Meinerz, C. A biomechanical comparison of successful and unsuccessful power clean attempts. *Sports Biomechanics* **2017**, *16*, 272-282.
39. Gourgoulis, V.; Aggeloussis, N.; Garas, A.; Mavromatis, G. Unsuccessful vs. successful performance in snatch lifts: a kinematic approach. *Journal of Strength and Conditioning Research* **2009**, *23*, 486-494.
40. Kipp, K.; Harris, C. Patterns of barbell acceleration during the snatch in weightlifting competition. *Journal of Sports Sciences* **2015**, *33*, 1467-1471.
41. Suchomel, T.J.; Nimphius, S.; Stone, M.H. The importance of muscular strength in athletic performance. *Sports Medicine* **2016**, *46*, 1419-1449, doi:10.1007/s40279-016-0486-0.
42. Cormie, P.; McBride, J.M.; McCaulley, G.O. Power-time, force-time, and velocity-time curve analysis of the countermovement jump: impact of training. *Journal of Strength & Conditioning Research* **2009**, *23*, 177-186.
43. Cormie, P.; McCaulley, G.O.; McBride, J.M. Power versus strength-power jump squat training: influence on the load-power relationship. *Medicine and Science in Sports and Exercise* **2007**, *39*, 996-1003.
44. Turner, A.N.; Jeffreys, I. The stretch-shortening cycle: proposed mechanisms and methods for enhancement. *Strength & Conditioning Journal* **2010**, *32*, 87-99.
45. Gourgoulis, V.; Aggeloussis, N.; Mavromatis, G.; Garas, A. Three-dimensional kinematic analysis of the snatch of elite Greek weightlifters. *Journal of Sports Sciences* **2000**, *18*, 643-652.
46. Häkkinen, K.; Kauhanen, H.; Komi, P. Biomechanical changes in the Olympic weightlifting technique of the snatch and clean and jerk from submaximal to maximal loads. *Scandinavian Journal of Sports Sciences* **1984**, *6*, 57-66.
47. Häkkinen, K.; Kauhanen, H.; Komi, P.V. Merkmale neuromuskulärer Leistungskapazität bei Gewichthebern nationalen und regionalen Niveaus. *Leistungssport* **1985**, *15*, 35-41.
48. Flores, F.J.; Sedano, S.; Redondo, J.C. Optimal load and power spectrum during snatch and clean: differences between international and national weightlifters. *International Journal of Performance Analysis in Sport* **2017**, *17*, 521-533.
49. Garhammer, J. Power production by Olympic weightlifters. *Medicine and Science in Sports and Exercise* **1980**, *12*, 54-60.
50. Garhammer, J. A comparison of maximal power outputs between elite male and female weightlifters in competition. *International Journal of Sport Biomechanics* **1991**, *7*, 3-11.

51. Stone, M.H.; Sands, W.A.; Pierce, K.C.; Carlock, J.; Cardinale, M.; Newton, R.U. Relationship of maximum strength to weightlifting performance. *Medicine and Science in Sports and Exercise* **2005**, *37*, 1037-1043.
52. Lucero, R.A.J.; Fry, A.C.; LeRoux, C.D.; Hermes, M.J. Relationships between barbell squat strength and weightlifting performance. *International Journal of Sports Science and Coaching* **2019**, *0*, 1-7, doi:10.1177/1747954119837688.
53. Bernstein, N. *The coordination and regulation of movements*; Pergamon Press: New York, NY, 1967.
54. Enoka, R.M. Muscular control of a learned movement: The speed control system hypothesis. *Experimental Brain Research* **1983**, *51*, 135-145.
55. Kipp, K.; Redden, J.; Sabick, M.B.; Harris, C. Weightlifting performance is related to kinematic and kinetic patterns of the hip and knee joints. *Journal of Strength and Conditioning Research* **2012**, *26*, 1838-1844.
56. Carson, H.J.; Collins, D.; Jones, B. A case study of technical change and rehabilitation: intervention design and interdisciplinary team interaction. *International Journal of Sport Psychology* **2014**, *45*, 57-78.
57. Payton, C.J. Motion Analysis Using Video. In *Biomechanical Evaluation of Movement in Sport and Exercise*, Payton, C.j.B., R M, Ed. Routledge: New York, NY, 2008.
58. Chiu, H.-T.; Wang, C.-H.; Cheng, K.B. The three-dimensional kinematics of a barbell during the snatch of Taiwanese weightlifters. *Journal of Strength and Conditioning Research* **2010**, *24*, 1520-1526.
59. Dæhlin, T.E.; Krosshaug, T.; Chiu, L.Z. Enhancing digital video analysis of bar kinematics in weightlifting: A case study. *Journal of Strength and Conditioning Research* **2017**, *31*, 1592-1600.
60. Garhammer, J.; Whiting, W.C. Comparison of three data smoothing techniques for determining weightlifting kinematics. *Scandinavian Journal of Sports Sciences* **1989**, *11*, 97-104.
61. Wood, G.A. Data smoothing and differentiation procedures in biomechanics. *Exercise and Sport Sciences Reviews* **1982**, *10*, 308-362.
62. Cunanan, A.J.; DeWeese, B.H.; Wagle, J.P.; Carroll, K.M.; Sausaman, R.; Hornsby III, W.G.; Haff, G.G.; Triplett, N.T.; Pierce, K.C.; Stone, M.H. The general adaptation syndrome: a foundation for the concept of periodization. *Sports Medicine* **2018**, *48*, 787-797, doi:10.1007/s40279-017-0855-3.
63. Suchomel, T.J.; Comfort, P.; Lake, J.P. Enhancing the force-velocity profile of athletes using weightlifting derivatives. *Strength and Conditioning Journal* **2017**, *39*, 10-20.



© 2019 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

CHAPTER 4

SUMMARY AND FUTURE DIRECTIONS

This dissertation was designed to 1) update to the scientific knowledge of weightlifting technique and performance, 2) improve coaches' ability to conduct and interpret technique analysis, and 3) enhance transferability of weightlifting in training to improve sport performance. The methods used in this dissertation involved video recording of lifts performed during the 2015 World Weightlifting Championship (WWC) and 2017 Pan-American Weightlifting Championship (PAWC), with subsequent analysis using the free, open-source software program Kinovea (version 0.8.27) and a custom Labview program. These methods were chosen because they offer a valid, reliable, and inexpensive way to collect and analyze a large volume of data. Secondarily, these methods demonstrate a practicable method for coaches to incorporate technique analysis into their daily practice. The total time to analyze a single video file was less than 2 minutes. This process can be reduced to 15 to 30 seconds by using Kinovea's built-in graphics and analysis functions.

The heaviest successful snatch attempts by 319 weightlifters competing in the A sessions at WWC and PAWC were analyzed during this project, which is the largest sample of athletes of this caliber included in a study since 1997 (Hiskia, 1997). This dissertation builds upon the findings of previous research to conclude that weightlifting technique assessed by barbell trajectory has changed since the 1980's. Most notably, the type 3 trajectory was most commonly observed, even among top-three finishers, in the present study. These changes may reflect changes in coaching philosophy, teaching methods, and/or training methods. The results of this dissertation also suggest that the barbell kinematic variables analyzed cannot be used to reliably differentiate international weightlifters by competition or ranking. These results further suggest

that optimal technique is likely individualized. Therefore, coaches may be interested in developing individual technique profiles to evaluate, monitor, or predict weightlifting performance and ability. The results of this dissertation also suggest that strength may be the primary determinant of weightlifting ability. Thus, once individual weightlifting technique is established, coaches should seek to develop maximum strength and other related characteristics using a periodized approach while incorporating complementary methods to refine technique.

It is apparent that individuals exhibit technique differences between attempts; however, the degree of intra-individual variation in weightlifting technique is unclear. Therefore, an important objective for future research is to establish the reliability of various indices of weightlifting technique for diverse samples to aid coaches in developing technique profiles. While top international weightlifters represent the most highly skilled performers of the weightlifting movements, it is also necessary to characterize weightlifting technique across a broad range of levels to better understand potential differences between levels. Thus, more research dedicated to examining weightlifting technique among different levels is needed. Although there is a high degree of similarity between the snatch and the clean, there is a lack of studies examining technical and biomechanical parameters of the clean. Direct comparison between the snatch and the clean, especially intra-individual comparison, also merits investigation. Additionally, future research can help to elucidate technical and biomechanical differences between successful and unsuccessful attempts.

REFERENCES

- Adesida, Y., Papi, E., & McGregor, A. H. (2019). Exploring the role of wearable technology in sport kinematics and kinetics: a systematic review. *Sensors, 19*(7), 1597. DOI: 10.3390/s19071597
- Aján, T., & Baroga, L. (1988). Weightlifting: fitness for all sports. Budapest, Hungary: International Weightlifting Federation.
- Akkuş, H. (2012). Kinematic analysis of the snatch lift with elite female weightlifters during the 2010 World Weightlifting Championship. *Journal of Strength and Conditioning Research, 26*(4), 897-905.
- Antoniuk, O. V., Pavlyuk, O. S., Chopyk, T. V., & Pavlyuk, E. A. (2016). Characteristics of barbell trajectory in snatch, fulfilled by elite female weight-lifters. *Pedagogics, Psychology, Medical-Biological Problems of Physical Training and Sports, 20*(6), 4-8. DOI:10.15561/18189172.2016.0601
- Antoniuk, O., Vynogradskyi, B., Pavlyuk, O., Chopyk, T., Pavlyuk, Y., & Soltyk, O. (2017). Peculiarities of barbell motion trajectory during snatch lift of elite female weightlifters. *Journal of Physical Education and Sport, 17*(1), 402-406.
- Arabatzi, F., & Kellis, E. (2012). Olympic Weightlifting Training Causes Different Knee Muscle–Coactivation Adaptations Compared with Traditional Weight Training. *Journal of Strength and Conditioning Research, 26*(8), 2192-2201.
- Arend, S., & Higgins, J. R. (1976). A strategy for the classification, subjective analysis, and observation of human movement. *Journal of Human Movement Studies, 2*, 36-52.

- Bartonietz, K. E. (1996). Biomechanics of the snatch: Toward a higher training efficiency. *Strength and Conditioning Journal*, 18(3), 24-31.
- Baumann, W., Gross, V., Quade, K., Galbierz, P., & Schwirtz, A. (1988). The snatch technique of world class weightlifters at the 1985 world championships. *International Journal of Sport Biomechanics*, 4(1), 68-89.
- Bazyler, C. D., Mizuguchi, S., Zourdos, M. C., Sato, K., Kavanaugh, A. A., DeWeese, B. H., . . . Stone, M. H. (2017). Characteristics of a National Level Female Weightlifter Peaking for Competition: A Case Study. *Journal of Strength and Conditioning Research*, 32(1), 3029-3038.
- Bernstein, N. (1967). The coordination and regulation of movements. New York, NY: Pergamon Press.
- Bober, T. (1979). Biomechanical aspects of sports techniques. Paper presented at the Seventh International Congress of Biomechanics, Warsaw, Poland.
- Bosch, F. (2015). Strength Training and Coordination: an Integrative Approach. Rotterdam, Netherlands: 2010Publishers.
- Bruenger, A. J., Smith, S. L., Sands, W. A., & Leigh, M. R. (2007). Validation of instrumentation to monitor dynamic performance of Olympic weightlifters. *Journal of Strength and Conditioning Research*, 21(2), 492-499.
- Burdett, R. G. (1982). Biomechanics of the snatch technique of highly skilled and skilled weightlifters. *Research Quarterly for Exercise and Sport*, 53(3), 193-197.

- Carson, H. J., Collins, D., & Jones, B. (2014). A case study of technical change and rehabilitation: intervention design and interdisciplinary team interaction. *International Journal of Sport Psychology, 45*(1), 57-78.
- Chaouachi, A., Hammami, R., Kaabi, S., Chamari, K., Drinkwater, E. J., & Behm, D. G. (2014). Olympic weightlifting and plyometric training with children provides similar or greater performance improvements than traditional resistance training. *Journal of Strength and Conditioning Research, 28*(6), 1483-1496.
- Chen, S. K., Wu, M. T., Huang, C. H., Wu, J. H., Guo, L. Y., & Wu, W. L. (2013). The analysis of upper limb movement and EMG activation during the snatch under various loading conditions. *Journal of Mechanics in Medicine and Biology, 13*(1), 1-13. DOI: 10.1142/S0219519413500103.
- Chiu, H.-T., Wang, C.-H., & Cheng, K. B. (2010). The three-dimensional kinematics of a barbell during the snatch of Taiwanese weightlifters. *Journal of Strength and Conditioning Research, 24*(6), 1520-1526.
- Chiu, L. Z. F., Schilling, B. K., Fry, A. C., & Salem, G. J. (2008). The influence of deformation on barbell mechanics during the clean pull. *Sports Biomechanics, 7*(2), 260-273.
- Cormie, P., McBride, J. M., & McCaulley, G. O. (2009). Power-time, force-time, and velocity-time curve analysis of the countermovement jump: impact of training. *Journal of Strength and Conditioning Research, 23*(1), 177-186.
- Cormie, P., McCaulley, G. O., & McBride, J. M. (2007). Power versus strength-power jump squat training: influence on the load-power relationship. *Medicine and Science in Sports and Exercise, 39*(6), 996-1003.

- Courel-Ibáñez, J., Martínez-Cava, A., Morán-Navarro, R., Escribano-Peñas, P., Chavarren-Cabrero, J., González-Badillo, J. J., & Pallarés, J. G. (2019). Reproducibility and repeatability of five different technologies for bar velocity measurement in resistance training. *Annals of Biomedical Engineering*, 1-19. DOI: 10.1007/s10439-019-02265-6.
- Cunanan, A. J., DeWeese, B. H., Wagle, J. P., Carroll, K. M., Sausaman, R., Hornsby III, W. G., . . . Stone, M. H. (2018). The general adaptation syndrome: a foundation for the concept of periodization. *Sports Medicine*, 48(4), 787-797. DOI:10.1007/s40279-017-0855-3
- Cunanan, A. J., Mizuguchi, S., Sato, K., Pierce, K. C., & Stone, M. H. (In review). Methods to evaluate the weightlifting pull: A practical review. *Professional Strength and Conditioning*.
- Czaplicki, A., Dziewiecki, K., Mazur, Z., & Blajer, W. (2019). Assessment of internal loads in the joints of the lower extremities during the snatch in young weightlifters. *International Journal of Structural Stability and Dynamics*, 19(5), 1-19. DOI: 10.1142/S0219455419410116.
- Dæhlin, T. E., Krosshaug, T., & Chiu, L. Z. (2017). Enhancing digital video analysis of bar kinematics in weightlifting: A case study. *Journal of Strength and Conditioning Research*, 31(6), 1592-1600.
- Enoka, R. M. (1979). The pull in Olympic weightlifting. *Medicine and Science in Sports*, 11(2), 131-137.
- Enoka, R. M. (1983). Muscular control of a learned movement: The speed control system hypothesis. *Experimental Brain Research*, 51(1), 135-145.

- Enoka, R. M. (1988). Load-and skill-related changes in segmental contributions to a weightlifting movement. *Medicine and Science in Sports and Exercise*, 20(2), 178-187.
- Enoka, R. M. (2015). *Neuromechanics of Human Movement* (5th ed.). Champaign, IL: Human Kinetics.
- Felici, F., Rosponi, A., Sbriccoli, P., Filligoi, G. C., Fattorini, L., & Marchetti, M. (2001). Linear and non-linear analysis of surface electromyograms in weightlifters. *European Journal of Applied Physiology*, 84, 337-342.
- Flores, F. J., Sedano, S., & Redondo, J. C. (2017). Optimal load and power spectrum during snatch and clean: differences between international and national weightlifters. *International Journal of Performance Analysis in Sport*, 17(4), 521-533.
- Flores, F. J., Sedano, S., de Benito, A. M., & Redondo, J. C. (2016). Validity and reliability of a 3-axis accelerometer for measuring weightlifting movements. *International Journal of Sports Science & Coaching*, 11(6), 872-879.
- Ford, L. E., Detterline, A. J., Ho, K. K., & Cao, W. (2000). Gender-and height-related limits of muscle strength in world weightlifting champions. *Journal of Applied Physiology*, 89(3), 1061-1064.
- Garhammer, I. (1993). A Review of Power Output Studies of Olympic and Powerlifting: Methodology, Performance. *Journal of Strength and Conditioning Research*, 7(2), 76-89.
- Garhammer, J. (1979). Performance evaluation of Olympic weightlifters. *Medicine and Science in Sports*, 11(3), 284-287.

- Garhammer, J. (1980). Power production by Olympic weightlifters. *Medicine and Science in Sports and Exercise*, 12(1), 54-60.
- Garhammer, J. (1981). Biomechanical characteristics of the 1978 world weightlifting champions. *International Series on Biomechanics*, 3, 300-304.
- Garhammer, J. (1982). Energy flow during Olympic weight lifting. *Medicine and Science in Sports and Exercise*, 14(5), 353-360.
- Garhammer, J. (1985). Biomechanical profiles of Olympic weightlifters. *International Journal of Sport Biomechanics*, 1(2), 122-130.
- Garhammer, J. (1989). Bar trajectories of world champion male and female weightlifters: coaching applications. Part 1. *International Olympic Lifter*, 10(5), 7-8.
- Garhammer, J. (1990). Bar trajectories of world champion male and female weightlifters: coaching applications. Part 2. *International Olympic Lifter*, 10(6), 12-13.
- Garhammer, J. (1991). A comparison of maximal power outputs between elite male and female weightlifters in competition. *International Journal of Sport Biomechanics*, 7(1), 3-11.
- Garhammer, J., & Gregor, R. (1992). Propulsion forces as a function of intensity for weightlifting and vertical jumping. *Journal of Applied Sport Science Research*, 6(3), 129-134.
- Garhammer, J., & Newton, H. (2013). Applied video analysis for coaches: Weightlifting examples. *International Journal of Sports Science & Coaching*, 8(3), 581-594.

- Garhammer, J., & Taylor, L. (1984). Center of pressure movements during weightlifting. Paper presented at the 2nd International Symposium on Biomechanics in Sports, Colorado Springs, USA.
- Garhammer, J., & Whiting, W. C. (1989). Comparison of three data smoothing techniques for determining weightlifting kinematics. *Scandinavian Journal of Sports Sciences*, *11*(3), 97-104.
- Gorton III, G. E., Herbert, D. A., & Gannotti, M. (2009). Assessment of the kinematic variability among 12 motion analysis laboratories. *Gait & Posture*, *29*, 398-402.
- Gourgoulis, V., Aggelousis, N., Mavromatis, G., & Garas, A. (2000). Three-dimensional kinematic analysis of the snatch of elite Greek weightlifters. *Journal of Sports Sciences*, *18*(8), 643-652.
- Gourgoulis, V., Aggeloussis, N., Antoniou, P., Christoforidis, C., Mavromatis, G., & Garas, A. (2002). Comparative 3-dimensional kinematic analysis of the snatch technique in elite male and female Greek weightlifters. *Journal of Strength and Conditioning Research*, *16*(3), 359-366.
- Gourgoulis, V., Aggeloussis, N., Garas, A., & Mavromatis, G. (2009). Unsuccessful vs. successful performance in snatch lifts: a kinematic approach. *Journal of Strength and Conditioning Research*, *23*(2), 486-494.
- Gourgoulis, V., Aggeloussis, N., Kalivas, V., Antoniou, P., & Mavromatis, G. (2004). Snatch lift kinematics and bar energetics in male adolescent and adult weightlifters. *Journal of Sports Medicine and Physical Fitness*, *44*(2), 126.

- Grieve, D. W. (1970). The defeat of gravity in weightlifting. *British Journal of Sports Medicine*, 5, 37-41.
- Hackett, D., Davies, T., Soomro, N., & Halaki, M. (2016). Olympic weightlifting training improves vertical jump height in sportspeople. *British Journal of Sports Medicine*, 50(865-872). DOI:10.1136/bjsports-2015-094951
- Haff, G. G., Carlock, J. M., Hartman, M. J., & Kilgore, J. L. (2005). Force-time curve characteristics of dynamic and isometric muscle actions of elite women olympic weightlifters. *Journal of Strength and Conditioning Research*, 19(4), 741.
- Haff, G. G., Jackson, J. R., Kawamori, N., Carlock, J. M., Hartman, M. J., Kilgore, J. L., . . . Stone, M. H. (2008). Force-time curve characteristics and hormonal alterations during an eleven-week training period in elite women weightlifters. *Journal of Strength and Conditioning Research*, 22(2), 433-446.
- Hakkinen, K., & Kauhanen, H. (1986). A biomechanical analysis of selected assistant exercises of weightlifting. *Journal of Human Movement Studies*, 12(6), 271-288.
- Häkkinen, K., Kauhanen, H., & Komi, P. (1984). Biomechanical changes in the Olympic weightlifting technique of the snatch and clean and jerk from submaximal to maximal loads. *Scandinavian Journal of Sports Sciences*, 6(2), 57-66.
- Hakkinen, K., Kauhanen, H., & Komi, P. V. (1985). Merkmale neuromuskulärer Leistungskapazität bei Gewichthebern nationalen und regionalen Niveaus. *Leistungssport*, 15(5), 35-41.

- Häkkinen, K., Komi, P. V., Alén, M., & Kauhanen, H. (1987). EMG, muscle fibre and force production characteristics during a 1 year training period in elite weight-lifters. *European Journal of Applied Physiology*, *56*, 419-427.
- Harbili, E. (2012). A gender-based kinematic and kinetic analysis of the snatch lift in elite weightlifters in 69-kg category. *Journal of Sports Science & Medicine*, *11*(1), 162.
- Hiskia, G. (1997). Biomechanical analysis of performance of World and Olympic champions. Paper presented at the Proceedings of Weightlifting Symposium, Ancient Olimpia, Greece.
- Ho, L. K., Lorenzen, C., Wilson, C. J., Saunders, J. E., & Williams, M. D. (2014). Reviewing current knowledge in snatch performance and technique: the need for future directions in applied research. *Journal of Strength and Conditioning Research*, *28*(2), 574-586.
- Ibarra, F. J. R., Encinas, M. T. S., Gómez, C. E. R., & Garcia, O. A. P. (2014). Design of a biomechanical model and a set of neural networks for monitoring weightlifting. *Research in Computing Science*, *80*, 31-42.
- Isaka, T., Okada, J., & Funato, K. (1996). Kinematic analysis of the barbell during the snatch movement of elite Asian weight lifters. *Journal of Applied Biomechanics*, *12*(4), 508-516.
- Issurin, V. (2013). Training transfer: scientific background and insights for practical application. *Sports Medicine*, *43*(8), 675-694.
- Jiang, H., Sun, D., Jampani, V., Yang, M. H., Learned-Miller, E., & Kautz, J. (2018). Super slomo: high quality estimation of multiple intermediate frames for video interpolation.

Paper presented at the IEEE Conference on Computer Vision and Pattern Recognition,
Salt Lake City, UT.

Kauhanen, H., Häkkinen, K., & Komi, P. V. (1984). A biomechanical analysis of the snatch and clean & jerk techniques of Finish elite and district level weightlifters. *Scandinavian Journal of Sports Sciences*, 6, 47-56.

Kipp, K., & Harris, C. (2017). Muscle-specific effective mechanical advantage and joint impulse in weightlifting. *Journal of Strength and Conditioning Research*, 31(7), 1905-1910.

Kipp, K., & Meinerz, C. (2017). A biomechanical comparison of successful and unsuccessful power clean attempts. *Sports Biomechanics*, 16(2), 272-282.

Kipp, K., Giordanelli, M., & Geiser, C. (2018). Predicting net joint moments during a weightlifting exercise with a neural network model. *Journal of Biomechanics*, 74, 225-229.

Kipp, K., Harris, C., & Sabick, M. B. (2011). Lower extremity biomechanics during weightlifting exercise vary across joint and load. *Journal of Strength and Conditioning Research*, 25(5), 1229-1234.

Kipp, K., Redden, J., Sabick, M. B., & Harris, C. (2012). Weightlifting performance is related to kinematic and kinetic patterns of the hip and knee joints. *Journal of Strength and Conditioning Research*, 26(7), 1838-1844.

Kipp, K., Redden, J., Sabick, M. B., & Harris, C. (2012b). Weightlifting performance is related to kinematic and kinetic patterns of the hip and knee joints. *Journal of Strength and Conditioning Research*, 26(7), 1838-1844.

- Kipp, K., Redden, J., Sabick, M., & Harris, C. (2012a). Kinematic and kinetic synergies of the lower extremities during the pull in Olympic weightlifting. *Journal of Applied Biomechanics*, 28(3), 271-278.
- Lee, S., DeRosia, K. D., & Lamie, L. M. (2017). Evaluating the contribution of lower extremity kinetics to whole body power output during the power snatch. *Sports Biomechanics*, 17(4), 554-556.
- Lee, S., DeRosia, K., Lamie, L., & Levine, N. (2017). Correlation profiles between lower extremity net joint torques and whole body power during the power clean. *International Journal of Sports Science*, 7(2), 66-72.
- Lees, A. (2002). Technique analysis in sports: a critical review. *Journal of Sports Sciences*, 20(10), 813-828.
- Liu, G., Fekete, G., Yang, H., Ma, J., Sun, D., Mei, Q., & Gu, Y. (2018). Comparative 3-dimensional kinematic analysis of snatch technique between top-elite and sub-elite male weightlifters in 69-kg category. *Heliyon*, 4(7).
- Liu, Y., & Chen, W. (2001). Foot pressure study during pulling phase of snatch lifting. Paper presented at the 19th International Symposium on Biomechanics in Sports, San Francisco, USA.
- Lucero, R. A. J., Fry, A. C., LeRoux, C. D., & Hermes, M. J. (2019). Relationships between barbell squat strength and weightlifting performance. *International Journal of Sports Science and Coaching*, 0(0), 1-7. DOI:10.1177/1747954119837688.
- Mair, P., & Wilcox, R. (2018). WRS2: Wilcox robust estimation and testing.

- Merriaux, P., Dupuis, Y., Boutteau, R., Vasseur, P., & Savatier, X. (2017). A study of Vicon system positioning performance. *Sensors, 17*, 1-18. DOI: 10.3390/s17071591
- Musser, L. J., Garhammer, J., Rozenek, R., Crussemeyer, J. A., & Vargas, E. M. (2014). Anthropometry and barbell trajectory in the snatch lift for elite women weightlifters. *Journal of Strength and Conditioning Research, 28*(6), 1636-1648.
- Payton, C. J. (2008). Motion Analysis Using Video. In C. j. B. Payton, R M (Ed.), *Biomechanical Evaluation of Movement in Sport and Exercise*. New York, NY: Routledge.
- Peake, J. M., Kerr, G., & Sullivan, J. P. (2018). A critical review of consumer wearables, mobile applications, and equipment for providing biofeedback, monitoring stress, and sleep in physically active populations. *Frontiers in Physiology, 9*, 1-19. DOI: 10.3389/fphys.2018.00743
- Robertson, D. G. E., Caldwell, G. E., Hamill, J., Kamen, G., & Whittlesey, S. N. (2004). *Research Methods in Biomechanics*. Champaign, IL: Human Kinetics.
- Rossi, S. J., Buford, T. W., Smith, D. B., Kennel, R., Haff, E. E., & Haff, G. G. (2007). Bilateral comparison of barbell kinetics and kinematics during a weightlifting competition. *International Journal of Sports Physiology and Performance, 2*(2), 150-158.
- Salaami, F., Jamshidi, N., Rostami, M., & Najarian, S. (2008). Power enhancement of weightlifters during snatch through reducing torque on joints by particle swarm optimization. *American Journal of Applied Sciences, 5*(12), 1670-1675.

- Sato, K., Smith, S. L., & Sands, W. A. (2009). Validation of an accelerometer for measuring sport performance. *Journal of Strength and Conditioning Research*, 23(1), 341-347.
- Schilling, B. K., Stone, M. H., O'bryant, H. S., Fry, A. C., Coglianese, R. H., & Pierce, K. C. (2002). Snatch technique of collegiate national level weightlifters. *Journal of Strength and Conditioning Research*, 16(4), 551-555.
- Schöllhorn, W. I. (2004). Applications of artificial neural nets in clinical biomechanics. *Clinical Biomechanics*, 19, 876-898.
- Seitz, L. B., Trajano, G. S., & Haff, G. G. (2014). The back squat and the power clean: elicitation of different degrees of potentiation. *International Journal of Sports Physiology and Performance*, 9, 643-649.
- Smidt, G. L. (1973). Biomechanical analysis of knee flexion and extension. *Journal of Biomechanics*, 6, 79-92.
- Srisen, P., Auephanwiriyakul, S., Theera-Umpon, N., & Chamnongkich, S. (2015). Kinect joints correction using optical flow for weightlifting videos. Paper presented at the 7th International Conference on Computational Intelligence, Modelling and Simulation, Kuantan, Malaysia.
- Stone, M. H. (2013). Overtraining in strength power athletes. Paper presented at the ACSM Annual Meeting, Indianapolis, IN.
- Stone, M. H., O'Bryant, H. S., Williams, F. E., Johnson, R. L., & Pierce, K. C. (1998). Analysis of bar paths during the snatch in elite male weightlifters. *Strength and Conditioning Journal*, 20(4), 30-38.

- Stone, M. H., Sands, W. A., Pierce, K. C., Carlock, J., Cardinale, M., & Newton, R. U. (2005). Relationship of maximum strength to weightlifting performance. *Medicine and Science in Sports and Exercise*, 37(6), 1037-1043.
- Stone, M. H., Stone, M., & Sands, W. A. (2007). Principles and Practice of Resistance Training. Champaign, IL: Human Kinetics.
- Storey, A., & Smith, H. K. (2012). Unique aspects of competitive weightlifting. *Sports Medicine*, 42(9), 769-790.
- Suarez, D. G., Wagle, J. P., Cunanan, A. J., Sausaman, R. W., & Stone, M. H. (2018). Dynamic correspondence of resistance training to sport: a brief review. *Strength and Conditioning Journal*. DOI: 10.1519/SSC.0000000000000458.
- Suchomel, T. J., Comfort, P., & Lake, J. P. (2017). Enhancing the force-velocity profile of athletes using weightlifting derivatives. *Strength and Conditioning Journal*, 39(1), 10-20.
- Suchomel, T. J., Comfort, P., & Lake, J. P. (2017). Enhancing the force-velocity profile of athletes using weightlifting derivatives. *Strength and Conditioning Journal*, 39(1), 10-20.
- Suchomel, T. J., Nimphius, S., & Stone, M. H. (2016). The importance of muscular strength in athletic performance. *Sports Medicine*, 46(10), 1419-1449. DOI:10.1007/s40279-016-0486-0.
- Tricoli, V., Lamas, L., Carnevale, R., & Ugrinowitsch, C. (2005). Short-term effects on lower-body functional power development: weightlifting vs. vertical jump training programs. *Journal of Strength and Conditioning Research*, 19(2), 433-437.

- Turner, A. N., & Jeffreys, I. (2010). The stretch-shortening cycle: proposed mechanisms and methods for enhancement. *Strength and Conditioning Journal*, 32(4), 87-99.
- van der Kruk, E., & Reijne, M. M. (2018). Accuracy of human motion captures systems for sport applications; state-of-the-art review. *European Journal of Sport Science*, 18(6), 806-819. DOI: 10.1080/17461391.2018.1463397.
- Verkhoshansky, Y. V., & Siff, M. C. (2009). Supertraining (6th ed.). Denver, CO: Supertraining Institute.
- Vorobyev, A. N. (1978). A Textbook on Weightlifting (W. J. Brice, Trans.). Budapest, Hungary: International Weightlifting Federation.
- Wagner, J. F. (2018). About motion measurement in sports based on gyroscopes and accelerometers - an engineering point of view. *Gyroscopy and Navigation*, 9(1), 1-18.
- Wei, G., Tian, F., Tang, G., & Wang, C. (2012). A wavelet-based method to predict muscle forces from surface electromyography signals in weightlifting. *Journal of Bionic Engineering*, 9, 48-58.
- Westenburg, T. M., Smith, S. L., Conrad, T. M., & Leigh, M. R. (2004). Instrumentation to monitor dynamic performance of Olympic weightlifters. Paper presented at the The Engineering of Sport 5, Davis, California.
- Whitehead, P. N., Schilling, B. K., Stone, M. H., Kilgore, J. L., & Chiu, L. Z. (2014). Snatch technique of United States national level weightlifters. *Journal of Strength and Conditioning Research*, 28(3), 587-591.

Winchester, J. B., Erickson, T. M., Blaak, J. B., & McBride, J. M. (2005). Changes in bar-path kinematics and kinetics after power-clean training. *Journal of Strength and Conditioning Research*, 19(1), 177.

Wood, G. A. (1982). Data smoothing and differentiation procedures in biomechanics. *Exercise and Sport Sciences Reviews*, 10, 308-362.

Zahran, K., Vieten, M., & Riehle, H. (2002). Quantification of technical skills in weightlifting. Paper presented at the XX International Symposium on Biomechanics in Sport, Cáceres, Spain.

VITA

AARON J. CUNANAN

- Education: B.S. Sociology, Louisiana State University Shreveport, Shreveport, Louisiana, 2010
M.S. Kinesiology and Wellness, Louisiana State University Shreveport, Shreveport, Louisiana, 2012
Ph.D. Sport Physiology and Performance, East Tennessee State University, Johnson City, Tennessee, 2019
- Professional Experience: Graduate Assistant, Louisiana State University Shreveport, Department of Kinesiology and Health Science, 2010-2012
Assistant Coach, USA Weightlifting Center for High Performance and Development; Shreveport, Louisiana, 2011-2016
Adjunct Instructor, Louisiana State University Shreveport; Shreveport, Louisiana, 2012-2017
Exercise Physiologist, Eberhardt Physical Therapy; Shreveport, Louisiana, 2012-2016
Doctoral Fellow, East Tennessee State University, Department of Sport, Exercise, Recreation, and Kinesiology, 2016-2019
Assistant Coach, U.S. Olympic Training Site at East Tennessee State University; Johnson City, Tennessee, 2016-2019
- Publications: Suarez DG, Wagle JP, Cunanan AJ, Sausaman RW, & Stone HM. (2019). Dynamic Correspondence of Resistance Training to Sport: A Brief Review. *Strength and Conditioning Journal*, DOI: 10.1519/SSC.0000000000000458.
- Wagle JP, Carroll KM, Cunanan AJ, Wetmore A, Taber CB, DeWeese BH,...& Stone MH. (2018). Preliminary Investigation into the Effect of ACTN3 and ACE Polymorphisms on Muscle and Performance Characteristics. *Journal of Strength and Conditioning Research*, DOI: 10.1519/JSC.0000000000002809.
- Wagle JP, Cunanan AJ, Carroll KM, Sams ML, Wetmore A, Bingham GE,...& Stone MH. (2018). Accentuated Eccentric Loading and Cluster Set Configurations in the Back Squat: a Kinetic and Kinematic Analysis. *Journal of Strength and Conditioning Research*, 33(7S), S19-S25, DOI: 10.1519/JSC.0000000000002677.

Wagle JP, Taber CB, Carroll KM, Cunanan AJ, Sams ML, Wetmore A,...& Stone MH. (2018). Repetition-to-Repetition Differences Using Cluster and Accentuated Eccentric Loading in the Back Squat. *Sports*, 6(3), pii: E59. DOI: 10.3390/sports6030059.

Cunanan AJ, DeWeese BH, Wagle JP, Carroll KM, Sausaman R, Hornsby III WG,...& Stone MH. (2018). The General Adaptation Syndrome: A Foundation for the Concept of Periodization. *Sports Medicine*, 48(4), 787-797.

Wagle JP, Taber CB, Cunanan AJ, Bingham GE, Carroll KM, DeWeese BH,...& Stone MH. (2017). Accentuated Eccentric Loading for Training and Performance: A Review. *Sports Medicine*, 47(12), 2473-2495.

Wagle JP, Carroll KM, Cunanan AJ, Taber CB, Wetmore A, Bingham GE,...& Stone MH. (2017). Comparison of the Relationship between Lying and Standing Ultrasonography Measures of Muscle Morphology with Isometric and Dynamic Force Production Capabilities. *Sports*, 5(4), pii: E88. DOI: 10.3390/sports5040088.

Honors and Awards:

2017-2018 Graduate Assistant of the Year, Department of Sport, Exercise, Recreation, and Kinesiology, East Tennessee State University, 2018