

East Tennessee State University

Digital Commons @ East Tennessee State University

ETSU Faculty Works

Faculty Works

2-22-2019

The Influence of Immigrant Generation on Obesity Among Asian Americans in California from 2013 to 2014

Shaoqing Gong
Jiaotong University

Kesheng Wang
East Tennessee State University

Ying Li
East Tennessee State University, liy005@etsu.edu

Arsham Alamian
East Tennessee State University, alamian@etsu.edu

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>



Part of the [Environmental Public Health Commons](#), and the [Epidemiology Commons](#)

Citation Information

Gong, Shaoqing; Wang, Kesheng; Li, Ying; and Alamian, Arsham. 2019. The Influence of Immigrant Generation on Obesity Among Asian Americans in California from 2013 to 2014. *PLoS ONE*. Vol.14(2). <https://doi.org/10.1371/journal.pone.0212740> ISSN: 1932-6203

This Article is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

The Influence of Immigrant Generation on Obesity Among Asian Americans in California from 2013 to 2014

Copyright Statement

© 2019 Gong et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

2-22-2019

The Influence of Immigrant Generation on Obesity Among Asian Americans in California from 2013 to 2014

Shaoqing Gong
Jiaotong University

Kesheng Wang
East Tennessee State University

Ying Li
East Tennessee State University, liy005@etsu.edu

Arsham Alamian
East Tennessee State University, alamian@etsu.edu

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>

Citation Information

Gong, Shaoqing; Wang, Kesheng; Li, Ying; and Alamian, Arsham. 2019. The Influence of Immigrant Generation on Obesity Among Asian Americans in California from 2013 to 2014. *PLoS ONE*. Vol.14(2). <https://doi.org/10.1371/journal.pone.0212740> ISSN: 1932-6203

This Article is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

The Influence of Immigrant Generation on Obesity Among Asian Americans in California from 2013 to 2014

Copyright Statement

© 2019 Gong et al. This document was originally published in *PeerJ*.


RESEARCH ARTICLE

The influence of immigrant generation on obesity among Asian Americans in California from 2013 to 2014

Shaoqing Gong^{1*}, Kesheng Wang², Ying Li³, Arsham Alamian²

1 Institute of Health Administration and Policy, School of Public Policy and Administration, Xi'an Jiaotong University, Xi'an, Shaanxi Province, China, **2** Department of Biostatistics and Epidemiology, College of Public Health, East Tennessee State University, Johnson City, Tennessee, United States of America, **3** Department of Environment Health, College of Public Health, East Tennessee State University, Johnson City, Tennessee, United States of America

* gongshaoqing@mail.xjtu.edu.cn



OPEN ACCESS

Citation: Gong S, Wang K, Li Y, Alamian A (2019) The influence of immigrant generation on obesity among Asian Americans in California from 2013 to 2014. PLoS ONE 14(2): e0212740. <https://doi.org/10.1371/journal.pone.0212740>

Editor: Catherine Mary Schooling, CUNY, UNITED STATES

Received: January 24, 2018

Accepted: February 10, 2019

Published: February 22, 2019

Copyright: © 2019 Gong et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: We thank the support of School of Public Policy and Administration, Xi'an Jiaotong University and College of Public Health of East Tennessee State University. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Objectives

We aimed to examine the association between immigrant generation and obesity among Californian adults and Asian Americans.

Methods

We pooled weighted data ($n = 2,967$) on Asian Americans from the 2013–2014 California Health Interview Survey. Overweight and obesity were defined using body mass indices (BMI) of 25 kg/m² and 30 kg/m², respectively, in non-Asians, compared with BMI of 23 kg/m² (for being overweight) and 27.5 kg/m² (for being obese) in Asians. First-generation or immigrant Asian Americans were defined as those born outside of the U.S. Second-generation Asian Americans were defined as those born in the U.S. with at least one foreign-born parent. All other Asian participants were classified as third-generation or higher. Multiple logistic regression analyses were used with adjustment for age, sex, family income, smoking status, marital status, education, physical activity, and fast food consumption.

Results

Overall, 23.3% of the Asian population was obese, and 40.0% was overweight. The percentage of 1st, 2nd, and 3rd generation were 72.7%, 22.6%, and 4.6%, respectively. Overall, 1st generation of Asians had lower odds of being obese compared to Whites (OR = 0.34, 95% CI = 0.26–0.45). Multiple logistic regression analyses showed that overall, 2nd generation (OR = 1.69, 95%CI = 1.10–2.60) and 3rd generation (OR = 2.33, 95%CI = 1.29–4.22) Asians had higher odds of being obese compared to 1st generation Asians. Among Chinese, compared to the 1st generation, the 3rd generation had increased likelihood of being obese (OR = 6.29, 95%CI = 2.38–16.6).

Conclusion

Compared to Whites, Hispanics, and Blacks, Asian immigrants are less likely to be obese. Among Asians, 2nd and 3rd generations were more likely to be obese compared to 1st generation. The obesity rate seems to increase the longer Asian immigrants remain in the U.S.

Introduction

Asian immigrants first came to the United States (U.S.) in significant numbers more than a century and a half ago, mainly as low-skilled male laborers who mined, farmed and built the railroads. They endured generations of officially sanctioned racial prejudice. A first major wave of Asian immigration occurred in the late 19th century, primarily in Hawaii and the West Coast. Since the Immigration and Nationality Act of 1965, Asian American demographics changed rapidly. Over the decades, this modern wave of immigrants from Asia has increasingly become more skilled and educated; a new wave of new immigrants to the U.S. in 2010 were mainly from Asia [1]. In 2014, among a total 42.4 million immigrants in the U.S., Asian Americans accounted for 42.4% [2]. The immigrant population continues to grow. Of all the immigrant populations, Asian Americans were the second largest immigrant group, accounting for 28% of all foreign born populations [3]. In 2011, 79% of Asian Americans aged 18 years and older were foreign-born [4]. Asian Americans are one of the fastest growing populations in the U.S. The Asian population in the U.S. increased by 46% between 2000 and 2010 [5] and will double in population size with a projected increase to more than 43 million people by 2050 [6]; their numbers as a proportion of the U.S. population are projected to grow from 5.8% in 2011 to 9% by 2050 [5].

Immigrants' health is a function of influences derived from the sending country, the receiving country, and the migration and resettlement experience itself [7]. The existence of a healthy immigrant effect, i.e., that immigrants are on average healthier than the native born, is a widely cited phenomenon across a multitude of literatures including epidemiology and the social sciences [8]. For example, Tu and Colleagues have shown that immigrants to Canada experience lower risk of CVD compared to long-term residents, indicating a healthy immigrant effect, although this protective effect appears to erode slightly over time [7]. Among the few studies that focused on Asians, previous studies that imply the healthy immigrant effect show some evidence that Chinese immigrants have a lower asthma prevalence [9, 10], lower body mass index [10], and overall better general health [11, 12] than native born residents. In the U.S., obesity has become an epidemic over the past two decades [13, 14]. According to the 2013–2014 National Health and Nutrition Examination Survey (NHANES) the age-adjusted prevalence of obesity in the U.S. was 35.0% for men and 40.4% for women [14]. Compared to 2001, obesity prevalence in 2011–2012 has increased among Chinese (3.8% versus 6.1%), Japanese (9.0% versus 15.0%), Filipino (8.8% versus 18.7%), Vietnamese (3.4% versus 6.8%), Southeast Asians (5.8% versus 19.7%), and multiple Asian groups (5.8% versus 12.3%) [15]. Increase in obesity prevalence has also been observed in other ethnic groups (e.g., Whites, Blacks) [16]. However, to date, there is very limited research examining the influence of immigrant generation on obesity among Asian Americans. Identifying factors that have propelled the increase of obesity among Asian Americans is important if we are to effectively combat and prevent obesity nationwide [17].

Research has shown that compared to their native-born cohorts, newly arrived immigrants have better health, but their health declines the longer they remain in the U.S. and become

more acculturated [18]. Acculturation is the process by which a minority group adopts the cultural lifestyle and behaviors of the host country [19]. Studies show that some immigrants might possess a health advantage due to migration selectivity and a protective native culture; this protective buffer dissipates by the third generation [20]. Antecol (2006) examined the association between length of residency of immigrants and their obesity. The author found that within 10–15 years after immigration study participants' BMIs became similar as those of U.S. born; immigrants' BMIs were 2–5% lower than those of U.S. born once they arrived in the U.S. [21].

The aim of this research is to analyze the association of immigration generation and obesity among Asian Americans in California. We analyzed data from a large population-based sample that is representative of Asian ethnic groups: Chinese, Filipino, Vietnamese, Korean, and Japanese [22]. In contrast, most national surveys on health sample a small number of Asian Americans. Further, we pooled weighted data from the latest cycle of California population, i.e., 2013–2014. The results of this study can help to refine the diverse risk profile for obesity among Asian Americans while also contributing to the overall understanding of the impact of migration on chronic health conditions.

Methods

Participants

The California Health Interview Survey (CHIS) is a population-based telephone survey designed to produce health data representative of California's non-institutionalized population. The survey has been conducted biennially since 2001 and continually beginning 2011, and covers comprehensive aspects of health. A detailed technical report of the survey can be found at website <http://healthpolicy.ucla.edu/chis/Pages/default.aspx>. For CHIS 2013–2014, the landline/list sample household response rate was 14.8 percent; the cell sample household response rate was 16.6 percent. Data on Asian Americans ($n = 2,967$) of 40,240 whole sample (also including Whites (25,643), Hispanics ($n = 7,996$), Blacks ($n = 1,764$)) were obtained from CHIS 2013–2014. CHIS oversampled Asian Americans to increase the precision of estimates for those ethnic groups. Interviews were conducted in five languages [English, Spanish, Chinese (both Mandarin and Cantonese), Vietnamese, and Korean].

We used data collected from Californian adult participants aged 18 to 85 and restricted the sample to Asian descent living in the U.S. and U.S.-born other ethnic groups (i.e., Whites, Hispanics, and Blacks), where they were used as the reference group in our study, respectively. We then excluded the following subjects: 1) Asians who self-identified as being of an ethnicity other than the following large ethnic groups of Asian Americans (Chinese, Filipino, Vietnamese, Korean, and Japanese) or as being of multiple Asian sub-ethnicities; 2) foreign-born Asians who were missing information on the nativity of either parent for studying obesity as outcome variables.

Study variables

Obesity. BMI was calculated using self-reported height and weight. In order to account for racial differences in body fat percentage at the same BMI level, we applied race-specific BMI standards. We used BMI of 25 kg/m^2 and 30 kg/m^2 as the cut-off points for the three weight categories (underweight/normal weight ($\text{BMI} < 25 \text{ kg/m}^2$), overweight ($25 \text{ kg/m}^2 \leq \text{BMI} < 30 \text{ kg/m}^2$), and obese ($\text{BMI} > 30 \text{ kg/m}^2$) in Whites, compared with BMI of 23 kg/m^2 (for being overweight) and 27.5 kg/m^2 (for being obese) in Asians [23].

Generational status. First-generation or immigrant Asian Americans were defined as those born outside of the U.S. Second-generation Asian Americans were defined as those born

in the U.S. with at least one foreign-born parent. All other Asian participants were classified as third-generation or more [20].

Covariates. Age was treated as a continuous variable. Sex and race/ethnicity were treated as categorical variables. Due to small sample size for some subgroups, race/ethnicity included Chinese, Filipino, Vietnamese, Korean, and Japanese. Family income was reported by the adult respondent, usually a parent, and was examined as a percentage of the federal poverty level (FPL), which adjusted for total household income and number of members in the household. Family income was categorized as below 100%, 100% to 299%, or 300% and above of the FPL. Smoking status was defined as current smoker or not current smoker. Marital status was defined as married, never married, or other. Education attainment included three categories: high school, college, and graduate. Physical activity was defined as walking at least 10 minutes for either transportation or leisure over the past 7 days. Fast food consumption was determined by response to the following question: “In the past 7 days, how many times did you eat fast food? Include fast food meals eaten at work, at home, or at fast-food restaurants, carryout or drive through” Response categories included: never, 1–2 times, and ≥ 3 times.

Statistical analysis

To investigate the characteristics of immigrant generation (e.g., 2nd generation and 3rd generation versus 1st generation) among Asian Americans in California, and the association with obesity, we merged public-use data files of adult samples from the CHIS 2013–2014. All analyses were accounted for complex sampling designs and non-response by employing weights developed by CHIS. Weights are included with the data files and are based on the State of California’s Department of Finance population estimates and projections, adjusted to remove the population living in group quarters (such as nursing homes, prisons, etc.) and thus not eligible to participate in CHIS. When the weights are applied to the data, the results represent California’s residential population during that year for the age group corresponding to the data file in use. Additional information on how to use the CHIS sampling weights, including sample code, is available at: <http://healthpolicy.ucla.edu/chis/analyze/Pages/sample-code.aspx>.

The descriptive statistics for participant characteristics were reported by Asian ethnicity. Wald Chi-square test was used to examine categorical variables while analysis of variance was used to compare continuous variables between racial/ethnic groups. Potential confounders were controlled if they fulfilled the following three basic properties: 1) be associated with the exposure, 2) be a risk factor for the outcome, and 3) not in the causal pathway of interest. Confounders will also always be associated with the exposure; however, the relationship may or may not be causal, i.e., confounder does not necessarily need to be a common cause of both immigrant generation and obesity in this study [24]. We further fit logistic regression models to obtain adjusted odds ratios (AORs) with 95% confidence interval (CI) of obesity for Asians from different subgroups versus other races (i.e., Whites, Hispanics, and Blacks, as the reference group, respectively). Furthermore, we examined the association between immigrant generation and obesity among Asian Americans using multiple logistic regression analyses that included variables with p value significant at 0.2 in univariate analysis. All tests of significance were two-tailed, with alpha level of 0.05. Analyses were performed using survey data analysis procedures in SAS (version 9.4; SAS Institute Inc., Cary, NC).

Results

Characteristics of Asian Americans

[Table 1](#) describes the sample characteristics of Asian adults in California by ethnicity. Overall, 23.3% of the participants were obese, 40.0% were overweight, and 2.4% were underweight. The

Table 1. Sample characteristics of Asian adults in California by ethnicity: CHIS 2013–2014 (n = 2,967).

	Overall	Chinese	Korean	Japanese	Filipino	Vietnamese
Weight status, n (%)						
Underweight (0–18.49)	131 (2.4)	47 (3.2)	9 (0.5)	22 (5.2)	8 (2.1)	33 (3.1)
Healthy weight (18.5–22.99)	1371 (34.3)	466 (38.0)	186 (44.2)	179 (36.4)	161 (25.6)	232 (50.3)
Overweight (23.0–27.49)	1530 (40.0)	435 (41.5)	190 (42.5)	193 (31.9)	236 (38.5)	216 (28.8)
Obesity (27.5+)	778 (23.3)	180 (17.3)	60 (12.8)	153 (26.5)	197 (33.8)	73 (17.9)
Immigrant generation of Asians, n (%)						
1 st generation	2672 (72.7)	796 (71.7)	387 (81.3)	118 (27.1)	394 (72.3)	509 (79.7)
2 nd generation	596 (22.6)	198 (22.0)	33 (16.1)	168 (34.8)	95 (26.0)	39 (20.1)
3 rd generation	224 (4.6)	49 (6.3)	9 (2.6)	159 (38.1)	12 (1.8)	1 (0.2)
Sex, n (%)						
Male	1673 (46.9)	477 (43.8)	197 (43.6)	216 (38.4)	248 (48.3)	249 (48.5)
Female	2137 (53.1)	651 (56.2)	248 (56.4)	331 (61.6)	354 (51.7)	305 (51.5)
Age						
18–44 years	1135 (56.2)	350 (60.9)	104 (51.6)	82 (41.5)	220 (52.6)	124 (50.0)
45–64 years	1443 (30.6)	467 (29.0)	131 (27.2)	214 (35.7)	213 (32.1)	233 (36.2)
65 years or above	1232 (13.2)	311 (10.1)	210 (21.2)	251 (22.8)	169 (15.4)	197 (13.8)
Family income, n (%)						
<100% FPL	754 (14.6)	204 (13.5)	105 (15.6)	40 (3.9)	83 (13.7)	244 (25.9)
100%–299% FPL	1146 (30.9)	336 (31.4)	177 (42.9)	119 (24.2)	194 (31.0)	184 (32.0)
≥300% FPL	1910 (54.5)	588 (55.1)	163 (41.5)	388 (71.9)	325 (55.3)	126 (42.2)
Smoking status, n (%)						
Current smoker	257 (8.8)	55 (9.3)	45 (14.1)	41 (5.2)	51 (9.9)	41 (9.0)
Not current smoker	3553 (91.2)	1073 (90.7)	400 (85.9)	506 (94.8)	551 (90.1)	513 (91.0)
Marital status, n (%)						
Married	2233 (54.7)	684 (56.9)	252 (57.2)	288 (52.2)	296 (46.8)	344 (57.2)
Others	815 (12.3)	212 (10.4)	124 (12.7)	158 (12.9)	149 (18.0)	112 (11.3)
Never married	762 (32.9)	232 (32.7)	69 (30.0)	101 (34.9)	157 (35.2)	98 (31.5)
Education, n (%)						
High school	1083 (27.2)	315 (30.1)	152 (33.9)	100 (24.5)	111 (18.0)	314 (44.1)
College	1939 (53.1)	522 (46.4)	223 (47.9)	324 (58.5)	426 (70.5)	200 (42.8)
Graduate	788 (19.7)	291 (23.5)	70 (18.2)	123 (17.1)	65 (11.5)	40 (13.2)
Physical activity,^a n (%)						
Yes	3058 (79.3)	910 (78.4)	360 (86.8)	411 (80.1)	488 (82.3)	444 (77.8)
No	752 (18.7)	218 (21.6)	85 (13.2)	136 (19.9)	114 (17.7)	110 (22.2)
Fast food, n (%)						
Never	1982 (43.8)	617 (48.5)	228 (39.9)	245 (41.7)	226 (35.6)	399 (55.6)
1–2 times	1340 (37.2)	389 (35.3)	165 (48.2)	220 (38.5)	251 (36.2)	115 (29.7)
≥3 times	488 (19.0)	122 (16.2)	52 (11.9)	82 (19.7)	125 (28.3)	40 (14.7)

Abbreviation: FPL = federal poverty level.

<https://doi.org/10.1371/journal.pone.0212740.t001>

percentage of 1st, 2nd, and 3rd generation were 72.7%, 22.6%, and 4.6%, respectively. The majority of the Asian population was aged 18–44 years, had high family income, was not current smoker, was married, had college or graduate degrees, engaged in physical activity, and consumed fast foods during past seven days. Among Asian subgroups, the prevalence of obesity was highest in Filipino (33.8%), followed by Japanese (26.5%), Vietnamese (17.9%), Chinese (17.3%), and Korean (12.8%) (p<0.0001). Koreans had the highest percentage of 1st

generation (81.3%), while Japanese had the lowest percentage (27.1%). As for the 2nd generation, the percentage was highest in Japanese (34.8%) and lowest in Koreans (16.1%). As for the 3rd generation, Japanese had the highest percentage (38.1%) while Vietnamese had the lowest percentage (0.2%) (all $p < 0.05$).

Multiple logistic regression analyses for the association between immigration generations and obesity among Californian adults

Table 2 shows results from multiple logistic regression analyses for the association between immigration generations and obesity among Californian adults (Asian Americans vs. Whites, Hispanics, and Blacks, respectively). Compared to Whites, overall, 1st generation of Asians had lower odds of being obese (OR = 0.34, 95% CI = 0.26–0.45). Due to small sample size, Asians were divided into Chinese and other Asian groups. Among Chinese, compared to Whites, 1st generation (OR = 0.26, 95% CI = 0.14–0.46) and 2nd generation (OR = 0.24, 95% CI = 0.09–0.64) had reduced likelihood of being obese, but such association disappeared for 3rd generation (OR = 1.08, 95% CI = 0.33–3.49). Among other Asian groups, compared to Whites, being of 1st generation (OR = 0.36, 95% CI = 0.25–0.53) was inversely related to obesity, but such association disappeared for 2nd generation (OR = 0.74, 95% CI = 0.38–1.47) and 3rd generation (OR = 0.78, 95% CI = 0.37–1.63).

Compared to Hispanics, overall, 1st generation (OR = 0.24, 95% CI = 0.18–0.32) and 2nd generation (OR = 0.46, 95% CI = 0.26–0.81) of Asians had lower odds of being obese. Among Chinese, compared to Hispanics, 1st generation (OR = 0.16, 95% CI = 0.09–0.28) and 2nd generation (OR = 0.20, 95% CI = 0.08–0.52) had reduced likelihood of being obese, but such association disappeared for 3rd generation (OR = 0.78, 95% CI = 0.29–2.11). Among other Asian groups, compared to Hispanics, being of 1st generation (OR = 0.26, 95% CI = 0.18–0.38) was associated with decreased odds of being obese, but such association disappeared for 2nd generation (OR = 0.55, 95% CI = 0.28–1.09) and 3rd generation (OR = 0.62, 95% CI = 0.29–1.34).

Compared to Blacks, overall, 1st generation (OR = 0.16, 95% CI = 0.12–0.22), 2nd generation (OR = 0.26, 95% CI = 0.14–0.46), and 3rd generation (OR = 0.41, 95% CI = 0.22–0.78) of

Table 2. Multiple logistic regression analyses for the association between immigration generations and obesity among Californian adults, CHIS 2013–2014.

	All Asians AOR (95% CI)	Chinese AOR (95% CI)	Other Asians AOR (95% CI)
Asians vs. Whites*			
1 st generation	0.34 (0.26–0.45)	0.26 (0.14–0.46)	0.36 (0.25–0.53)
2 nd generation	0.59 (0.34–1.04)	0.24 (0.09–0.64)	0.74 (0.38–1.47)
3 rd generation	0.86 (0.45–1.64)	1.08 (0.33–3.49)	0.78 (0.37–1.63)
Asians vs. Blacks*			
1 st generation	0.16 (0.12–0.22)	0.11 (0.06–0.21)	0.17 (0.11–0.26)
2 nd generation	0.26 (0.14–0.46)	0.11 (0.04–0.29)	0.32 (0.16–0.66)
3 rd generation	0.41 (0.22–0.78)	0.46 (0.17–1.28)	0.39 (0.19–0.81)
Asians vs. Hispanics*			
1 st generation	0.24 (0.18–0.32)	0.16 (0.09–0.28)	0.26 (0.18–0.38)
2 nd generation	0.46 (0.26–0.81)	0.20 (0.08–0.52)	0.55 (0.28–1.09)
3 rd generation	0.68 (0.35–1.29)	0.78 (0.29–2.11)	0.62 (0.29–1.34)

Abbreviations: AOR, adjusted odds ratio; CI, confidence interval.

*The multiple regression analyses were used to adjust for age, sex, family income, smoking status, marital status, education, physical activity, and fast food consumption.

<https://doi.org/10.1371/journal.pone.0212740.t002>

Asians had lower odds of being obese. Among Chinese, compared to Blacks, 1st generation (OR = 0.11, 95% CI = 0.06–0.21) and 2nd generation (OR = 0.11, 95% CI = 0.04–0.29) had reduced likelihood of being obese, but such association disappeared for 3rd generation (OR = 0.46, 95% CI = 0.17–1.28). Among other Asian groups, compared to Blacks, 1st generation (OR = 0.17, 95% CI = 0.11–0.26), 2nd generation (OR = 0.32, 95% CI = 0.16–0.66) and 3rd generation (OR = 0.39, 95% CI = 0.19–0.81) were associated with decreased odds of being obese.

Multiple logistic regression analyses for the association between immigration generations and obesity among Asian adults

Table 3 shows results from multiple logistic regression analyses for the association between immigration generations and obesity among Asian adults. Overall, 2nd generation (OR = 1.69, 95% CI = 1.10–2.60) and 3rd generation (OR = 2.33, 95% CI = 1.29–4.22) were associated with higher odds of being obese compared to 1st generation. Among Chinese, compared to 1st generation, 3rd generation had increased likelihood of being obese (OR = 6.29, 95% CI = 2.38–16.6). Among other Asian groups, compared to 1st generation, 2nd generation were associated with increased likelihood of being obese (OR = 1.99, 95% CI = 1.14–3.47).

Discussion

This study is among the very few studies to examine the association between immigrant generation and obesity among Asian Americans. We found that, using WHO Asian cut points for obesity, overall, 23.3% of the Asians were obese, and 40.0% were overweight. The percentage of 1st, 2nd, and 3rd generation in the study were 72.7%, 22.6%, and 4.6%, respectively. Compared to other races (i.e., Whites, Hispanics, and Blacks), Asians were less likely to be obese, especially for 1st generation of Asian Americans. Although 2nd and 3rd generations of Asians were also associated with reduced obesity prevalence as compared to other races, the magnitude of the association decreased compared to the 1st generation of Asians. For example, 1st generation, 2nd generation, and 3rd generation of Asians had lower odds of being obese compared to Blacks. This finding is further supported by the observation that among Asian adults, 2nd generation and 3rd generation were associated with higher odds of being obese compared to 1st generation. These findings are consistent with previous studies, e.g., compared to their native-born cohorts, newly arrived immigrants' health declines the longer they remain in the U.S. and become more acculturated [17]. All the evidence suggests an increased risk of being obese among immigrants due to lifestyle changes closer to the U.S. born population with

Table 3. Multiple logistic regression analyses for the association between immigrant generation and obesity among Asian Americans, CHIS 2013–2014.

	All Asians AOR (95% CI)	Chinese AOR (95% CI)	Other Asians AOR (95% CI)
Immigrant generation*			
1 st generation (ref)			
2 nd generation	1.69 (1.10–2.60)	1.47 (0.57–3.80)	1.99 (1.14–3.47)
3 rd generation	2.33 (1.29–4.22)	6.29 (2.38–16.6)	1.71 (0.85–3.44)

Abbreviations: AOR, adjusted odds ratio; CI, confidence interval.

*The multiple regression analyses were used to adjust for age, sex, family income, smoking status, marital status, education, physical activity, and fast food consumption.

<https://doi.org/10.1371/journal.pone.0212740.t003>

increasing generation, i.e., indicating that acculturation to U.S. lifestyles contributes to their heightened obesity risk.

Asian Americans have been reported to be less obese than other major U.S. ethnic population groups [25]; the latter may be due to the high proportion of immigrants in the Asian American population. Studies on obesity in Asian Americans have found that compared to U.S.-born Asian Americans, foreign-born Asian Americans are significantly less overweight and obese, though increased duration of residence in the U.S. among foreign-born Asian Americans is correlated with increased obesity [26–28]. Studies of Asian Americans by immigrant generation have observed a trend of increasing obesity with later generations [25, 29]. Although it is assumed that U.S. environmental characteristics and acculturation to western lifestyles (e.g., diet and exercise) may explain the phenomenon, very limited studies have been conducted to link immigrant generation and subsequent obesity in Asian American adults. The similarities and differences with findings of previous studies on migrant status and obesity may be due to study designs (e.g., cross-sectional, cohort studies), sample size for the study population, and measures of study variables (e.g., definition of migrants status, cut-off points used for defining obesity among Asians).

Although immigrants likely arrive in the U.S. with adverse social and economic profile, they have health advantage, which may reflect migration selectivity [27, 30, 31] or the protective culture of immigrants. This encourages healthy behaviors and strong social support systems [32, 33]. Nevertheless, immigrants and their subsequent generations lose at least some of this initial health advantage [21, 34]. With regard to obesity, the differences in its prevalence and related risk factors have been reported between immigrants and U.S. born population [29], leading to different obesity prevalence between Asians and other races.

Immigration selection, Asian culture (diet and life style), and a tight social network among first-generation families may have a protective role in health outcome, e.g., they may protect male immigrants from an increased risk of type 2 diabetes in the U.S. [20]. BMI is, in particular, lower in many sending countries than in North America or Western Europe [25]. This study is mainly focused on Asians, and compared Asians with other ethnic groups (Whites, Hispanics, and Blacks, as the reference group, respectively). For example, compared to Asians immigrants, White immigrants may be less likely to change significant diet and life style in later generations. Thus, using Whites or other ethnic groups as reference groups that did not exclude their immigrants may not significantly bias the results. Acculturation is considered to likely modify health and behavioral risks of immigrants, leading to decreased odds of being obese from 1st to 3rd generations [29]; this is consistent with a recent study where health advantage in some immigrants due to migration selectivity and a protective native culture seem to dissipate by later generations [20]. The association of immigrant generation and obesity was also observed in Chinese and other Asian groups.

In addition, some studies have suggested a positive association between immigrant generation and other health outcomes. For example, a study found that immigrant generation is associated with increased diabetes risk among adults of Mexican origin. Compared with 1st generation Mexican immigrants, U.S. born second- and third-generation individuals had higher diabetes risk [35].

Immigrants' advantage in body composition is driven by the selection of healthy people to immigrate to the U.S. and the selection of unhealthy people to return to their home country. Immigrants' initial body composition advantage, however, is gradually eroded by the exposure to the high risk of obesity in the U.S. [36]. Previous studies have reported that immigrants have a lower BMI than the native born [27, 37], and further document that immigrants who have spent less than 10 years in the U.S. have a significant lower BMI than those with a longer duration of residence [27]. Compared with duration of migration, age at immigration is also

important. It is likely that healthy behaviors (e.g., less exposure to smoking, better diets, and more light or moderate intensity physical activity) were culturally ingrained prior to immigration [10]. Immigrants may have developed these habits before immigrating, and likely have retained them if they immigrate at older age and live in the potentially protective ethnic enclave, which could be protective because they facilitate access to services, food, and social opportunities [38]. Roshania et al. (2012) shows that immigrants younger than 20 at arrival in the U.S. may be at higher risk of overweight/obesity with increasing duration of residence than those who arrive at later ages. Obesity prevention among young U.S. immigrants should be a priority [39].

The strengths of this study include the large CHIS sample that used people from many ethnic groups and provides health-related information for most large and small racial and ethnic population in particular Asian subgroups. In addition, CHIS is conducted in different languages to represent diverse population. Some limitations also need our attention. First, our analyses were limited to the CHIS public use file, and thus we did not have access to information such as parents' national origins or documentation status. Second, CHIS only interviews respondents in residential settings, which do not include homeless individuals and people living in group quarters such as group homes, dormitories, jails, and prisons. Third, the original sample size is small, and due to small sample sizes for some subgroups, Asians group was only dichotomized into Chinese and other Asians as a whole, providing limited information for Japanese, Koreans, Filipino, and Vietnamese. Fourth, our data are subject to self-report bias. Self-reported height and weight used in this study may underestimate the prevalence of obesity [40]. Measured obesity rates were reported to be dramatically higher compared to self-reported estimates [41] and were consistent with overall national obesity prevalence of 34% from NHANES 2007–2008 [42], highlighting the importance of using measured height and weight when determining population estimates. Fifth, the use of a new definition of a confounder may need some attention [43, 44], which does go deeper and beyond the methods employed in the current manuscript. The new definition of a confounder pertains to approaches that include creating a framework of concepts and methods for colliders, instrumental variables, d-separations, backdoor paths and, notably, Directed Acyclic Graphs [43]. This is because the proposed new definition of a confounder is said to 1) better distinguish between mediators and confounders, so that direct and indirect effects are more clear and 2) help limit adjusting for a mediator that generate selection bias, as in the birth weight paradox [43]. Additional benefits of using the new methods and definitions of confounders are described elsewhere [44]. Finally, CHIS is a California-based random-digit dial survey of population health. Low response rates for the years included in this study limit generalizability. Additionally, our results are not generalizable to the U.S. population since CHIS only samples the California population.

Conclusion

Compared to non-immigrants, immigrants are more likely to have obesity. Among Asians, 2nd and 3rd generations were associated with higher likelihood of being obese compared to 1st generation. The obesity rate seems to increase the longer Asians remain in the U.S. due to being more acculturated.

Supporting information

S1 File.
(RAR)

Acknowledgments

The authors are grateful to the support of data from the 2013–2014 California Health Interview Survey. We also thank the support of School of Public Policy and Administration, Xi'an Jiao-tong University and College of Public Health of East Tennessee State University.

Author Contributions

Conceptualization: Shaoqing Gong, Ying Li, Arsham Alamian.

Formal analysis: Shaoqing Gong, Kesheng Wang.

Supervision: Kesheng Wang, Ying Li, Arsham Alamian.

Writing – original draft: Shaoqing Gong.

Writing – review & editing: Kesheng Wang, Ying Li, Arsham Alamian.

References

1. Pew Research Center. "The Rise of Asian Americans". Pew Research Center's Social & Demographic Trends Project. June 19, 2012. Updated Edition, April 4, 2013. <http://www.pewsocialtrends.org/2012/06/19/the-rise-of-asian-americans/>
2. Mossaad, N. 2016. U.S. lawful permanent residents: 2014. Annual Flow Report. Office of Immigration Statistics. U.S. Department of Homeland Security. https://www.dhs.gov/sites/default/files/publications/LPR%20Flow%20Report%202014_508.pdf
3. Grieco EM, Acosta YD, Patricia de la Cruz G, Gambino C, Gryn T, Larsen, et al. 2012. The foreign-born population in the United States: 2010 American community survey reports (ACS-19). www.census.gov/prod/2012pubs/acs-19.pdf
4. U.S. Census Bureau. 2015. American FactFinder—Results. Washington, DC: U.S. Census Bureau. http://factfinder.census.gov/faces/tableservices/jsf/pages/productview.xhtml?pid=ACS_11_1YR_C05003D&prodType=table
5. Pew Research Center. 2008. U.S. Population Projections: 2005–2050. Washington, DC: Pew Hispanic Center, 2008. <http://www.pewhispanic.org/2008/02/11/us-population-projections-2005-2050/>
6. Yi SS, Kwon SC, Wyatt L, Islam N, Trinh-Shevrin C. Weighing in on the hidden Asian American obesity epidemic. *Prev Med*, 2015; 73: 6–9. <https://doi.org/10.1016/j.ypmed.2015.01.007> PMID: 25602909
7. Tu JV, Chu A, Rezai MR, et al. Incidence of major cardiovascular events in immigrants to Ontario, Canada: the CANHEART immigrant study. *Circulation*. 2015; 132:1549–1559. <https://doi.org/10.1161/CIRCULATIONAHA.115.015345> PMID: 26324719
8. Kennedy S, Kidd MP, McDonald JT, Biddle N. The healthy immigrant effect: Patterns and evidence from four countries. *Int. Migration & Integration*. 2015; 16(2):317–332.
9. Wang H-Y, Wong GWK, Chen Y-Z, Ferguson AC, Greene JM, Ma Y, et al. Prevalence of asthma among Chinese adolescents living in Canada and in China. *CMAJ*. 2008; 179:1133–1142. <https://doi.org/10.1503/cmaj.071797> PMID: 19015564
10. Corlin L, Woodin M, Thanikachalam M, Lowe L, Brugge Q. Evidence for the healthy immigrant effect in older Chinese immigrants: a cross-sectional study. *BMC Public Health*. 2014; 14: 603. <https://doi.org/10.1186/1471-2458-14-603> PMID: 24928348
11. Frisbie WP, Cho Y, Hummer RA. Immigration and the Health of Asian and Pacific Islander adults in the United States. *Am J Epidemiol*. 2001; 153:372–380. PMID: 11207155
12. Sheth T, Nair C, Nargundkar M, Anand S, Yusuf S. Cardiovascular and cancer mortality among Canadians of European, south Asian and Chinese origin from 1979 to 1993 an analysis of 1.2 million deaths. *Can Med Assoc J*. 1999; 161:132–138.
13. Flegal KM, Carroll MD, Kit BK, Ogden CL. Prevalence of obesity and trends in the distribution of body mass index among US adults, 1999–2010. *JAMA*. 2012; 307(5): 491–497. <https://doi.org/10.1001/jama.2012.39> PMID: 22253363
14. Flegal KM, Kruszon-Moran D, Carroll MD, Fryar CD, Ogden CL. Trends in obesity among adults in the United States, 2005–2014. *JAMA*. 2016; 315(21): 2284–2291. <https://doi.org/10.1001/jama.2016.6458> PMID: 27272580

15. Wolstein J, Babey SH, Diamant AL. Obesity in California. Los Angeles, CA: UCLA Center for Health Policy Research, 2015. <http://healthpolicy.ucla.edu/publications/Documents/PDF/2015/obesityreport-jun2015.pdf>
16. Jackson CL, Szklo M, Yeh HC, Wang NY, Dray-Spira R, Thorpe R, et al. Black-white disparities in overweight and obesity trends by educational attainment in the United States, 1997–2008. *J Obes.* 2013;140743.
17. Desir S. Acculturation factors associated with the prevalence of obesity in immigrant children and adolescent. Thesis. College of Nursing and the Burnett Honors College. The university of Central Florida. 2015. http://etd.fcla.edu/CF/CFH0004776/Desir_Suzeline_201504_BSN.pdf
18. Van Hook J, Baker E. Big boys and little girls: gender, acculturation, and weight among young children of immigrants. *J Health Soc Behav.* 2010; 51: 200–214. <https://doi.org/10.1177/0022146510372347> PMID: 20617759
19. Schaefer SE, Salazar M, Bruhn C, Saviano D, Boushey C, Van Loan MD. Influence of Race, acculturation, and socioeconomic status on tendency toward overweight in Asian-American and Mexican-American early adolescent females. *J Immigr Minor Health.* 2009; 11(3): 188–197. <https://doi.org/10.1007/s10903-008-9150-6> PMID: 18506624
20. Huang ZJ, Zheng C. Type 2 diabetes among 6 Asian ethnic groups in California: the nexus of ethnicity, gender, and generational status. *J Health Care Poor Underserved.* 2015; 26(2 Suppl): 16–35. <https://doi.org/10.1353/hpu.2015.0061> PMID: 25981086
21. Antecol BK. Unhealthy assimilation: why do immigrants converge to American health status levels? *Demography.* 2006; 43(2): 337–360. PMID: 16889132
22. Hoeffel EM, Rastogi S, Kim MO, Shahid H. The Asian population: 2010. Washington DC: U.S. Census Bureau. 2012. www.census.gov/prod/cen2010/briefs/c2010br11.pdf
23. World Health Organization (WHO) Expert Consultation. Appropriate body-m ass index for Asian populations and its implications for policy and intervention strategies. *Lancet.* 2004; 363(9403):157–163. [https://doi.org/10.1016/S0140-6736\(03\)15268-3](https://doi.org/10.1016/S0140-6736(03)15268-3) PMID: 14726171
24. Westreich D, Greenland S. The table fallacy: presenting and interpreting confounder and modifier coefficients. *Am J Epidemiol.* 2013; 177(4):292–298. <https://doi.org/10.1093/aje/kws412> PMID: 23371353
25. Bates LM, Acevedo-Garcia D, Alegría M, Krieger N. Immigration and generational trends in body mass index and obesity in the United States: results of the National Latino and Asian American Survey, 2002–2003. *Am J Public Health.* 2008; 98(1): 70–77. <https://doi.org/10.2105/AJPH.2006.102814> PMID: 18048787
26. Kaushal N. Adversities of acculturation? Prevalence of obesity among immigrants. *Health Econ.* 2009; 18(3): 291–303. <https://doi.org/10.1002/hec.1368> PMID: 18464286
27. Goel MS, McCarthy EP, Phillips RS, Wee CC. Obesity among US immigrant subgroups by duration of residence. *JAMA.* 2004; (23): 2860–2867. <https://doi.org/10.1001/jama.292.23.2860> PMID: 15598917
28. Sanchez-Vaznaugh EV, Kawachi I, Subramanian SV, Sanchez BN, Acevedo-Garcia D. Do socioeconomic gradients in body mass index vary by race/ethnicity, gender, and birthplace? *Am J Epidemiol.* 2009; 169(9): 1102. <https://doi.org/10.1093/aje/kwp027> PMID: 19299405
29. Singh GK, Kogan MD, Yu SM. Disparities and overweight prevalence among US immigrant children and adolescents by generational status. *J Community Health.* 2009; 34: 271–281. <https://doi.org/10.1007/s10900-009-9148-6> PMID: 19333745
30. Palloni A, Arias E. Paradox lost: explaining the Hispanic adult mortality advantage. *Demography.* 2004; 41(3): 385–415. PMID: 15461007
31. Rubalcava LN, Teruel GM, Thomas D, Goldman N. The healthy migrant effect: new findings from the Mexican Family Life Survey. *Am J Public Health.* 2008; 98(1): 78–84. <https://doi.org/10.2105/AJPH.2006.098418> PMID: 18048791
32. Kawachi I, Berkman L. Social cohesion, social capital, and health.: Berkman L, Kawachi I, editors., *Social Epidemiology.* Oxford: Oxford University Press. 2000; 174–190.
33. Cattell V. Poor people, poor places, and poor health: the mediating role of social networks and social capital. *Soc Sci Med.* 2001; 52(10): 1501–1516. PMID: 11314847
34. Angel RJ, Angel JL, Diaz Venegas CD, Bonazzo C. Shorter stay, longer life: age at migration and mortality among the older Mexican-origin population. *J Aging Health.* 2010; 22(7): 914–931. <https://doi.org/10.1177/0898264310376540> PMID: 20682948
35. Afable-Munsuz A, Mayeda ER, Pérez-Stable EJ, Haan MN. Immigrant generation and diabetes risk among Mexican Americans: the Sacramento area Latino study on aging. *Am J Public Health.* 2004; 104 Suppl 2: S234–250.
36. Hao L, Kim J. Immigration and the American obesity epidemic. *Int Migr Rev.* 2009; 43(2): 237–262. <https://doi.org/10.1111/j.1747-7379.2009.00764.x> PMID: 27158177

37. Jasso G; Massey DS.; Rosenzweig MR.; SmithJP. Immigrant Health—Selectivity and Acculturation. In: Anderson Norman B.; Bulatao Randy A.; Cohen Barney, editors. *Critical Perspectives on Racial and Ethnic Differences in Health in Late Life*. Washington, DC: National Academy Press; 2004. p. 227–266.
38. Chung T. Asian American Health: discrepancies, convergence, and enclave-specific trends. In *Handb Asian Am Health Tak Not Tak Action*. Volume 2. Edited by BATERMAN W, ABESAMIS-MENDOZA N, HO-ASJOE H. Santa Barbara, CA: Greenwood Publishing Group; 2010:575–622.
39. Roshaiia R, Narayan V, Oza-Frank. Age at arrival and risk of obesity among US immigrants. *Obesity*. 2008; 16(12):2669–2675. <https://doi.org/10.1038/oby.2008.425> PMID: 18846044
40. Connor GS, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev*. 2007; 8(4): 307–326. <https://doi.org/10.1111/j.1467-789X.2007.00347.x> PMID: 17578381
41. Befort CA, Nazir N, Perri MG. Prevalence of obesity among adults from rural and urban areas of the United States: findings from NHANES (2005–2008). *J Rural Health*. 2012; 28(4), 392–397. <https://doi.org/10.1111/j.1748-0361.2012.00411.x> PMID: 23083085
42. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA*. 2010; 303(3): 235–241. <https://doi.org/10.1001/jama.2009.2014> PMID: 20071471
43. Porta M, Vineis P, Bolúmar F. The current deconstruction of paradoxes: one sign of the ongoing methodological "revolution". *Eur J Epidemiol*. 2015; 30(10):1079–1087. <https://doi.org/10.1007/s10654-015-0068-8> PMID: 26164615
44. Pearl J. *Causality: Models, Reasoning, and Inference*, 2nd ed. Cambridge Univ. Press, Cambridge. 2009.