

East Tennessee State University

## Digital Commons @ East Tennessee State University

---

ETSU Faculty Works

Faculty Works

---

10-29-2018

### Preliminary Scale of Reference Values for Evaluating Reactive Strength Index-Modified in Male and Female NCAA Division I Athletes

Christopher J. Sole

*The Citadel-The Military College of South Carolina*

Timothy J. Suchomel

*Carroll University*

Michael H. Stone

*East Tennessee State University, stonem@etsu.edu*

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>



Part of the [Exercise Physiology Commons](#), and the [Sports Sciences Commons](#)

---

#### Citation Information

Sole, Christopher J.; Suchomel, Timothy J.; and Stone, Michael H.. 2018. Preliminary Scale of Reference Values for Evaluating Reactive Strength Index-Modified in Male and Female NCAA Division I Athletes. *Sports*. Vol.6(4). 133. <https://doi.org/10.3390/sports6040133> ISSN: 2075-4663

This Article is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact [digilib@etsu.edu](mailto:digilib@etsu.edu).

---

## Preliminary Scale of Reference Values for Evaluating Reactive Strength Index-Modified in Male and Female NCAA Division I Athletes

### Copyright Statement

© 2018 by the authors. This document was originally published in *Sports*.

### Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Brief Report

# Preliminary Scale of Reference Values for Evaluating Reactive Strength Index-Modified in Male and Female NCAA Division I Athletes

Christopher J. Sole <sup>1,\*</sup>, Timothy J. Suchomel <sup>2</sup> and Michael H. Stone <sup>3</sup>

<sup>1</sup> Department of Health and Human Performance, The Citadel-The Military College of South Carolina, Charleston, SC 29409, USA

<sup>2</sup> Department of Human Movement Sciences, Carroll University, Waukesha, WI 53186, USA; tsuchome@carrollu.edu

<sup>3</sup> Center of Excellence for Sport Science and Coach Education, Department of Exercise and Sport Science, East Tennessee State University, Johnson City, TN 37614, USA; stonem@etsu.edu

\* Correspondence: csole@citadel.edu; Tel.: +1-843-953-6386

Received: 29 September 2018; Accepted: 19 October 2018; Published: 29 October 2018



**Abstract:** The purpose of this analysis was to construct a preliminary scale of reference values for reactive strength index-modified ( $RSI_{mod}$ ). Countermovement jump data from 151 National Collegiate Athletic Association (NCAA) Division I collegiate athletes (male  $n = 76$ ; female  $n = 75$ ) were analyzed. Using percentiles, scales for both male and female samples were constructed. For further analysis, athletes were separated into four performance groups based on  $RSI_{mod}$  and comparisons of jump height (JH), and time to takeoff (TTT) were performed.  $RSI_{mod}$  values ranged from 0.208 to 0.704 and 0.135 to 0.553 in males and females, respectively. Males had greater  $RSI_{mod}$  ( $p < 0.001$ ,  $d = 1.15$ ) and JH ( $p < 0.001$ ,  $d = 1.41$ ) as compared to females. No statistically significant difference was observed for TTT between males and females ( $p = 0.909$ ,  $d = 0.02$ ). Only JH was found to be statistically different between all performance groups. For TTT no statistical differences were observed when comparing the top two and middle two groups for males and top two, bottom two, and middle two groups for females. Similarities in TTT between sexes and across performance groups suggests JH is a primary factor contributing to differences in  $RSI_{mod}$ . The results of this analysis provide practitioners with additional insight as well as a scale of reference values for evaluating  $RSI_{mod}$  scores in collegiate athletes.

**Keywords:** countermovement jump; jump height; time to takeoff; force platform; athlete monitoring

## 1. Introduction

When assessing athlete testing and training data for the purpose of performance monitoring, coaches and sport scientists are often faced with questions regarding the worth of the data they have collected. Questions such as, “Is that score good?” or “How does that score compare to peer and aspirant performers?” are common as athlete performance data are reviewed. Unfortunately, answers to questions such as these are not always immediately apparent. In order to find answers, or determine the worth of data, coaches and sport scientists must engage in the process of evaluation.

Although the evaluation of data can be achieved through several means, one common approach is through a norm-referenced perspective. In this form of evaluation, an individual athlete’s score is compared to other scores considered representative of a population, and are commonly presented using percentiles. Overall, norm-referenced evaluation provides coaches and practitioners with a general idea of how an individual compares to a group or population. Normative data for various fitness and performance tests have been previously published for a variety of populations [1]. Depending on

the purpose of the testing, this method of evaluation can be useful, such as in talent identification, or grouping individuals based on their abilities. However, if data for a specific measure or population do not exist, this process is not possible.

With the growing interest in vertical jump testing in athlete performance monitoring [2], as well as the increased accessibility of technology such as portable force platforms [3–7], many new variables derived from a single vertical jump have arisen from the literature. Although the addition of new variables used to characterize jump performance is not necessarily negative, it does however present interpretation and evaluation challenges for practitioners. One such variable is reactive strength index-modified ( $RSI_{mod}$ ) [8]. Reactive strength index-modified can be calculated from a standing countermovement jump (CMJ) and represents the ratio of jump height (JH) to movement time, referred to as time to takeoff (TTT). Since it was first introduced [8],  $RSI_{mod}$  has been reported to be both a reliable [9] and valid [10,11] indication of the athlete's lower-body impulsive or "explosive" ability. Furthermore, considering  $RSI_{mod}$  takes into consideration an outcome variable (i.e., JH) and a process variable (i.e., TTT), it appears to be a simple and effective means for evaluating jumping strategy and ultimately neuromuscular functional state (i.e., adaptation or fatigue) [12].

Although a variety of normative data have been published related to CMJ's criterion variable of jump height [13–16], normative data have yet to be published on many specific CMJ variables such as  $RSI_{mod}$ . If  $RSI_{mod}$  is to be effectively used in athlete performance testing and monitoring, more information is needed related to the interpretation and evaluation of this variable. The purpose of this report is twofold (1) to present a preliminary scale of reference values for  $RSI_{mod}$ , and (2) to examine differences in the constituents of  $RSI_{mod}$  (JH and TTT) across the scale in an effort to provide additional context to aid in the evaluation of this variable.

## 2. Materials and Methods

### 2.1. Participants

This analysis was completed retrospectively using archived data that were collected as part of an ongoing athlete performance monitoring program [17]. Additionally, all athletes provided written informed consent at the time of data collection. The methods and scope of this analysis were reviewed and approved by the University's institutional review board. Countermovement jump data from a total of 151 collegiate athletes (male  $n = 76$ ; female  $n = 75$ ) were included in this analysis. All athletes were NCAA Division I, representing a variety of sports including baseball ( $n = 29$ ; height =  $182.1 \pm 6.2$  cm, body mass  $88.0 \pm 9.0$  kg), men's tennis ( $n = 7$ ; height =  $176.9 \pm 9.0$  cm, body mass =  $74.7 \pm 9.5$  kg), men's soccer ( $n = 25$ ; height =  $179.5 \pm 6.8$  cm, body mass =  $78.5 \pm 9.2$  kg), men's track and field ( $n = 15$ ; height =  $183.1 \pm 6.3$  cm; body mass =  $94.4 \pm 29.0$  kg), women's tennis ( $n = 11$ ; height =  $167.6 \pm 5.6$  cm, body mass =  $68.6 \pm 12.7$  kg), women's soccer ( $n = 22$ ; height =  $166.1 \pm 6.2$  cm, body mass =  $63.9 \pm 8.1$  kg), women's track and field ( $n = 13$ ; height =  $166.5 \pm 7.2$  cm, body mass =  $67.1 \pm 20.5$  kg), softball ( $n = 14$ ; height =  $168.5 \pm 6.7$  cm, body mass =  $70.4 \pm 10.1$  kg), and women's volleyball ( $n = 15$ ; height =  $175.3 \pm 7.5$  cm, body mass =  $70.7 \pm 7.7$  kg). Athletes ranged in age from 18–23 years.

### 2.2. Data Collection

In order to avoid any possible influence of fatigue from training or competition, all CMJ data included in this analysis were selected from the athlete's preseason testing sessions. All athletes were injury-free at the time of data collection. All data were collected under the same standardized testing protocol. Specifically, participants completed a general warm-up consisting of 25 jumping-jacks, one set of five repetitions of mid-thigh clean pulls with a 20 kg barbell, and three sets of five repetitions of mid-thigh clean pulls with barbell totaling 40 kg for females and 60 kg for males. Immediately following the general warm-up, participants began jump testing where they completed a specific warm-up consisting of two submaximal CMJs performed at 50% and 75% of their perceived maximum effort. Following the specific warm-up, participants performed two maximal CMJs with approximately

60 seconds of rest between each jump. Briefly, each participant stood motionless on the force platform and then received a countdown of “3, 2, 1, jump!” After the jump command, participants performed a rapid countermovement to a self-selected depth before propelling themselves upward with the intent of jumping as high as possible. To control for arm swing, all jumps were performed while holding a near-weightless ( $\leq 0.5$  kg) plastic bar across the shoulders, approximately between the seventh cervical and third thoracic vertebra [9,18]. All jumps were performed on a force platform (0.91 m  $\times$  0.91 m, RoughDeck HP, Rice Lake Weighing Systems, Rice Lake, WI, USA) sampling at 1000 Hz.

### 2.3. Data Analysis

Following data collection, voltage data obtained from the force platform were converted to the vertical component of the ground reaction force using laboratory calibrations. Force-time curves were then constructed. All data were collected and analyzed using custom programs (LabVIEW version 15, National Instruments, Austin, TX, USA). To reduce noise in the signal, a digital low-pass Butterworth filter with a cutoff frequency of 10 Hz was used. From the force-time data,  $RSI_{mod}$  was then calculated using procedures outlined by previous authors [8,9,19].

Equation (1)


$$RSI_{mod} = \frac{\text{CMJ height (m)}}{\text{time to takeoff (s)}} \quad (1)$$

The CMJ height (JH) was calculated from the vertical displacement of the jumper’s center of mass estimated from flight time [20] and TTT was defined as the time interval between the initiation of the countermovement and the instant when the jumper left the force platform. A threshold of 10 N was used to identify the beginning and end of this period.


### 2.4. Statistical Analyses

Relative and absolute reliability of  $RSI_{mod}$  and its constituent variables were assessed using a two-way mixed-effect model intraclass correlation coefficient (ICC) and typical error expressed as a coefficient of variation (CV%) performed between the two CMJ trials. Reliability was found to be acceptable with high test-retest correlation and low CV% for all variables for both males ( $RSI_{mod}$ : ICC = 0.963, CV% 7.6%; JH: ICC = 0.978, CV% = 4.7%; TTT: ICC 0.899, CV% = 5.4%) and females ( $RSI_{mod}$ : ICC = 0.967, CV% 7.9%, JH: ICC = 0.978, CV% = 4.1%; TTT: ICC 0.892, CV% = 6.0%), therefore the mean of the CMJ trials was used for all analyses [21,22].

Once averaged,  $RSI_{mod}$  data were aggregated by sex to form male (n = 76) and female (n = 75) groups.  $RSI_{mod}$  scores for both males and females were assessed and found to be normally distributed (males: Kolmogorov-Smirnov (D (76) = 0.090,  $p > 0.200$ ; Skewness: 0.170, SE = 0.276; Kurtosis:  $-0.154$ , SE = 0.545; females: Kolmogorov-Smirnov (D (75) = 0.064,  $p > 0.200$ ; Skewness: 0.221, SE = 0.277; Kurtosis:  $-0.272$ , SE = 0.548). Additionally, no outliers were identified [23,24] for either group. Male and female scales were then constructed using percentile rank. In order to allow for further analysis of the scales, male and female participants were then ranked based on  $RSI_{mod}$  scores and then divided into four performance groups using quartiles as cut points (Figures 1 and 2). Independent samples t-tests were used to compare differences in  $RSI_{mod}$ , JH, and TTT between males and females. To compare differences between each performance group, a series of one-way analysis of variances (ANOVAs) were used to examine the four performance groups with Bonferroni post hoc analysis used when appropriate. Levene’s test was used to assess equality of variance for all group comparisons. To provide an indication of the practical significance of any observed differences, Cohen’s  $d$  effect sizes were calculated and interpreted in accordance with the scale developed by Hopkins [25]. The critical alpha was set at  $p < 0.05$  for all analyses. All statistical analyses were performed using SPSS version 23 (IBM, Armonk, NY, USA) and Microsoft Excel 2013 (Microsoft Corporation, Redmond, WA, USA).



**Figure 1.** Ranked reactive strength index-modified (RSI<sub>mod</sub>) scores and performance groups for male athletes. Note: U = upper performance group; UM = upper-middle performance group; LM = lower-middle performance group; L = lower performance group.



**Figure 2.** Ranked reactive strength index-modified scores and performance groups for female athletes. Note: U = upper performance group; UM = upper-middle performance group; LM = lower-middle performance group; L = lower performance group.

### 3. Results

Descriptive statistics for RSI<sub>mod</sub> and its constituents for male and female athletes are displayed in Table 1. The preliminary reference value scales for RSI<sub>mod</sub> for both male and female athletes are displayed in Tables 2 and 3. Descriptive statistics for male and female athletes stratified by performance group are displayed in Table 4.

**Table 1.** Descriptive statistics for reactive strength index-modified for male and female athletes (n = 151, mean  $\pm$  standard deviation).

	Male (n = 76)	Female (n = 75)
RSI <sub>mod</sub> (m/s)	0.424 $\pm$ 0.102	0.314 $\pm$ 0.089
JH (m)	0.36 $\pm$ 0.07	0.27 $\pm$ 0.06
TTT (s)	0.868 $\pm$ 0.105	0.870 $\pm$ 0.114

Note: RSI<sub>mod</sub> = reactive strength index-modified; JH = jump height; TTT = time to takeoff

**Table 2.** Reference value scale for reactive strength index-modified for male collegiate athletes (n = 76).

Percentile	RSI <sub>mod</sub> (m/s)
97	0.630
95	0.604
90	0.547
85	0.523
80	0.508
75	0.492
70	0.487
65	0.476
60	0.461
55	0.448
50	0.419
45	0.398
40	0.376
35	0.369
30	0.366
25	0.352
20	0.331
15	0.316
10	0.308
5	0.257
3	0.216

**Table 3.** Reference value scale for reactive strength index-modified for female collegiate athletes (n = 75).

Percentile	RSI <sub>mod</sub> (m/s)
97	0.497
95	0.461
90	0.434
85	0.413
80	0.391
75	0.379
70	0.366
65	0.351
60	0.333
55	0.315
50	0.308
45	0.293
40	0.279
35	0.273
30	0.266
25	0.248
20	0.241
15	0.214
10	0.202
5	0.173
3	0.139

**Table 4.** Descriptive statistics for reactive strength index-modified between performance groups (mean  $\pm$  standard deviation).

Male Performance Group				Female Performance Group					
Performance Group	n	RSI <sub>mod</sub> (m/s)	JH (m)	TTT (s)	Performance Group	n	RSI <sub>mod</sub> (m/s)	JH (m)	TTT (s)
U	20	0.549 $\pm$ 0.057	0.43 $\pm$ 0.05	0.795 $\pm$ 0.087	U	19	0.429 $\pm$ 0.045	0.34 $\pm$ 0.03	0.792 $\pm$ 0.077
UM	18	0.464 $\pm$ 0.021	0.39 $\pm$ 0.04	0.850 $\pm$ 0.067	UM	18	0.343 $\pm$ 0.023	0.29 $\pm$ 0.04	0.848 $\pm$ 0.117
LM	20	0.379 $\pm$ 0.019	0.33 $\pm$ 0.04	0.871 $\pm$ 0.100	LM	20	0.277 $\pm$ 0.017	0.24 $\pm$ 0.03	0.880 $\pm$ 0.092
L	18	0.296 $\pm$ 0.044	0.28 $\pm$ 0.05	0.964 $\pm$ 0.090	L	18	0.203 $\pm$ 0.036	0.19 $\pm$ 0.03	0.964 $\pm$ 0.100

Note: RSI<sub>mod</sub> = reactive strength index-modified; JH = jump height; TTT = time to takeoff, U = upper performance group; UM = upper-middle performance group; LM = lower-middle performance group, L = lower performance group.

Statistically significant differences were observed when comparing male and female athletes with regard to RSI<sub>mod</sub> ( $t(149) = 7.09, p < 0.001, d = 1.15$ ) and JH ( $t(149) = 8.64, p < 0.001, d = 1.41$ ). No statistically significant difference was observed for TTT between male and female athletes ( $t(149) = 0.11, p = 0.909, d = 0.02$ ). Statistically significant differences were observed for JH and TTT between individual performance groups for both male (JH: ( $F_{(3,72)} = 43.68, p < 0.001$ ); TTT: ( $F_{(3,72)} = 12.17, p < 0.001$ ) and female athletes (JH: ( $F_{(3,71)} = 63.90, p < 0.001$ ); TTT: ( $F_{(3,71)} = 10.11, p < 0.001$ )). Specifically, post hoc analyses for the males revealed that JH was statistically different ( $p < 0.05$ ) between all four performance groups with moderate to very large effect sizes observed ranging from  $d = 0.86$  to  $2.96$ . For TTT all comparisons were found to be statistically significant with effect sizes ranging from moderate to large ( $d = 0.81$  to  $1.44$ ), with the exception of the comparisons of upper and upper-middle groups and upper-middle and lower-middle groups that were found not to be statistically different exhibiting moderate ( $d = 0.71$ ) and small ( $d = 0.25$ ) effect sizes, respectively. Similarly, for females, statistically significant ( $p < 0.05$ ) differences were identified between all four performance groups for JH with large to very large effect sizes observed ranging from  $d = 1.26$  to  $4.50$ . For TTT, all groups were found to be statistically different ( $p < 0.05$ ) with moderate to large effect sizes observed ranging from  $d = 1.05$  to  $1.93$  with the exception of comparisons of upper and upper middle ( $d = 0.57$ ), upper-middle and lower-middle ( $d = 0.31$ ), and lower-middle and lower ( $d = 0.87$ ) groups, who were determined not to be statistically different, exhibiting small to moderate effect sizes.

#### 4. Discussion

The purpose of this analysis was to construct a preliminary scale of reference values for evaluating RSI<sub>mod</sub>. Additionally, this report included an analysis of the variables used to calculate RSI<sub>mod</sub> in effort to improve the interpretation of this variable. RSI<sub>mod</sub> values ranged from  $0.208$  to  $0.704$  and  $0.135$  to  $0.553$  in males and females, respectively. When comparing male and female athletes overall, males exhibited statistically greater RSI<sub>mod</sub> values. This finding is in agreement with previous research that examined RSI<sub>mod</sub> differences between male and female athletes [9]. On average, male athletes were found to have RSI<sub>mod</sub> values 29.8% greater than their female counterparts. Interestingly, when comparing TTT values males and females were found to be quite similar, exhibiting only a 0.2% difference on average. This similarity in TTT between males and females is in agreement with previous investigations examining the temporal structure of the CMJ [26,27]. The observed similarities in TTT values and statistically different RSI<sub>mod</sub> values, indicates that the primary factor influencing the sex differences observed in RSI<sub>mod</sub> may be attributed to jump height. In fact, in the present analysis, there was an approximate 30% difference in JH between males and females. The strong influence of JH on RSI<sub>mod</sub> over TTT can be further illustrated by examining the relationships between these variables. The relationship between RSI<sub>mod</sub> and TTT was  $r = -0.60$  for females and  $r = -0.40$  for males, whereas the relationship between RSI<sub>mod</sub> and JH was  $r = 0.87$  and  $r = 0.89$  for males and females, respectively. Although not explicitly examined in the present study, there are several reasons why differences in JH may have existed between males and females. Two reasons may be due to differences in muscular strength or in jump strategy. Previous research has indicated that RSI<sub>mod</sub>



displayed strong relationships with maximal isometric strength [28]; however, it should be noted that additional research has indicated that  $RSI_{mod}$  differences existed between males and females despite controlling for strength level and using it as a covariate [29]. Furthermore, McMahon et al. [30], as well as Sole et al. [26], have indicated that male participants applied a larger concentric impulse and achieved a greater velocity throughout the concentric phase, which ultimately leads to a greater jump height. It is clear that although differences in  $RSI_{mod}$  and JH may exist between males and females, further research is needed to determine the factors that produce these differences.

A secondary analysis of the present data scale involved a comparison of four performance groups representing the upper, upper-middle, lower-middle, and lower male and female athletes, as determined by  $RSI_{mod}$ . When examining both JH and TTT between the performance groups, only JH was found to be statistically different between all groups. In contrast, no statistical differences and only small to moderate effect sizes were observed when comparing the top two (upper and upper-middle) and middle two (upper-middle and lower-middle) groups for males in TTT values. Similarly for females, no statistical differences and only small to moderate effect sizes were observed between the top two (upper and upper-middle), bottom two (lower and lower-middle), and middle two (upper- and lower-middle) groups. These results indicate that TTT alone was not enough to differentiate between groups, as was the case with JH. As mentioned above, the current findings may be partially explained by muscular strength differences between each group. An abundance of research supports the idea that stronger individuals produce superior jump performances compared to weaker individuals [31]. Another possible explanation for the current findings may be due to differences in musculotendinous stiffness characteristics. Kipp et al. [32] indicated that vertical stiffness was strongly correlated to the traditional measure of reactive strength index (drop jump height/ground contact time). Given that the  $RSI_{mod}$  and reactive strength index variants are strongly correlated [33], it is possible that vertical stiffness may also play a role in JH performance during a CMJ. To the authors' knowledge, no study has yet to investigate the influence of stiffness on  $RSI_{mod}$ .

The reference values presented in the current study were constructed using athletes from various sporting disciplines. While the use of multiple sports within a single scale may be viewed as a limitation, it should be noted that no other study has used a sample size as large as the one used within the current study. As displayed in Figures 1 and 2, the ranking and grouping of athletes by  $RSI_{mod}$  resulted in a disproportionate sport representation within groups. Although this lack of homogeneity within performance groups may be viewed as problematic, it also may simply highlight sport specific differences in jumping strategies as noted by previous reports [34]. Moreover, it should be noted that previous research indicated that within-team differences based on player position may exist when examining  $RSI_{mod}$  [35]. Thus, while the  $RSI_{mod}$  reference values produced in this study may serve as an initial step in  $RSI_{mod}$  comparisons between athletes, it is important for future research to continue to collect normative  $RSI_{mod}$  data for different sports and levels of sports so that additional scales may be developed. Furthermore, additional data will allow for the comparison of athletes within a single sport or between different levels of the same sport [1].

Although the scales and analyses provide insight for evaluating  $RSI_{mod}$  and its constituent parts, practitioners should be cognizant of some potential limitations of this data. The present scales and analyses used athlete data collected from a single university, and although all athletes were competitive at the NCAA Division I level, this sample may not be representative of all collegiate athletes. Thus, as noted above, it is important that if practitioners see value in using  $RSI_{mod}$  as a monitoring tool, additional data for male and female athletes is needed. A second limitation of the present study may be related to the specific data analysis procedures used to calculate  $RSI_{mod}$ . Although all data were collected using force plates sampling at an optimal sampling frequency [36] and analyzed using identical procedures, the present study used arbitrary thresholds to identify the initiation of the countermovement (unweighting phase) as well as takeoff and landing events. Furthermore, the jump height values used to calculate  $RSI_{mod}$  were estimated using time in air. Recently researchers have identified this as potentially problematic as it relates to  $RSI_{mod}$  as any errors in identifying TTT

or JH will ultimately influence  $RSI_{mod}$  [12]. Future analyses should consider using more robust methods [22,37] for identifying these key time points during the jump, in effort to reduce potential error. Although this limitation is valid, within this analysis it may be partially obviated by the fact that all CMJ analyses were completed using identical procedures.

## 5. Conclusions

The results of the present analysis provide practitioners with preliminary scales of reference values for interpreting  $RSI_{mod}$  scores in collegiate athletes. A primary finding of this report was that the difference in  $RSI_{mod}$  between male and female Division I athletes is largely attributed to differences in jump height. Interestingly, when compared as a whole, there were no discernable differences in TTT between male and female athletes. Analysis of the  $RSI_{mod}$  scales indicate that there are clear performance differences between upper performers and lower male and female performers, in both  $RSI_{mod}$  and its constituent parts. Given the importance of comparing performances between individuals, these reference values may provide a valuable resource for practitioners seeking to evaluate their athlete's performance.

**Author Contributions:** C.J.S. and T.J.S. conceived and designed the experiments; C.J.S. and T.J.S. performed the experiments; C.J.S. analyzed the data; C.J.S. and M.H.S. contributed materials and analysis tools; C.J.S., T.J.S., and M.H.S. wrote the paper.

**Funding:** No financial support was provided for the completion of this project.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Hoffman, J.R. *Norms for Fitness, Performance, and Health*; Human Kinetics: Champaign, IL, USA, 2006; ISBN-13: 9780736054836.
- McMahon, J.J.; Suchomel, T.J.; Lake, J.P.; Comfort, P. Understanding the key phases of the countermovement jump force-time curve. *Strength Cond. J.* **2018**, *40*, 96–106. [[CrossRef](#)]
- Linthorne, N. Analysis of standing vertical jumps using a force platform. *Am. J. Phys.* **2001**, *69*, 1198–1204. [[CrossRef](#)]
- Lake, J.; Mundy, P.; Comfort, P.; McMahon, J.J.; Suchomel, T.J.; Carden, P. Concurrent validity of a portable force plate using vertical jump force-time characteristics. *J. Appl. Biomech.* **2018**, *34*, 410–413. [[CrossRef](#)] [[PubMed](#)]
- Walsh, M.S.; Ford, K.R.; Bangen, K.J.; Myer, G.D.; Hewett, T.E. The validation of a portable force plate for measuring force-time data during jumping and landing tasks. *J. Strength Cond. Res.* **2006**, *20*, 730–734. [[PubMed](#)]
- Loturco, I.; Pereira, L.; Kobal, R.; Cal Abad, C.; Fernandes, V.; Ramirez-Campillo, R.; Suchomel, T.J. Portable force plates: A viable and practical alternative to rapidly and accurately monitor elite sprint performance. *Sports* **2018**, *6*, 61. [[CrossRef](#)] [[PubMed](#)]
- Peterson Silveira, R.; Stergiou, P.; Carpes, F.P.; Castro, F.A.d.S.; Katz, L.; Stefanyshyn, D.J. Validity of a portable force platform for assessing biomechanical parameters in three different tasks. *Sports Biomech.* **2017**, *16*, 177–186. [[CrossRef](#)] [[PubMed](#)]
- Ebben, W.P.; Petushek, E.J. Using the reactive strength index modified to evaluate plyometric performance. *J. Strength Cond. Res.* **2010**, *24*, 1983–1987. [[CrossRef](#)] [[PubMed](#)]
- Suchomel, T.J.; Bailey, C.A.; Sole, C.J.; Grazer, J.L.; Beckham, G.K. Using reactive strength index-modified as an explosive performance measurement tool in division I athletes. *J. Strength Cond. Res.* **2015**, *29*, 899–904. [[CrossRef](#)] [[PubMed](#)]
- Kipp, K.; Kiely, M.T.; Geiser, C.F. Reactive strength index modified is a valid measure of explosiveness in collegiate female volleyball players. *J. Strength Cond. Res.* **2016**, *30*, 1341–1347. [[CrossRef](#)] [[PubMed](#)]
- McMahon, J.J.; Jones, P.A.; Suchomel, T.J.; Lake, J.; Comfort, P. Influence of the reactive strength index modified on force-and power-time curves. *Int. J. Sports Physiol. Perform.* **2017**, *13*, 220–227. [[CrossRef](#)] [[PubMed](#)]

12. McMahon, J.J.; Lake, J.P.; Comfort, P. Reliability of and relationship between flight time to contraction time ratio and reactive strength index modified. *Sports* **2018**, *6*, 81. [CrossRef] [PubMed]
13. Nikolaidis, P. Age-related differences in countermovement vertical jump in soccer players 8–31 years old: The role of fat-free mass. *Am. J. Sports Sci. Med.* **2014**, *2*, 60–64.
14. Patterson, D.D.; Peterson, D.F. Vertical jump and leg power norms for young adults. *Meas. Phys. Educ. Exerc. Sci.* **2004**, *8*, 33–41. [CrossRef]
15. Castagna, C.; Castellini, E. Vertical jump performance in Italian male and female national team soccer players. *J. Strength Cond. Res.* **2013**, *27*, 1156–1161. [CrossRef] [PubMed]
16. Gabbett, T.J. Physiological characteristics of junior and senior rugby league players. *Br. J. Sport Med.* **2002**, *36*, 334–339. [CrossRef]
17. Sole, C.J.; Kavanaugh, A.A.; Reed, J.P.; Israetel, M.A.; Devine, L.E.; Ramsey, M.W.; Sands, W.A.; Stone, M.H. The sport performance enhancement group: A five-year analysis of interdisciplinary athlete development. In Proceedings of the 8th Annual Center of Excellence for Sport Science and Coach Education Coaches and Sport Science College, Johnson City, TN, USA, 13–14 December 2013.
18. Kraska, J.M.; Ramsey, M.W.; Haff, G.G.; Fethke, N.; Sands, W.A.; Stone, M.E.; Stone, M.H. Relationship between strength characteristics and unweighted and weighted vertical jump height. *Int. J. Sports Physiol. Perform.* **2009**, *4*, 461–473. [CrossRef] [PubMed]
19. Suchomel, T.J.; Sole, C.J.; Stone, M.H. Comparison of methods that assess lower-body stretch-shortening cycle utilization. *J. Strength Cond. Res.* **2016**, *30*, 547–554. [CrossRef] [PubMed]
20. Moir, G.L. Three different methods of calculating vertical jump height from force platform data in men and women. *Meas. Phys. Educ. Exerc. Sci.* **2008**, *12*, 207–218. [CrossRef]
21. Henry, F.M. “Best” versus “average” individual scores. *Res Q. Exerc. Sport* **1967**, *38*, 317–320. [CrossRef]
22. Kennedy, R.A.; Drake, D. Improving the signal-to-noise ratio when monitoring countermovement jump performance. *J. Strength Cond. Res.* **2018**, in press. [CrossRef] [PubMed]
23. Hoaglin, D.C.; Iglewicz, B. Fine-tuning some resistant rules for outlier labeling. *J. Am. Stat. Assoc.* **1987**, *82*, 1147–1149. [CrossRef]
24. Hoaglin, D.C.; Iglewicz, B.; Tukey, J.W. Performance of some resistant rules for outlier labeling. *J. Am. Stat. Assoc.* **1986**, *81*, 991–999. [CrossRef]
25. A New View of Statistics. Available online: <http://www.sportsci.org/resource/stats/effectmag.html> (accessed on 20 September 2018).
26. Sole, C.J.; Mizuguchi, S.; Sato, K.; Moir, G.L.; Stone, M.H. Phase characteristics of the countermovement jump force-time curve: A comparison of athletes by jumping ability. *J. Strength Cond. Res.* **2018**, *32*, 1155–1165. [CrossRef] [PubMed]
27. Laffaye, G.; Wagner, P.P.; Tomblason, T.I.L. Countermovement jump height: Gender and sport-specific differences in the force-time variables. *J. Strength Cond. Res.* **2014**, *28*, 1096–1105. [CrossRef] [PubMed]
28. Beckham, G.K.; Suchomel, T.J.; Bailey, C.A.; Sole, C.J.; Grazer, J.L. The relationship of the reactive strength index-modified and measures of force development in the isometric mid-thigh pull. In Proceedings of the XXXII International Conference of Biomechanics in Sports, Johnson City, TN, USA, 12–16 July 2014.
29. Beckham, G.K.; Suchomel, T.J.; Sole, C.J.; Bailey, C.A.; Grazer, J.L.; Kim, S.B.; Talbot, K.B.; Stone, M.H. Influence of sex and maximum strength on reactive strength index-modified. In Proceedings of the National Strength and Conditioning Association’s 41st Annual Meeting, Indianapolis, IN, USA, 11–14 July 2018.
30. McMahon, J.J.; Rej, S.J.; Comfort, P. Sex differences in countermovement jump phase characteristics. *Sports* **2017**, *5*, 8. [CrossRef] [PubMed]
31. Suchomel, T.J.; Nimphius, S.; Stone, M.H. The importance of muscular strength in athletic performance. *Sports Med.* **2016**, *46*, 1419–1449. [CrossRef] [PubMed]
32. Kipp, K.; Kiely, M.T.; Giordanelli, M.D.; Malloy, P.J.; Geiser, C.F. Biomechanical determinants of the reactive strength index during drop jumps. *Int. J. Sports Physiol. Perform.* **2018**, *13*, 44–49. [CrossRef] [PubMed]
33. McMahon, J.J.; Suchomel, T.J.; Lake, J.P.; Comfort, P. Relationship between reactive strength index variants in rugby league players. *J. Strength Cond Res.* **2018**, in press. [CrossRef] [PubMed]
34. Suchomel, T.J.; Sole, C.J.; Bailey, C.A.; Grazer, J.L.; Beckham, G.K. A comparison of reactive strength index-modified between six U.S. Collegiate athletic teams. *J. Strength Cond. Res.* **2015**, *29*, 1310–1316. [CrossRef] [PubMed]

35. Bailey, C.A.; Suchomel, T.J.; Beckham, G.K.; Sole, C.J.; Grazer, J.L. A comparison of baseball positional differences with reactive strength index-modified. In Proceedings of the XXXIInd International Conference of Biomechanics in Sports, Johnson City, TN, 12–16 July 2014.
36. Street, G.; McMillan, S.; Board, W.; Rasmussen, M.; Heneghan, J.M. Sources of error in determining countermovement jump height with the impulse method. *J. Appl. Biomech.* **2001**, *17*, 43–54. [[CrossRef](#)]
37. Owen, N.J.; Watkins, J.; Kilduff, L.P.; Bevan, H.R.; Bennett, M.A. Development of a criterion method to determine peak mechanical power output in a countermovement jump. *J. Strength Cond. Res.* **2014**, *28*, 1552–1558. [[CrossRef](#)] [[PubMed](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).