

East Tennessee State University

Digital Commons @ East Tennessee State University

ETSU Faculty Works

Faculty Works

1-1-2017

Identification of PP1 as the First Phosphatase for IRF7

Shunbin Ning

East Tennessee State University, nings1@etsu.edu

Ling Wang

East Tennessee State University, wangl3@etsu.edu

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>



Part of the [Internal Medicine Commons](#)

Citation Information

Ning, Shunbin; and Wang, Ling. 2017. Identification of PP1 as the First Phosphatase for IRF7. *Journal of Cell Signaling*. Vol.2(2). 146-146. <https://doi.org/10.18632/oncotarget.16385> ISSN: 2576-1471

This Article is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

Identification of PP1 as the First Phosphatase for IRF7

Copyright Statement

© 2017 Ning S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Creative Commons License



This work is licensed under a [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/).

Identification of PP1 as the First Phosphatase for IRF7

Shunbin Ning^{1,2}, Ling Wang^{1,2*}

¹Center of Excellence for Inflammation, Infectious Diseases and Immunity, Quillen College of Medicine, East Tennessee State University, Johnson City, USA

²Department of Internal Medicine, Quillen College of Medicine, East Tennessee State University, Johnson City, USA

*Corresponding author: Ling Wang, Center of Excellence for Inflammation, Infectious Diseases and Immunity, Quillen College of Medicine, East Tennessee State University, Johnson City, USA, Tel: 423-439-8063; E-mail: wangl3@etsu.edu

Received date: March 13, 2017; Accepted date: April 11, 2017; Published date: April 19, 2017

Copyright: © 2017 Ning S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Commentary

Interferon (IFN) regulatory factor 7 (IRF7) is phosphorylated and activated in response to pathogenic infections for production of type I IFNs. The IFN production has to be turned off soon after infection. While there are a panel of kinases have been identified for IRF7 phosphorylation, no phosphatase has been reported for IRF7 dephosphorylation that may play a pivotal role in turning off IFN production. We have recently addressed this critical question by identification of protein phosphatase 1 (PP1) as the first phosphatase for IRF7.

Main content: The host innate immune system defends against invading pathogens initially by triggering signaling pathways mediated by the transmembrane receptors TLRs [1], and cytoplasmic receptors that include RLRs [2,3], NLRs [4], cGAS [5,6], IFI16 [7], DDX41 [8,9], DHX9/36 [10,11], RNA polymerase III [12], TRIM5α [13], ISG56 [14], LRRFIP1 [15], MRE11 [16], amongst others. Interferon (IFN) regulatory factor 7 (IRF7) is phosphorylated and activated downstream of many of these innate immune pathways for induction of IFN-I gene expression (especially IFNαs) [17]. The innate immune system also comprises of lymphocytes-mediated epigenetic memory, which defends reinfection and involves ATF7-mediated chromatin regulation [18,19].

IRF7 is required not only for IFN priming at early stage, but also for IFN amplification at later stages when robust IFN-I production depends on a positive regulatory circuit between IRF7 and IFN-I [20-22]. This robust reaction is turned off soon after infection under normal physiological conditions, but excessive production of IFN-I is fatal to the cell. Thus, regulation of IRF7 phosphorylation is of paramount importance for controlling antiviral innate immunity. However, no phosphatase for negative regulation of IRF7 phosphorylation and activity has been reported.

In our recent study [23], we have identified a conserved protein phosphatase 1 (PP1)-binding motif in human and mouse IRF7 proteins, and shown that PP1 physically interacts with IRF7. Exogenous expression of PP1 subunits (PP1α, β or γ) ablates IKKε-stimulated IRF7 phosphorylation and dramatically attenuates IRF7 transcriptional activity. Inhibition of PP1 activity significantly increases IRF7 phosphorylation and IRF7-mediated IFNα production in response to NDV infection or Toll-like receptor 7 (TLR7) challenge, leading to impaired viral replication. In addition, IFN treatment, TLR challenges and viral infection induce PP1 expression.




Figure 1: PP1 abrogates IRF7-mediated type I IFN response in antiviral immunity. IRF7 is phosphorylated by IKKs or IRAK1 downstream of TLR signaling pathways in response to viral infection. PP1 targets IRF7 for dephosphorylation and attenuates type I IFN production in these signaling pathways. Also, PP1 promotes IL6 production. In addition, IFN treatment, TLR challenges and viral infection induce PP1 expression. Correspondingly, we have identified IFN-responsive elements and IRF-binding motifs in the PP1 gene promoters, and the transcriptional regulation is currently under investigated.

Our results are the first to identify PP1 as a phosphatase that targets key activating phosphorylation sites of IRF7, attenuating its activity and blocking the IFN-I response during viral infection (Figure 1). Thus, our study has addressed an important knowledge gap regarding IRF7-mediated IFN-I innate immune response, and has broad significance in IFN-mediated antiviral innate immunity and IRF7-mediated pathogenesis [17]. In future follow-up studies, we will validate our findings in *in vivo* systems, and develop strategies to control PP1 phosphatase activity during viral infection for potential clinical interventions.

References

1. Kawai T, Akira S (2010) The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat Immunol* 11: 373-384.
2. Rehwinkel J, Reis e Sousa C (2010) RIGorous detection: exposing virus through RNA sensing. *Science* 327: 284-286.
3. Hornung V, Latz E (2010) Intracellular DNA recognition. *Nat Rev Immunol* 10: 123-130.
4. Sha W, Mitoma H, Hanabuchi S, Bao M, Weng L, et al. (2014) Human NLRP3 inflammasome senses multiple types of bacterial RNAs. *Proc Natl Acad Sci U S A* 111: 16059-16064.
5. Sun L, Wu J, Du F, Chen X, Chen ZJ (2013) Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the Type I interferon pathway. *Science* 339: 786-91.
6. Wu J, Sun L, Chen X, Du F, Shi H, et al. (2013) Cyclic GMP-AMP is an endogenous second messenger in innate immune signaling by cytosolic DNA. *Science* 339: 826-830.
7. Unterholzner L, Keating SE, Baran M, Horan KA, Jensen SB, et al. (2010) IFI16 is an innate immune sensor for intracellular DNA. *Nat Immunol* 11: 997-1004.
8. Parvatiyar K, Zhang Z, Teles RM, Ouyang S, Jiang Y, et al. (2012) DDX41 recognizes the bacterial secondary messengers cyclic di-GMP and cyclic di-AMP to activate a type I interferon immune response. *Nat. Immunol* 13:1155-61.
9. Zhang Z, Yuan B, Bao M, Lu N, Kim T, et al. (2011) The helicase DDX41 senses intracellular DNA mediated by the adaptor STING in dendritic cells. *Nat Immunol* 12: 959-965.
10. Kim T, Pazhoor S, Bao M, Zhang Z, Hanabuchi S, et al. (2010) Aspartate-glutamate-alanine-histidine box motif (DEAH)/RNA helicase A helicases sense microbial DNA in human plasmacytoid dendritic cells. *Proc Natl Acad Sci USA* 107: 15181-6.
11. Zhang Z, Kim T, Bao M, Facchinetti V, Jung SY, et al. (2011) DDX1, DDX21, and DHX36 Helicases Form a Complex with the Adaptor Molecule TRIF to Sense dsRNA in Dendritic Cells. *Immunity* 34: 866-78.
12. Chiu YH, Macmillan JB, Chen ZJ (2009) RNA polymerase III detects cytosolic DNA and induces type I interferons through the RIG-I pathway. *Cell* 138: 576-591.
13. Pertel T, Hausmann S, Morger D, Züger S, Guerra J, et al. (2011) TRIM5 is an innate immune sensor for the retrovirus capsid lattice. *Nature* 472: 361-365.
14. Pichlmair A, Lassnig C, Eberle CA, Górna MW, Baumann CL, et al. (2011) IFIT1 is an antiviral protein that recognizes 5'-triphosphate RNA. *Nat Immunol* 12: 624-630.
15. Yang P, An H, Liu X, Wen M, Zheng Y, et al. (2010) The cytosolic nucleic acid sensor LRRFIP1 mediates the production of type I interferon via a beta-catenin-dependent pathway. *Nat Immunol* 11: 487-494.
16. Kondo T, Kobayashi J, Saitoh T, Maruyama K, Ishii KJ, et al. (2013) DNA damage sensor MRE11 recognizes cytosolic double-stranded DNA and induces type I interferon by regulating STING trafficking. *Proc Natl Acad Sci USA* 110: 2969-74.
17. Ning S, Pagano JS, Barber GN (2011) IRF7: activation, regulation, modification and function. *Genes Immun* 12: 399-414.
18. Yoshida K, Maekawa T, Zhu Y, Guillet CR, Chatton B, et al. (2015) The transcription factor ATF7 mediates lipopolysaccharide-induced epigenetic changes in macrophages involved in innate immunological memory. *Nat Immunol* 16: 1034-43.
19. Fang TC, Schaefer U, Mecklenbrauker I, Stienen A, Dewell S, et al. (2012) Histone H3 lysine 9 di-methylation as an epigenetic signature of the interferon response. *JEM* 209: 661-9.
20. Marie I, Durbin JE, Levy DE (1998) Differential viral induction of distinct interferon- α genes by positive feedback through interferon regulatory factor-7. *EMBO J* 17: 6660-9.
21. Sato M, Hata N, Asagiri M, Nakaya T, Taniguchi T, et al. (1998) Positive feedback regulation of type I IFN genes by the IFN-inducible transcription factor IRF-7. *FEBS Lett* 441: 106-110.
22. Honda K, Taniguchi T (2006) IRFs: master regulators of signalling by Toll-like receptors and cytosolic pattern-recognition receptors. *Nat Rev Immunol* 6: 644-58.
23. Wang L, Zhao J, Ren J, Hall KH, Moorman JP, et al. (2016) Protein phosphatase 1 abrogates IRF7-mediated type I IFN response in antiviral immunity. *Eur J Immunol* 46: 2409-2419.