

East Tennessee State University

Digital Commons @ East Tennessee State University

ETSU Faculty Works

Faculty Works

7-3-1905

Pediatric Telebehavioral Health in Rural Primary Care: An Electronically-Mediated Warm Handoff

J. A. Correll

East Tennessee State University

Jodi Polaha

East Tennessee State University, polaha@etsu.edu

Follow this and additional works at: <https://dc.etsu.edu/etsu-works>



Part of the [Family Medicine Commons](#)

Citation Information

Correll, J. A.; and Polaha, Jodi. 1905. Pediatric Telebehavioral Health in Rural Primary Care: An Electronically-Mediated Warm Handoff. *Collaborative Family Healthcare Association*, Philadelphia, PA. <https://slideplayer.com/slide/1690675/>

This Presentation is brought to you for free and open access by the Faculty Works at Digital Commons @ East Tennessee State University. It has been accepted for inclusion in ETSU Faculty Works by an authorized administrator of Digital Commons @ East Tennessee State University. For more information, please contact digilib@etsu.edu.

Pediatric Telebehavioral Health in Rural Primary Care: An Electronically-Mediated Warm Handoff

Pediatric Telebehavioral Health in Rural Primary Care: An Electronically-Mediated Warm Handoff

Jennifer A. Correll, M.A., LPC

Jodi Polaha, Ph.D.

East Tennessee State University

Outline

- Background
- Clinic description
- Year 1 outcomes
- Barriers
- Future directions



Background

- **21%** (Polaha, Dalton, & Allen, 2011)
- **63%** (Polaha, Dalton, & Allen)
- **Rural areas have less access to evidence-based care** (Hoagwood, et al., 2000; Palmer, et al, 2010).

Background

- Children with depression responded as well to CBT treatment delivered via videoconferencing as those receiving face-to-face treatment (Nelson, et al., 2003).
- ADHD was one of the most commonly treated child/adolescent diagnoses (Palmer, et al, 2010).
- Telehealth was identified as one of the most promising avenues of collaborative service delivery for rural and underserved youth with ADHD.

Southern Appalachian Telebehavioral Health Clinic

- Began August 2010
- 8 primary care systems across TN and school-based health clinics in NC.
- Distal clinics range from 45 miles to over 400 miles from ETSU.





Southern Appalachian Telebehavioral Health Clinic

- Initially offered specialty services
 - Co-located model



Southern Appalachian Telebehavioral Health Clinic

- Move towards more collaborative in November 2010.
 - 1 day of open hours per week
 - Continued to offer specialty care appointments
 - Referral increase!
- Even more collaborative in March 2011.
 - 2 days of open hours per week
 - Continued to offer specialty care appointments
 - HUGE referral increase!

Southern Appalachian Telebehavioral Health Clinic

- Moved completely away from a specialty care model in Year 2.
 - Providers utilize electronically-mediated warm hand offs to refer patients for services
 - Provide brief, problem focused interventions operating on a 15 minute hour.
 - Care coordination with provider

Southern Appalachian Telebehavioral Health Clinic

- Initial consult requires no paperwork and the patient is not charged.
- Patients placed on a sliding scale and complete a brief information packet if an additional appointment required.
- Addition of an ADHD management clinic
Summer 2011
 - Monitor behavioral response to medication
 - Provide behavioral recommendations to improve functioning at school/home

Southern Appalachian Telebehavioral Health Clinic: Clinic Data Year 1

- Descriptive and satisfaction data collected on 38 patients

	<i>Age (M)</i>	<i>SD</i>
Open Hours	8.9	3.6
Specialty Clinic	12.1	4.6
Overall	9.6	3.9

Southern Appalachian Telebehavioral Health Clinic: Clinic Data Year 1

- Gender
 - Female-47.4%
 - Male-52.6%
- Ethnicity
 - White-94.7%

Southern Appalachian Telebehavioral Health Clinic: Clinic Data Year 1

Referral Problem	Percentage
Noncompliant/Oppositional Behavior	41%
ADHD	15.4%
Toileting Concerns	12.8%
Depression	10.3%
Anxiety	7.7%

Southern Appalachian Telebehavioral Health Clinic: Clinic Data

Satisfaction Survey

Item	<i>M (SD)</i>
I could see the psychologist clearly during the telemedicine visit.	5.74 (.50)
I had no trouble hearing the psychologist when she spoke to me.	5.68 (.70)
I was able to speak freely with the psychologist and ask questions.	5.74 (.64)
The psychologist was able to ask me questions.	5.89 (.34)
The doctor seemed to understand my problem.	5.50 (.89)
I was comfortable with and understood what the psychologist told me about my complaint.	5.52 (.72)
The camera or other equipment embarrassed me or made me feel uncomfortable.	1.60 (1.19)
The telemedicine visit makes receiving care more accessible .	5.42 (.85)

Southern Appalachian Telebehavioral Health Clinic: Clinic Data

Item	<i>M (SD)</i>
I would prefer a telemedicine visit now rather than waiting for a face-to-face appointment with the same doctor.	4.89 (1.20)
I would have traveled to another city to see a specialist if I had not used telemedicine	4.87 (1.65)
Traveling to another hospital would have cut into my work/school or my child's school time.	5.00 (1.33)
Traveling would affect my wages for that time.	3.26 (2.17)
I would experience other inconveniences in traveling.	4.45 (2.02)
I would prefer a face-to-face visit with the specialist rather than a teleconsultation with a specialist.	3.42 (1.88)
This telemedicine visit was as good as a face-to-face encounter.	5.29 (.89)
Overall, I am satisfied with telemedicine.	5.47 (.65)

Southern Appalachian Telebehavioral Health Clinic: Clinic Data

- PSC was administered to all clients who completed a referral packet.
- Initial and follow-up PSC scores available for 6 patients.
 - Initial scores $M=25.5$, $SD=16.02$
 - Follow-up scores $M=17.3$, $SD=9.73$
 - Reduction in PSC scores for patients seen in specialty clinic.

Challenges

- Reduction in referrals over time
- Difficulty remaining top of mind for providers
- Record sharing
 - No access to clinics EMR
 - Rely on fax to exchange information
- Billing
 - Some patients reluctant to pay fee for service out of pocket
 - Insurance reimbursement challenges

Future directions

- Sustainability
 - Contracts with individual clinics
 - Billing infrastructure
- Increase focus on evidence based care of disruptive behavior via videoconferencing
- Collaborative care of disruptive behavior disorders such as ADHD
 - Serving as the link between primary care and local schools.

Conclusions

- Telebehavioral health can be an effective means of treatment for children and adolescents with behavioral concerns.
- Record sharing and reduction in top of mind presence serve as barriers to collaborative care.
- Addressing these barriers could improve collaboration.

Contact Information

Jodi Polaha, PhD.
polaha@etsu.edu

Jennifer Correll, M.A., LPC
correllj@etsu.edu

References

- Hoagwood, K., Kelleher, K.J., Feil, M., & Comer, D.M. (2000). Treatment services for children with ADHD: A national perspective. *Journal of the American Academy of Child and Adolescent Psychiatry*, 39, 198-206.
- Palmer, N.B., Myers, K.M., Stoep, A., McCarty, C.A., Geyer, J.R., & DeSalvo, A. (2010). Attention-Deficit/Hyperactivity Disorder and telemental health. *Curr Psychiatry Rep*, 15, 409-417. doi: 10.1007/s11920-010-0123-8.
- Polaha, J. Dalton, W.T., Allen, S., (2011). The prevalence of emotional and behavioral problems in pediatric primary care serving rural children. *Journal of Pediatric Psychology*, 36, 652-660. doi: 10.1093/jepsy/jsqn6.
- Nelson, E.L., Barnard, M., & Cain, S. (2003). Treating childhood depression over videoconferencing. *Telemed J E Health*, 9, 49-55.